

## **INVESTIGATIONS IN FISH CONTROL**

- 1. Laboratories and methods  
for screening fish-control chemicals**
- 2. Preliminary observations  
on the toxicity of antimycin A  
to fish and other aquatic animals**



*No. 186*

**United States Department of the Interior  
Fish and Wildlife Service  
Bureau of Sport Fisheries and Wildlife**

Bureau Circular 185

<u>Page</u>	<u>Column</u>	<u>Line</u>
8	1	39
11	2	3
13	1	47

Bureau Circular 186

1	1	16
2	1	46
2	1	48
3	Fig. 1	
5	2	17
6	2	2
8	1	2
10	2	23
10	2	28
17	1	--

ERRATA

<u>properties</u>	<u>Item</u>	<u>of properties</u>
<u>properties</u>	instead of	<u>porperties</u>
<u>or</u>	"	" <u>on</u>
<u>fiberglass</u>	"	" <u>fibergalss</u>
<u>are</u>	"	" <u>is</u>
<u>empirical</u>	"	" <u>emperical</u>
"	"	" "
<u>Methylethyl-</u> <u>acetic acid</u>	"	" <u>Methylethyl-</u> <u>actetic acid</u>
( <u>Table 15</u> )	"	" ( <u>Table 2</u> )
<u>were</u>	"	" <u>was</u>
<u>occurred</u>	"	" <u>occured</u>
<u>12°</u>	"	" <u>12</u>
<u>17°</u>	"	" <u>17</u>
Literature citation No. 5: add <u>Biochim. Biophys. Acta., Vol. 53</u>		

# **INVESTIGATIONS IN FISH CONTROL**

## **1. Laboratories and methods for screening fish-control chemicals**

**Bureau Circular 185**

## **2. Preliminary observations on the toxicity of antimycin A to fish and other aquatic animals**

**Bureau Circular 186**



Washington, D.C. • June 1964

Investigations in Fish Control are reports on the results of work at the Bureau's Fish Control Laboratories at La Crosse, Wis., and Warm Springs, Ga. The two reports presented here are the first of several reports that are planned on work now under way.

## CONTENTS

	Pages
Laboratories and methods for screening fish-control chemicals, by Robert E. Lennon and Charles R. Walker . . . . .	1-15
Preliminary observations on the toxicity of antimycin A to fish and other aquatic animals, by Charles R. Walker, Robert E. Lennon, and Bernard L. Berger . . . . .	1-18



# INVESTIGATIONS IN FISH CONTROL

## 1. Laboratories and methods for screening fish-control chemicals

By Robert E. Lennon, Fishery Research Biologist  
Charles R. Walker, Chemist  
Bureau of Sport Fisheries and Wildlife  
La Crosse, Wis.



Bureau of Sport Fisheries and Wildlife

Circular 185

Washington, D.C. • June 1964

## CONTENTS

	Page
Abstract . . . . .	1
Facilities . . . . .	3
Bioassay laboratories . . . . .	3
Other laboratory facilities . . . . .	4
Fish-holding facilities . . . . .	5
Outdoor bioassay pools . . . . .	5
Methods . . . . .	6
Test chemicals . . . . .	8
Test fishes . . . . .	8
Responses of test fish . . . . .	10
Screening . . . . .	12
Recording and reporting results . . . . .	14
Summary . . . . .	14
Literature cited . . . . .	15



## Laboratories and methods for screening fish-control chemicals

By Robert E. Lennon, Fishery Research Biologist  
and Charles R. Walker, Chemist  
Bureau of Sport Fisheries and Wildlife  
La Crosse, Wis.

**Abstract.**--This report describes the physical and technical facilities and the procedures of the Fish Control Laboratories at La Crosse, Wis., and Warm Springs, Ga. The laboratories emphasize screening of chemicals to find a variety of fishery management tools. Preliminary Screening ascertains whether a chemical in three concentrations has a desirable biological activity on eight species of fish in reconstituted water at 12° and 17° C. Delineative Screening ascertains effective concentrations (EC100) on eight species in reconstituted water (the method for which is described) at 12°, 17°, 22°, and 27°. Intensive Screening of promising fish-control agents ascertains effects on 24 species of fish and on other aquatic organisms, at different temperatures and in waters of various qualities, in the laboratory and in the field.

Fish Control Laboratories were established by the Bureau of Sport Fisheries and Wildlife at La Crosse, Wis., in 1959 and at Warm Springs, Ga., in 1963. The mission of the Laboratories is the development of means for efficient manipulation of fresh-water fish. In particular, safe and economical controls--chemical, biological, electrical, or mechanical--are sought for undesirable populations in standing and flowing waters. The objectives are sufficiently broad to encompass investigation and development of any new tools that may be useful in fishery management, fish culture, or fishery research.

At La Crosse, the buildings of the National Fish Hatchery were remodeled and expanded in 1960-62 to provide a large research facility (figs. 1 and 2). The subsidiary Laboratory at Warm Springs is new construction on the grounds of the National Fish Hatchery (fig. 3). Their locations offer contrasting advantages to be exploited through close coordination in the research on fish control:

### La Crosse

Northern fishes.  
Cold climate.  
Cold water.  
Hard water.

### Warm Springs

Southern fishes.  
Warm climate.  
Warm water.  
Soft water.

In equipping and staffing the laboratories, early recognition was given to the potentials of chemical control agents. The bioassay (wet), chemistry, and physiology laboratories are concerned with general and selective toxicants, attractants, repellants, anesthetics, sterilants, spawning inducers, osmoregulators, marking dyes, medications for diseases, and sedatives and decontaminants for fish distribution. Emphasis is on finding selective toxicants for longnose and shortnose gars, gizzard shad, goldfish, carp, squawfishes, white sucker, black bullhead, rock bass, green sunfish, pumpkinseed, yellow perch, and freshwater drum.

Ample justification for research on selective piscicides is contained in fishery literature.




Figure 1.--The Fish Control Laboratory at La Crosse, Wis.



Figure 2.--The fish holding house at La Crosse, Wis.



Figure 3.--The Fish Control Laboratory (foreground) and wet laboratory holding house at Warm Springs, Ga. Construction and grading were incomplete when the photograph was taken.

LesVeaux (1959) listed 30 States that need control of certain troublesome fishes. Among the qualities desired in selective toxicants are specificity to certain life stages or to certain fish, low cost, ease and safety of application, rapid degradation to nontoxic residues, harmlessness to warm-blooded animals, and effectiveness at low temperatures. Applegate et al. (1961) reported on an effective and selective sea lamprey larvicide. Loosanoff, MacKenzie, and Shearer (1960) showed the possibilities of controlling certain shellfish predators with chemicals in marine environments. These and other studies stimulated interest in investigations to find selective toxicants for various fresh-water fish. The American Fisheries Society, for example, resolved at its 88th annual meeting in 1958 to recommend, to the Secretary of the Interior, an expansion of research in fish control. Congress in the same year made the first appropriation for establishment of the Fish Control Laboratory at La Crosse.

## FACILITIES

### Bioassay laboratories

At La Crosse and Warm Springs there are wet laboratories for large-scale screening of

chemicals against fish. Fiberglass or aluminum troughs serve as water baths for bioassay vessels, and fiberglass or concrete tanks hold selected fish for experiments (figs. 4 and 5). Ground water is used for the water baths and fish holding, and temperatures are adjusted by means of thermostatically controlled immersion heaters or refrigeration units.

Deionized water of at least 1 million ohms resistivity is reconstituted according to a formula developed at the Bureau's Fish-Pesticide Research Laboratory and is employed as a test medium in the bioassay vessels. The following chemicals are added per liter of deionized water: 30 mg. of calcium sulfate, 30 mg. of magnesium sulfate, 48 mg. of sodium bicarbonate, and 3 mg. of potassium chloride.

Glass vessels are preferred for bioassays in the laboratory. Most of the screening is done in economical 1-gallon pickle jars which are used once and discarded. Five-gallon water buckets of glass are employed in advanced screening. They are reused after thorough decontamination and washing which include several steps:

- a. Rinse jar with tap water.






Figure 4.--View of wet laboratory at La Crosse showing batteries of bioassay vessels in concrete, aluminum, and fiberglass tanks.

- b. Add 6.3 grams of activated charcoal (1 gram/3 liters), fill jar with deionized water, and let stand over night.
- c. Empty and rinse; wash with strong detergent in hot tap water and rinse thoroughly.
- d. Sponge entire jar with 10- to 14-percent hydrochloric acid and rinse twice with deionized water.

Whenever residual contamination is detected or suspected, the jar is discarded. All discarded bioassay glassware is smashed to prevent further use.

All test solutions are discarded into a floor drain which continually carries at least 300 g.p.m. of waste water from the fish holding tanks. The dilution has been found to be more than sufficient to eliminate hazards.

All fish used in the bioassays are disposed of in gas-fired incinerators of complete-combustion type.

#### Other laboratory facilities

Each Fish Control Laboratory has chemistry, biochemistry, and physiology laboratories as adjuncts to the bioassay facilities. Chemicals for testing are received in the chemistry laboratories, stored in fire- and explosion-resistant vaults, and prepared in proper solutions and dilutions for bioassay. Compounds showing promise as fish-control agents are investigated in the biochemistry laboratories to evolve methods for application, effective and economical formulations, possibilities for potentiation, means for minimizing side effects and hazards, and techniques for detoxification. In the final stages of development, a control agent is studied in the



Figure 5.--View of a wet laboratory at Warm Springs showing a battery of bioassay vessels in a fiberglass tank,

physiology laboratories to define its mode of action on fishes and other organisms, any chronic effects, the fate of residues in live animals, and the risks, if any, to consumers of treated fish.

#### Fish-holding facilities

Large quantities of fish are required for the chemical screening programs. At La Crosse, for example, 498,000 fish of 34 species were used in 1963. Most are obtained from Federal, State, or private hatcheries and rearing stations; it is more satisfactory to arrange small, frequent deliveries than attempt to maintain large quantities on hand

for long periods in usable, disease-free condition. A holding house and outside pools are provided at each Laboratory for the maintenance, feeding, sorting, and grading of the experimental fish (fig. 3 and 6).

#### Outdoor bioassay pools

An intermediate step between laboratory testing and field trials of promising fish-control agents is essential to detect and evaluate some of the physical or chemical factors that influence the performance of a candidate agent in natural waters. Raceways and portable plastic pools are located at each Laboratory for this purpose.





Figure 6.--Interior view of fish holding house at La Crosse.

Inexpensive vinyl wading pools, 9 and 10 feet in diameter, 2.5 feet deep, and about 1,000 gallons in capacity, are set up as described by Lawrence and Blackburn (1962) in outdoor testing areas (figs. 7 and 8). Bottom soils of various types, pond or ground waters, aquatic plants, invertebrates, fish, and amphibians are used in them as needed during chemical trials. Contaminated vinyl liners are economically replaced. The pools at La Crosse are employed only during the warm season, but those at Warm Springs are in operation all year.

Ten- and 20-foot concrete raceways are used for extraordinary tests in running or standing water. Disposable vinyl liners are used when necessary to avoid harmful contamination of the raceways.

## METHODS

The static bioassay is the first approach in screening chemicals for control agents. Constant-flow bioassays are reserved for advanced stages of testing.

Bliss (1957) defined a bioassay as a determination of the potency of a physical, chemical, or biological agent by means of a biological indicator. Noting its development during the past 20 or 30 years by scientists from many and diverse fields, he listed principles which characterize the modern bioassay: (1) Potency is a property of the drug, not of the response; (2) potency is relative, not absolute; (3) the assayed potency of an unknown is only an estimate of its true value; and (4) both the reliability



Figure 7.--Vinyl bioassay pools on levee at La Crosse.



Figure 8.--Vinyl bioassay pools at Warm Springs.

and efficiency of an assay are linked inseparably with its design. Observance of these principles overcomes some major disadvantages of the bioassay as a research tool.

Fish have long been employed as biological indicators in bioassays of water pollutants, insecticides, herbicides, detergents, and other substances. The standards recommended by Doudoroff et al. (1951), Henderson and Tarzwell (1957), and Henderson (1960) for such tests have been widely accepted and applied, although difficulties in comparing studies have arisen because of the many kinds of fish involved. Douglas and Irwin (1962) pointed out that the results of independent bioassays often cannot be related because the comparative resistance of the many test fishes has not been established. They noted that certain species have been more useful in toxicity bioassays than others, and they emphasized the need for knowledge about the reactions of different species of fish when exposed to a particular toxicant.

The large body of literature on methods and results of bioassays with fish has been helpful in defining the screening programs of the Fish Control Laboratories. In general, the practical methods proposed by the investigators cited above are followed but with some modifications since we seek more in biological activities than acute toxicity only.

### Test chemicals

The test chemicals are selected by staff chemists and biologists and are contributed by industry. Preference is given to compounds that have demonstrated biological activity or are suspected of possessing a useful activity against fish. It is also desirable to have as much information as possible before screening on the nature and properties of each chemical, its shelf life and stability, its solubility, and its potential hazards to investigators. Security is respected, and precautions are taken with compounds and test data to protect the rights of contributors.

The chemicals obtained for screening are arbitrarily classified as follows in order to

facilitate scheduling of tests, observation of responses, and reporting of results:

1. Natural organic products:
  - a. Animal extracts (steroids, proteins, etc.).
  - b. Plant extracts (rotenoids, alkaloids, phenols, etc.).
  - c. Fermentation products (antibiotics, etc.).
2. Synthetic organic products:
  - a. Halogenated hydrocarbons.
  - b. Nitrogen-bearing hydrocarbons (salicylanilides, carbamates, triazines, etc.).
  - c. Phosphorus-bearing hydrocarbons.
  - d. Sulfur-bearing hydrocarbons (mercapto, thioates, thiozines, etc.).
  - e. Miscellaneous compounds and combinations.
3. Inorganic products.

The progress of a chemical through testing and development is depicted in figure 9. A compound is introduced into Preliminary Screening to detect whether it has activity against fish. If it has, it is advanced into Delineative Screening, where the effective concentrations are defined and the possible usefulness of the substance is declared. If results are favorable, the candidate is referred to Intensive Screening, where it is fully evaluated in the laboratory and in the field as a fishery tool. Compounds which fail to meet requirements at any stage of screening are shelved or discarded immediately.

### Test fishes

It is of utmost importance that the validity and comparability of the bioassays be assured by using only fish of selected species, of certain sizes, and in good condition as biological indicators. The criteria are defined and strictly observed for each stage of screening. Therefore, the rate of progress of a chemical through screening and development depends largely on the availability of the prescribed fishes.




Figure 9.-Schematic on the progress of chemicals through screening.



TABLE 1.--Time required by certain fish held at 12° C. to empty the digestive tracts after food is withheld

Species	Number of fish		Voiding time (hours)
	Tested	Per pound	
Rainbow trout.....	90	2,000	36
Do.....	30	380	60
Do.....	12	22	84
Goldfish.....	30	207	48
River shiner.....	30	226	36
White sucker.....	12	76	60
Green sunfish.....	45	355	60
Pumpkinseed.....	45	177	84
Bluegill.....	60	368	84
Longear sunfish.....	45	465	72

Lots of hatchery fish are requested for delivery at least 2 weeks prior to use in bioassays. They are placed in the care of fish culturists, and everything possible is done to minimize stresses. During the first 10 days they are fed, prophylactically or therapeutically treated as necessary, and observed to evaluate them as test animals. Lots in which the mortalities exceed 10 percent within the period are not moved into the bioassay program.

The fish for experiments are carefully graded to desired and uniform size and transferred into the wet laboratories or outside pools 3 or 4 days before use. Food is withheld for as long as 96 hours before screening, depending on the life stage and species of fish. Generally, young fish and certain species require less time to empty the intestinal tract than others (table 1). The fish spend at least 24 hours in water similar to the test medium before being introduced into the bioassay vessels.

A help in maintaining comparable results in bioassays is use of a recognized reference toxicant against a sample of fish from each test lot. We employ para-, para'-DDT (1,1,1-trichloro-2,2-bis(p-chlorophenyl) ethane), and the test is made coincidentally with the introduction of a lot of fish into the screening program; the test is repeated biweekly if the lot remains on hand. The sensitivity (EC50) of the sample is determined for comparison with regression curves established for the species from experience or literature. Although costly, the test provides the only measure of relative sensitivity of fish used in the bioassays.

The value of reference tests was demonstrated during shakedown trials of facilities

and methods at La Crosse. For example, goldfish from Missouri and Wisconsin had such widely divergent sensitivities that they bracketed those of other species. Also, the sensitivities within a lot of fish differ significantly if some specimens are exposed to presumably harmless electric shocks like those experienced in electrofishing. Moreover, the stress of disease, malnutrition, temperature change, or altered water quality may influence sensitivity to a toxicant. Holding time is another factor, and we observed that the sensitivities changed greatly in a lot of goldfish which was retained for 2 months. Saila (1953) emphasized the significance of this and showed that the resistance of his mosquito-fish to rotenone decreased in rough proportion to the length of time they were held.

#### Responses of test fish

We recognize that the observer may be the greatest source of error in bioassays with fish. The criteria of response are arbitrary and subject to individual interpretation. Moreover, they are complicated at the Fish Control Laboratories because effects in addition to and more subtle than acute toxicity are sought. To achieve individual and mutual consistency, the observers are trained to use definite criteria of response.

Chemicals may have either short-term or long-term effects on fishes. Those of short term are typically identified with the following:

1. Acute toxicity:
  - a. Selective toxicant.
  - b. General toxicant.
2. Movement:
  - a. Attractant.
  - b. Repellant.
3. Facilitating capture, handling, or transport:
  - a. Anesthetic.
  - b. Sedative.
  - c. Osmoregulator.
4. Marking live fish:
  - a. Immersion stain.
  - b. Internal stain.

5. Therapy or prevention of disease:
  - a. Bactericide.
  - b. Parasiticide.
  - c. Fungicide.
  - d. Prophylactic.
  - e. Antiseptic.

- c. Inverted.
- d. On side.
- e. Against tank sides on bottom.

The long-term activities of potential control chemicals may be difficult to detect and evaluate. They are associated with--

1. Control of reproduction:
  - a. Hormonal spawning inducer.
  - b. Hormonal spawning inhibitor.
  - c. Sterilant.
2. Control of growth and development:
  - a. Growth stimulant.
  - b. Growth inhibitor.

The criteria employed to evaluate the responses of fish are rather refined, and are applicable to short-term responses. They are used as necessary at each observation period to identify fully the reactions of the fish.

They are--

#### General behaviour:

1. No observable difference from control.
2. Quiescent.
3. Excitable.
4. Irritated.
5. Surfacing.
6. Sounding.
7. Twitching.
8. Motionless:
  - a. Tetany.
  - b. Flaccidity.
9. Swimming:
  - a. Erratic; convulsive.
  - b. Gyration; skittering.

#### Integument:

1. Pigmentation:
  - a. No observable change.
  - b. Light discoloration.
  - c. Dark discoloration.
  - d. Varidiscoloration.
2. External mucosa:
  - a. No observable change.
  - b. Shedding; patchy.
  - c. Copious exudate.
  - d. Coagulation.
3. Hemorrhagic.

#### Respiration:

1. Respiratory rate:
  - a. No observable change.
  - b. Rapid.
  - c. Slow.
  - d. Irregular.
  - e. Ceased.
2. Gulping air.
3. Structures, organs:
  - a. No observable change.
  - b. Mouth gaping.
  - c. Hemorrhage in gills.
  - d. Irritative response.
  - e. Copious mucus in gills.

#### Alimentary responses:

1. No observable change.
2. Egurgitating mucus or other material.
3. Defecating mucus or other material.

#### Nervous responses:

1. No observable change.

2. Sensitivity to stimuli:

	<u>Positive</u>	<u>Negative</u>
a. Exterior movement..	_____	_____
b. Light.....	_____	_____
c. Sound.....	_____	_____
d. Touch.....	_____	_____
e. Electric probe .....	_____	_____

Moribundity:

1. No motion.
2. No respiration.
3. Distended operculum.
4. Opaque eyes.
5. Death.

Recovery:

1. Complete.
2. Incomplete.

Additional and more definitive observations are made of fish at the in-test, recovery, or postmortem stages in the biochemistry and physiology laboratories. Their objectives may be, for example, the modes of action and side effects of control agents.

Screening

Preliminary Screening.--Preliminary Screening is designed to detect whether a selected chemical at 0.1, 1, and 10 p.p.m. has a biological activity (see list of criteria above) against eight species of fish within 48 hours. It is a static bioassay in 1-gallon jars containing 2.5 liters of pre-aerated, reconstituted water. No artificial aerating is done during the test. The temperatures of water

baths are 12° C. at La Crosse and 17° C. at Warm Springs.

The fish include the following species:

<u>La Crosse</u>	<u>Warm Springs</u>
Rainbow trout.	Carp.
Goldfish.	Black bullhead.
White sucker.	Green sunfish.
Yellow perch.	Bluegill.

Goldfish from the same gene pool and from the same source are employed for occasional toxicity checks between the two laboratories.

At least 10 fish of each species are used with each concentration of chemical and as controls. They weigh between half a gram and 2 grams each, but the weight range of the fish in each test does not exceed 15 percent. They are distributed at one species per jar about 16 hours before the bioassay at a loading of 1 gram or less of fish per liter of test medium. Thus, there may be but one or two fish per jar, and up to 10 jars may be needed for each concentration of chemical.

A stock solution of a test compound is prepared within an hour of the bioassay. The solvents preferred are water, acetone, and ethanol, in that order. Because of their toxicity to fish, precautions are taken that the concentrations of acetone and ethanol do not exceed 4 parts per thousand in the bioassay media. Furthermore, whenever these solvents are used, equal volumes of them are applied in control vessels. Thus the volume in controls is equal to the highest volume used in a dosage series. Small aliquots of a stock solution are thoroughly mixed into bioassay media to avoid improper dilution or stratification.

The responses of the fish in bioassays and controls are observed routinely at 0.75, 1.5, 3, 6, 24, and 48 hours, but so far as possible they are recorded on the first day and thereafter as often as the nature of a candidate compound warrants.

If a compound has a desirable activity against the fish, it is held for further screenings, which are discussed briefly below.



**Delineative Screening.**--The effective concentrations (EC100) of a chemical against fish are determined in Delineative Screening. The vessels may be jars or troughs for static or flowing water trials. The water, aeration, species and size of fish, duration of tests, and controls are the same as in Preliminary Screening. A compound is first bioassayed at 12° C. to define concentrations which evoke all-or-none responses from the fish. The approach to these concentrations may be direct by bracketing and interpolation, or it may be indirect by probit analysis, whichever is the shorter. The EC100, if seemingly practical, is confirmed by seven replicate trials.

A chemical which yields favorable results at 12° C. is retested at 17°, 22°, and 27° to evaluate the effects of temperature on the effective concentrations. At 22° and 27°, rainbow trout are omitted from the bioassays, and the loading rates for other species are half a gram or less per liter of test medium.

If a compound succeeds in these trials, its potential as a fish-control agent is estimated. Information is sought on its possible application; on the source and manufacturing costs; on possible hazards, conflicts, or limitations in use; and on the size of the market. Only if it continues to appear promising is the chemical promoted into the more elaborate and expensive Intensive Screening.

**Intensive Screening.**--Intensive Screening is directed toward development of a promising chemical as a fish-control agent and the definition of its advantages and limitations. These are broad objectives, and some subdivision of approaches is in order.

**A. Stages:** The advanced tests of a compound are accomplished in the wet laboratories, in outside pools, and finally in the field. They may include static or flowing water bioassays with durations of 24, 48, and 96 hours or longer. The vessels may include 1-gallon jars with 2.5 liters of test medium, 5-gallon jars with 15 liters of medium, and fiberglass troughs of 4-, 8-, and 16-foot lengths. The outdoor pools include 1,000-gallon vinyl units and 10- and 20-foot concrete

raceways. Proportioning pumps are used in the flowing bioassays to achieve consistent concentrations of chemical.

The initial trials of a compound in the field are conducted in waters closed to public use. These waters are found, for example, on the grounds of Federal and State fish hatcheries, on wildlife refuges, or on military reservations. Trials are manned or immediately supervised by the Laboratory staff. Subsequent experiments in the field may be accomplished by selected cooperators in State and Federal agencies.

**B. Varieties of fish:** Sixteen species in addition to those included in Preliminary and Delineative Screening are employed. They are:

1. Brook trout.
2. Northern pike.
3. Fathead minnow.
4. Brook stickleback.
5. Pumpkinseed.
6. Longear sunfish.
7. Smallmouth bass.
8. Walleye.
9. Gizzard shad.
10. Golden shiner.
11. Bigmouth buffalo.
12. Brown bullhead.
13. Channel catfish.
14. Redear sunfish.
15. Largemouth bass.
16. White crappie.

Other species may be used on occasion in advanced tests.

**C. Life stages:** Various life stages of fish, from egg to mature adult, may be involved in the more advanced bioassays to detect the presence, absence, or variation of response to a promising chemical. It may be a great advantage or disadvantage if a potential control agent is selective for one life stage and not for another.

**D. Other organisms:** The effects of a potential chemical tool on other aquatic life must be assessed at least minimally in the laboratory before it is tested in the field. Some forms such as water fleas, snails, and plants

2. Sensitivity to stimuli:

	<u>Positive</u>	<u>Negative</u>
a. Exterior movement..	_____	_____
b. Light.....	_____	_____
c. Sound.....	_____	_____
d. Touch.....	_____	_____
e. Electric probe .....	_____	_____

Moribundity:

1. No motion.
2. No respiration.
3. Distended operculum.
4. Opaque eyes.
5. Death.

Recovery:

1. Complete.
2. Incomplete.

Additional and more definitive observations are made of fish at the in-test, recovery, or postmortem stages in the biochemistry and physiology laboratories. Their objectives may be, for example, the modes of action and side effects of control agents.

Screening

Preliminary Screening.--Preliminary Screening is designed to detect whether a selected chemical at 0.1, 1, and 10 p.p.m. has a biological activity (see list of criteria above) against eight species of fish within 48 hours. It is a static bioassay in 1-gallon jars containing 2.5 liters of pre-aerated, reconstituted water. No artificial aerating is done during the test. The temperatures of water

baths are 12° C. at La Crosse and 17° C. at Warm Springs.

The fish include the following species:

<u>La Crosse</u>	<u>Warm Springs</u>
Rainbow trout.	Carp.
Goldfish.	Black bullhead.
White sucker.	Green sunfish.
Yellow perch.	Bluegill.

Goldfish from the same gene pool and from the same source are employed for occasional toxicity checks between the two laboratories.

At least 10 fish of each species are used with each concentration of chemical and as controls. They weigh between half a gram and 2 grams each, but the weight range of the fish in each test does not exceed 15 percent. They are distributed at one species per jar about 16 hours before the bioassay at a loading of 1 gram or less of fish per liter of test medium. Thus, there may be but one or two fish per jar, and up to 10 jars may be needed for each concentration of chemical.

A stock solution of a test compound is prepared within an hour of the bioassay. The solvents preferred are water, acetone, and ethanol, in that order. Because of their toxicity to fish, precautions are taken that the concentrations of acetone and ethanol do not exceed 4 parts per thousand in the bioassay media. Furthermore, whenever these solvents are used, equal volumes of them are applied in control vessels. Thus the volume in controls is equal to the highest volume used in a dosage series. Small aliquots of a stock solution are thoroughly mixed into bioassay media to avoid improper dilution or stratification.

The responses of the fish in bioassays and controls are observed routinely at 0.75, 1.5, 3, 6, 24, and 48 hours, but so far as possible they are recorded on the first day and thereafter as often as the nature of a candidate compound warrants.

If a compound has a desirable activity against the fish, it is held for further screenings, which are discussed briefly below.

**Delineative Screening.**--The effective concentrations (EC100) of a chemical against fish are determined in Delineative Screening. The vessels may be jars or troughs for static or flowing water trials. The water, aeration, species and size of fish, duration of tests, and controls are the same as in Preliminary Screening. A compound is first bioassayed at 12° C. to define concentrations which evoke all-or-none responses from the fish. The approach to these concentrations may be direct by bracketing and interpolation, or it may be indirect by probit analysis, whichever is the shorter. The EC100, if seemingly practical, is confirmed by seven replicate trials.

A chemical which yields favorable results at 12° C. is retested at 17°, 22°, and 27° to evaluate the effects of temperature on the effective concentrations. At 22° and 27°, rainbow trout are omitted from the bioassays, and the loading rates for other species are half a gram or less per liter of test medium.

If a compound succeeds in these trials, its potential as a fish-control agent is estimated. Information is sought on its possible application; on the source and manufacturing costs; on possible hazards, conflicts, or limitations in use; and on the size of the market. Only if it continues to appear promising is the chemical promoted into the more elaborate and expensive Intensive Screening.

**Intensive Screening.**--Intensive Screening is directed toward development of a promising chemical as a fish-control agent and the definition of its advantages and limitations. These are broad objectives, and some subdivision of approaches is in order.

**A. Stages:** The advanced tests of a compound are accomplished in the wet laboratories, in outside pools, and finally in the field. They may include static or flowing water bioassays with durations of 24, 48, and 96 hours or longer. The vessels may include 1-gallon jars with 2.5 liters of test medium, 5-gallon jars with 15 liters of medium, and fiberglass troughs of 4-, 8-, and 16-foot lengths. The outdoor pools include 1,000-gallon vinyl units and 10- and 20-foot concrete

raceways. Proportioning pumps are used in the flowing bioassays to achieve consistent concentrations of chemical.

The initial trials of a compound in the field are conducted in waters closed to public use. These waters are found, for example, on the grounds of Federal and State fish hatcheries, on wildlife refuges, or on military reservations. Trials are manned or immediately supervised by the Laboratory staff. Subsequent experiments in the field may be accomplished by selected cooperators in State and Federal agencies.

**B. Varieties of fish:** Sixteen species in addition to those included in Preliminary and Delineative Screening are employed. They are:

1. Brook trout.
2. Northern pike.
3. Fathead minnow.
4. Brook stickleback.
5. Pumpkinseed.
6. Longear sunfish.
7. Smallmouth bass.
8. Walleye.
9. Gizzard shad.
10. Golden shiner.
11. Bigmouth buffalo.
12. Brown bullhead.
13. Channel catfish.
14. Redear sunfish.
15. Largemouth bass.
16. White crappie.

Other species may be used on occasion in advanced tests.

**C. Life stages:** Various life stages of fish, from egg to mature adult, may be involved in the more advanced bioassays to detect the presence, absence, or variation of response to a promising chemical. It may be a great advantage or disadvantage if a potential control agent is selective for one life stage and not for another.

**D. Other organisms:** The effects of a potential chemical tool on other aquatic life must be assessed at least minimally in the laboratory before it is tested in the field. Some forms such as water fleas, snails, and plants

are cultured for this purpose. Others, such as freshwater scuds, damselfly and mayfly nymphs, and tadpoles, are collected in the field. Static bioassays in glass jars are conducted with chemical concentrations and experimental conditions similar to those of the fish assays.

E. Temperature: The influences of water temperature on the biological activity of a candidate chemical are defined for all species and different life stages from bioassays under ice and at 2°, 7°, 12°, 17°, 22°, and 27° C. It is especially desirable to develop compounds that may be applied effectively during cold seasons when recreational use of public waters is low.

F. Water quality: It is recognized that water quality may exert tremendous influence on the activity of an introduced chemical. Accordingly, the effectiveness of a candidate control is observed in hard and soft waters, in alkaline and acid waters, and in waters of low and high organic content, in the laboratory and field. Initially, the formula for reconstituting deionized water is altered to detect trends.

G. Formulations: Formulation is often the key to an effective biological activity or efficient application of a compound. Density and solubility, for example, may be extremely important factors in a fish-control agent. Various formulations with carriers, wetting agents, dispersing agents, potentiators, or inhibitors may be prepared or acquired for testing. Combinations of biologically active chemicals are tested to ascertain their potentials as multiple controls for fish and aquatic weeds, fish and parasites, or fish and amphibians.

Moreover, the practicality of a control agent might depend on or be enhanced by an applicator's ability to modify or arrest its activity efficiently at a given point. Hence, the Intensive Screening includes experiments with possible deactivators, detoxifiers, or decontaminants for potential controls.

Unsatisfactory results or hazardous side effects, and new information on manufacturing, marketing, or competitive control

chemicals may contribute at any point in Intensive Screening to the abandonment of further trials and development.

### Recording and reporting results

The results obtained with each chemical are analyzed soon after completion of the Preliminary, Delineative, and Intensive Screenings. They are furnished to the contributor of the chemical immediately. After screening, we evaluate the compound as a potential fishery management tool, to determine patent positions, to refer it for studies of residues and chronic effects, and to investigate requirements for clearance and labeling.

For compounds with negative results the screening data are segregated according to the classes of chemicals involved and will be published as soon as the accumulation of data warrants. The results with chemicals which succeed as fish-control agents will be reported individually and addressed primarily to fish managers or fish culturists. Thus, we plan to define the range of effective concentrations for certain target fishes in waters of different temperatures, different types, and of various qualities.

### **SUMMARY**

The Fish Control Laboratories at La Crosse, Wis., and Warm Springs, Ga., are equipped to find and develop chemicals which may be used as fishery management tools. Objectives of the research are general and selective toxicants, attractants, repellents, anesthetics, sterilants, spawning inducers, marking dyes, medications for diseases, and sedatives for fish distribution. The facilities include chemistry, physiology, and wet laboratories; fish holding structures; and outdoor pools and raceways. The program involves three stages of chemical screening.

Preliminary Screening detects whether a selected chemical at 0.1, 1, and 10 p.p.m. has an activity in static bioassays on eight species of fish in reconstituted water at 12° and 17° C.



Delineative Screening defines the effective concentrations (EC100) of a chemical in static or flowing bioassays on eight species of fish in reconstituted water at 12<sup>o</sup>, 17<sup>o</sup>, 22<sup>o</sup>, and 27<sup>o</sup> C.

Intensive Screening is reserved for chemicals which show great promise as fish-control agents. Effective concentrations are determined in the laboratory, in outdoor pools, and in the field against 24 or more species of fish in various life stages from egg to adult, against selected aquatic organisms, and in waters of different temperatures and various qualities. The modes of action, side effects, chronic effects, and deactivators also are studied, as necessary.

The results of screening are reported direct to contributors. They are also published by the Bureau. The effective concentrations of fishery tools on target fishes under certain conditions are given.

## LITERATURE CITED

- Applegate, Vernon C., John H. Howell, James W. Moffett, B. G. H. Johnson, and Manning A. Smith.  
1961. Use of 3-Trifluormethyl-4-nitrophenol as a selective sea lamprey larvicide. Great Lakes Fishery Commission, Technical Report 1, 35 p.
- Bliss, C. L.  
1957. Some principles of bioassay. *American Scientists*, vol. 45, No. 5, p. 449-466.
- Doudoroff, P., B. G. Anderson, G. E. Burdick, P. S. Galtsoff, W. B. Hart, R. Patrick, E. R. Strong, E. W. Surber, and W. M. Van Horn.  
1951. Bio-assay methods for the evaluation of acute toxicity of industrial wastes to fish. *Sewage and Industrial Wastes*, vol. 23, No. 11, p. 1380-1397.
- Douglas, Neil H., and W. H. Irwin.  
1962. Evaluation and relative resistance of sixteen species of fish as test animals in toxicity bioassays of petroleum refinery effluents. Department of Zoology, Oklahoma State University, Contribution No. 351. Mimeo, 40 p.
- Henderson, Croswell.  
1960. Bioassay procedures, aims, and equipment. Biological Problems in Water Pollution, Technical Report W60-3, The Robert A. Taft Sanitary Engineering Center, U.S. Public Health Service, p. 246-248.
- Henderson, Croswell, and Clarence M. Tarzwell.  
1957. Bio-assays for control of industrial effluents. *Sewage and Industrial Wastes*, vol. 29, No. 9, p. 1002-1017.
- Lawrence, J. M., and R. D. Blackburn.  
1962. Evaluating herbicidal activity of chemicals to aquatic plants and their toxicity to fish in the laboratory and in plastic pools. Auburn University. Mimeo, 23 p. (In press, Proceedings of 16th Annual Conference of Southeastern Association of Game and Fish Commissioners, Columbus, S.C.).
- LesVeaux, John F.  
1959. Summary report of survey to evaluate the need for specific fish toxicants in sport fishing waters. *Progressive Fish-Culturist*, vol. 21, No. 3, p. 99-110.
- Loosanoff, V. L., C. L. MacKenzie, Jr., and L. W. Shearer.  
1960. Use of chemicals to control shellfish predators. *Science*, vol. 131, No. 3412, p. 1522-1523.
- Saila, Saul B.  
1953. Bio-assay procedures for the evaluation of fish toxicants with particular reference to rotenone. *Transactions of the American Fisheries Society*, vol. 83, p. 104-114.

FOCUS  
FOCUS  
FOCUS  
FOCUS  
Fo@S

**INVESTIGATIONS IN FISH CONTROL**

**2. Preliminary observations  
on the toxicity of antimycin A  
to fish and other aquatic animals**

By Charles R. Walker, Chemist  
Robert E. Lennon, Fishery Research Biologist  
Bernard L. Berger, Chemist  
Bureau of Sport Fisheries and Wildlife



Bureau of Sport Fisheries and Wildlife

Circular 186

Washington, D.C. • June 1964



# CONTENTS

	Page
Abstract . . . . .	1
Antimycin . . . . .	1
Sources and uses . . . . .	1
Composition and structure . . . . .	2
Physical and chemical properties . . . . .	2
Biological activity . . . . .	2
Methods and materials . . . . .	4
Laboratory tests . . . . .	4
Field tests . . . . .	4
Results of laboratory studies . . . . .	6
Trouts . . . . .	7
Herrings . . . . .	7
Perches . . . . .	8
Pikes . . . . .	8
Suckers . . . . .	9
Sunfishes . . . . .	9
Sticklebacks . . . . .	9
Minnows and carps . . . . .	9
Fresh-water catfishes . . . . .	10
Other animals . . . . .	11
Results of field studies . . . . .	11
Tests in wading pools . . . . .	11
Tests in hatchery ponds . . . . .	14
Discussion of field studies . . . . .	15
Conclusions . . . . .	16
Literature cited . . . . .	17

# Preliminary observations on the toxicity of antimycin A to fish and other aquatic animals

By Charles R. Walker, Chemist  
Robert E. Lennon, Fishery Research Biologist  
and Bernard L. Berger, Chemist  
Bureau of Sport Fisheries and Wildlife

**Abstract.**--Antimycin A, an antifungal antibiotic, has been suggested for use as a fish toxicant. Preliminary tests were made to evaluate its effects at concentrations of 0.01 to 120 p.p.b. on 24 species of fresh-water fish in the laboratory and 25 species in outdoor pools. Responses of a select group of other animals and aquatic plants are discussed. The antibiotic is a powerful fish toxicant. Carp and other rough fish were killed by small concentrations in short exposures at cool and warm temperatures. Longnose gar, bowfin, black bullheads, and yellow bullheads were relatively resistant to the quantities tested. Plankton, aquatic plants, bottom fauna, salamanders, tadpoles, and turtles were not harmed by piscicidal concentrations. Antimycin A degrades rapidly in water, especially in the presence of free hydroxide. Detoxification occurred within 24 to 96 hours. Further studies are planned on the performance of antimycin A against various life stages of fish, on other aquatic animals, and in waters of differing qualities and temperatures. The process of detoxification and the fate of residues deserve further attention.

An objective of the Fish Control Laboratories is the development of new fish toxicants that can be used safely and economically in the management of fish populations. Antimycin A exhibits properties desired in a candidate fish toxicant. It is lethal to certain target fishes in low concentration and on short exposure; it works in cool and warm water and in the presence of aquatic plants; it degrades rapidly in water and appears to leave no harmful residue.

This report summarizes data obtained on antimycin A in the laboratory and small outdoor pools and larger hatchery ponds. Development and efficacy of the compound as a fishery tool is to be further investigated.

## ANTIMYCIN

### Sources and uses

Antimycin is an antifungal antibiotic isolated from the bacteria *Streptomyces* sp. and identified by Dunshee, Leben, Keitt, and Strong (1949) at the University of Wisconsin. Following this discovery, at least seven species of *Streptomyces* were found to be producers of antimycin. Burger, Teitel, and Grunberg crystalized the antibiotic from two species of *Streptomyces* (Strong, 1956). Later at the University of Wisconsin, another culture produced an antimycin-like product which showed promise as an antibiotic for plant pathogens (Lockwood et al., 1954).

Harada and associates (Nakayama et al., 1956) in Japan discovered an antimycin-producing culture of *Streptomyces kitazawaensis* which differed from the first culture at the University of Wisconsin, but both produce an antitumor substance (*carzinomyceticus*). Research at the University of Tokyo by Watanebe et al. (1957) on *S. blastmyceticus* yielded an antibiotic called blastmycin which consists largely of antimycin A<sub>3</sub>. Harada et al. (1959) devoted special attention to the antifungal property of blastmycin as a control for rice blast disease (*Piricularia oryzae*) in Japan.

Derse and Strong (1963) related that antimycin is an antibiotic of unusual chemical structure which is toxic to yeasts, other fungi, insects, and mammals, but not to bacteria. They also reported that it is extremely toxic to goldfish at 1 p.p.b. On the basis of this observation, on the rapid degradation of the chemical, and its much lower toxicity to higher animals, they suggested that antimycin may be useful in fish management.

#### Composition and structure

The complex structure of antimycin was elucidated by Dunshee et al. (1949), Tener et al. (1953), Strong (1956) and Strong et al. (1960), van Tamelen et al. (1959 and 1961), and Dickie et al. (1963). It is illustrated in figure 1.

Lockwood et al. (1954) described antimycin as a complex made up of several active fractions which they identified from paper chromatograms as A<sub>1</sub>, A<sub>2</sub>, A<sub>3</sub>, and A<sub>4</sub> according to increasing R<sub>F</sub> values. Liu and Strong (1959) determined that one or more of these R<sub>F</sub> values were represented in antimycin A-35, antimycin A-102, blastmycin, and virosin, and they investigated them. Further study by Dickie and his associates (1963) established that the fractions differ only in the alkyl side chain (R) in figure 1. The antimycin A<sub>1</sub> and A<sub>4</sub> fractions are probably isomeric with R = n-hexyl, and calculations of the elemental composition indicate that the empirical formula is C<sub>28</sub>H<sub>40</sub>N<sub>2</sub>O<sub>9</sub>. The A<sub>2</sub> and A<sub>3</sub> isomers bear the n-butyl side chain, and the empirical formula is perhaps C<sub>26</sub>H<sub>36</sub>N<sub>2</sub>O<sub>9</sub>. The percentage composition

of fractions or isomers is very important to the biological activity of the antimycin complex.

#### Physical and chemical properties


The fermentation extracts of antimycin are dark, tarry substances which upon further purification yield a fine crystalline material. This nitrogenous, phenolic complex is characterized by solubility in polar organic solvents including ethanol, acetone, and chloroform; slight solubility in nonpolar solvents including petroleum ether, benzene, and carbon tetrachloride; and relative insolubility in water and 5-percent solutions of hydrochloric acid, sodium bicarbonate, and sodium carbonate (Keitt, Leben, and Strong, 1953).

The infrared absorption spectrum of antimycin has been identified in isolates from several cultures, although the crystalline products appear to have different properties. These differences are attributed to the intricate composition of the antibiotic and the presence of impurities associated with samples (Strong, 1956). For example, blastmycin has almost the duplicate IR spectrum of antimycin A-35 isolate, but the melting points are 166°-167° and 140.5°-141.5° C. respectively. Blastmycin is composed primarily of the antimycin A<sub>3</sub> fraction with a trace of A<sub>4</sub> in contrast to antimycin A-35, antimycin A-102, and virosin, which contain additional subcompounds A<sub>1</sub> and A<sub>2</sub> (Strong, 1956; Liu and Strong, 1959).

Antimycin is susceptible to alkaline degradation as indicated in figure 1. Hydrolytic cleavage occurs at the lactone carbonyl sites on the cyclic diester and leads to the formation of antimycic acid or blastmycic and the neutral fragment (van Tamelen et al., 1961; Liu et al., 1960; and Tener et al., 1953). The degradation is rapid in water, and detoxification of 10 p.p.b. is accomplished within 7 days according to Derse and Strong (1963); it is accelerated in the presence of light, high alkalinity, and warm temperatures.

#### Biological activity

Antimycin is a powerful and highly selective inhibitor of the electron transport in oxidative



Antimycin A<sub>1</sub> ; R = n-hexyl : C<sub>28</sub>H<sub>40</sub>N<sub>2</sub>O<sub>9</sub>

Antimycin A<sub>3</sub> ; R = n-butyl : C<sub>26</sub>H<sub>36</sub>N<sub>2</sub>O<sub>9</sub>





Figure 1.--Structure of antimycin and the assumed process of breakdown under alkaline conditions in the laboratory.

phosphorylation systems (Strong, 1956). It retards the respiration of cells, and the selective action in the electron transport chain at the cytochrome b - (Coenzyme Q) - cytochrome c has made antimycin an indispensable reagent for enzyme studies. Its effects on the succinic-oxidase system have been described as the "antimycin-A-blocked factor." Gottlieb and Ramachandran (1961) illustrated the site of action of antimycin and ascocin as follows:

Substrate --> Pyridine nucleotide --> Flavoprotein -->



cytochrome c --> cytochrome a --> oxygen

Because of its extreme potency as an inhibitor of electron transport, Derse and Strong (1963) surmised that antimycin is absorbed into the gills and interferes with respiration in fishes.

## METHODS AND MATERIALS

Crystalline antimycin A was supplied by the Wisconsin Alumni Research Foundation from Kyowa Fermentation Company, Ltd., in Tokyo, Japan. This material was isolated from the culture of *Streptomyces kitazawaensis* and had the following fractions by weight: A<sub>1</sub>, 40 percent; A<sub>2</sub>, 20 percent; A<sub>3</sub>, 20 percent; and A<sub>4</sub>, 10 percent. Although the fraction A<sub>3</sub> amounts to only 20 percent, it accounts for about 60 percent of the biological activity.

Stock solutions were prepared with 100 milligrams of crystalline antimycin A dissolved in 1 liter of acetone. They were renewed with each series of bioassays, although tests indicated that solutions in acetone are relatively stable up to 24 days. Crystalline material stored at room temperature for 2 years also remained stable.

### Laboratory tests

The methods and facilities employed for evaluation of potential fish-control agents were described by Lennon and Walker (1964). The

bioassays of antimycin A were conducted in slightly alkaline and medium hard, reconstituted water at 12°, 17°, and 22° C. Twenty-four species of fish, representing nine families, were included (table 1). They were supplied by national fish hatcheries, the Wisconsin Conservation Department, and Ozark Fisheries, Inc., and each lot was graded to a desired size before use.

Aliquots of the stock solution of antimycin A were diluted and stirred into the 1- or 5-gallon bioassay vessels in the presence of fish. The responses of the fish to the toxicant were observed at 24, 48, 72, and 96 hours.

Other animals included in bioassays were water fleas (*Daphnia magna*), crayfish (*Cambarus* sp.), damselfly nymphs (*Ischnura* sp.), tiger salamander (*Ambystoma tigrinum*), and bullfrog tadpoles (*Rana catesbiana*). They were stocked in bioassay vessels as follow: 10 water fleas or 2 damselfly nymphs in each 16-ounce jar, 1 crayfish or 2 bullfrog tadpoles in each 1-gallon jar, and 1 adult tiger salamander in each 5-gallon jar.

### Field tests

Vinyl wading pools. -- Only a few outdoor bioassays were made in 1962 and 1963 because only small quantities of toxicant were available.

TABLE 1.--The 24 fishes used in laboratory tests of antimycin A

Common name	Technical name	Size range (grams)
Gizzard shad.....	<i>Dorosoma cepedianum</i>	12.0-15.0
Rainbow trout.....	<i>Salmo gairdneri</i>	1.0- 1.6
Brown trout.....	<i>Salmo trutta</i>	1.2- 1.4
Northern pike.....	<i>Esox lucius</i>	0.5- 0.6
Stoneroller.....	<i>Camptostoma anomalum</i>	3.0- 4.0
Goldfish.....	<i>Carassius auratus</i>	1.5- 2.4
Carp.....	<i>Cyprinus carpio</i>	0.6- 2.3
Golden shiner.....	<i>Notemigonus crysoleucas</i>	1.0- 2.2
Fathead minnow.....	<i>Pimephales promelas</i>	0.9- 1.8
White sucker.....	<i>Catostomus commersoni</i>	1.3- 2.8
Bigmouth buffalo.....	<i>Ictiobus cyprinellus</i>	1.6- 2.5
Black bullhead.....	<i>Ictalurus melas</i>	0.7- 2.3
Yellow bullhead.....	<i>Ictalurus natalis</i>	1.2- 2.2
Channel catfish.....	<i>Ictalurus punctatus</i>	1.5- 1.8
Brook stickleback.....	<i>Eucalia inconstans</i>	0.6- 1.0
Green sunfish.....	<i>Lepomis cyanellus</i>	0.8- 2.5
Pumpkinseed.....	<i>Lepomis gibbosus</i>	1.0- 2.3
Bluegill.....	<i>Lepomis macrochirus</i>	1.2- 2.4
Longear sunfish.....	<i>Lepomis megalotis</i>	1.0- 2.5
Largemouth bass.....	<i>Micropterus salmoides</i>	1.8- 2.9
White crappie.....	<i>Pomoxis annularis</i>	1.5- 3.0
Iowa darter.....	<i>Etheostoma exile</i>	0.6- 1.2
Yellow perch.....	<i>Perca flavescens</i>	0.6- 3.0
Walleye.....	<i>Stizostedion vitreum</i>	0.4- 0.8



The test vessels were 1,000-gallon wading pools similar to those described by Lawrence and Blackburn (1962). Some physical, chemical, and biological conditions characteristic of ponds were simulated or intrinsic. The physical aspects included bottom soils of sand and loam, naturally varying temperatures, turbidity, and natural light. The chemistry of the well water in the pools was modified by physical and biological factors.

Of the 18 pools, 9 had 3 inches of sand on the bottom, and 9 had 3 inches of silt loam. After the pools were filled, the following were introduced: *Sagittaria latifolia*, *Elodea canadensis*, *Myriophyllum heterophyllum*, *Potamogeton nodosus*, *P. pectinatus*, *Spirogyra* spp., and phytoplankton. They were established, and the water chemistry was stabilized, during the 4- to 8-week periods before fish were added. Fingerling and adult fish were stocked 1 to 2 weeks before applications of the toxicant.

The rate of detoxification of the antimycin was observed, and some of the killed fish were shipped to the Wisconsin Alumni Research Foundation for mammalian toxicity tests. Bottom fauna were sampled and quantitated. Data were obtained on water chemistry during

the course of tests according to standard methods (American Public Health Association et al., 1960).


**Hatchery ponds.**--The Wisconsin Conservation Department provided two ponds for tests at the Delafield Warmwater Fisheries Research Station in September 1963. The surface areas of ponds No. 2 and No. 5 are 0.47 and 0.78 acre respectively (fig. 2).

Pond No. 2 was stocked with 18 species of fish at the rate of 240 pounds per acre, and pond No. 5 with 19 species at 225 pounds per acre, 1 week before antimycin was applied. Samples of water, plankton, and bottom fauna were taken from each pond soon after the fish were stocked and again just before the ponds were drained (table 2).

TABLE 2.--Concentrations of antimycin A which caused all-or-none survival among rainbow trout and brown trout at selected water temperatures in 24 and 96 hours


Species	Number of fish	Temperature (°C)	Concentrations (p.p.b.) and survival			
			At 24 hours		At 96 hours	
			All	None	All	None
Rainbow trout....	1,829	12	0.10	0.60	0.02	0.08
Do.....	120	17	0.02	0.08	<0.02	0.04
Brown trout.....	348	12	0.10	0.40	<0.06	0.08
Do.....	120	17	0.02	0.06	<0.04	0.06

POND NO. 2



CONTOUR	ACRE- FEET
0 - 1 ft.	0.44
1 - 2 ft.	0.33
2 + ft.	0.01
<b>TOTAL</b>	<b>0.78</b>

POND NO. 5



CONTOUR	ACRE- FEET
0 - 1 ft.	0.68
1 - 2 ft.	0.63
2 - 3 ft.	0.42
3 + ft.	0.17
<b>TOTAL</b>	<b>1.90</b>

Figure 2.--Sketch of ponds No. 2 and No. 5 at the Delafield Warmwater Fisheries Research Station.

Two formulations of antimycin A were prepared for application at 10 p.p.b. Pond No. 2 received 9.72 grams of technical material in a carrier formulated by the S. B. Penick Company to make up a total volume of 300 ml. Pond No. 5 received 23.37 grams of technical material dissolved in 300 ml. of acetone as a carrier. Each aliquot was mixed with 2 gallons of water and applied to a pond surface with a hand-powered garden sprayer. The applications were made from a rowboat in late afternoon, and frequent observations were made during the next 8 hours. Observations and recovery of dead fish were made daily in the following 4 days.

### RESULTS OF LABORATORY STUDIES

We found that antimycin A is toxic to the 24 species of fish tested. The toxicity varies among species and is correlated with water temperature and time. Trends in sensitivity reflect taxonomic relationships of the fishes,

and variations in susceptibility among individuals was more pronounced in some species than others. The following remarks pertain principally to the concentrations which delineate the all-or-none survival  $EC_0$  to  $EC_{100}$  ranges, of fish at 24 or 96 hours in bioassays at 12°, 17°, or 22° C. Data are shown graphically in figures 3 and 4.

Among the 24 species, the group of fish most sensitive to antimycin A includes gizzard shad, rainbow trout, brown trout, white sucker, Iowa darter, yellow perch, and walleye. All survived exposure to 0.08 p.p.b. for 24 hours at 12° C; all perished at 0.8 p.p.b.

The group intermediate in sensitivity included northern pike, stoneroller, carp, golden shiner, fathead minnow, bigmouth buffalo, brook stickleback, green sunfish, pumpkinseed, bluegill, longear sunfish, largemouth bass, and white crappie (fig. 5). Concentrations of 0.1 and 1.6 p.p.b. defined their all-or-none survival in 24 hours at 12° C.




Figure 3.--The 24-hour responses of 24 fishes in the laboratory to antimycin A in p.p.b. The solid, plain, and cross hatched bars span the ranges between the  $EC_0$  and  $EC_{100}$  at 12°, 17°, and 22° C.






Figure 4.--The 96-hour responses of 16 fishes in the laboratory to antimycin A in p.p.b. The solid, plain, and crosshatched bars span the ranges between the EC<sub>0</sub> and EC<sub>100</sub> at 12°, 17°, and 22° C.

The more resistant group of fish was represented by goldfish, black bullhead, yellow bullhead, and channel catfish. The concentrations required for kills in 24 hours at 12° C were 20 p.p.b. for channel catfish, 80 p.p.b. for yellow bullhead, 100 p.p.b. for goldfish, and 120 p.p.b. for black bullhead.

Increases in water temperature or duration of exposure made significant differences in the toxicity of antimycin A to fish in the three groups. For example, the toxicity to goldfish was increased tenfold at the higher temperature of 17° C. Among catfishes, the toxicity was enhanced about twofold at 17°. At the maximum temperature of 22°, the black and yellow bullheads were about 10 times as tolerant to antimycin A as goldfish, but channel catfish were only slightly more resistant.

For more detailed discussion on the toxicity of antimycin A, the species are grouped according to their respective families. The

families, in turn, are presented in order of their sensitivity to the toxicant.

#### Trouts

Rainbow trout and brown trout were extremely sensitive to antimycin A (table 2). At 12°, the rainbow trout succumbed to 0.6 p.p.b. in 24 hours and to 0.08 p.p.b. in 96 hours. At the same temperature, brown trout were killed by 0.4 p.p.b. in 24 hours and by 0.08 p.p.b. in 96 hours. Both species tolerated concentrations of 0.1 p.p.b. for 24 hours. In 96-hour tests, the rainbow trout survived 0.02 p.p.b. whereas brown trout withstood 0.06 p.p.b.

#### Herrings

At 12° C., all gizzard shad died within 24 hours upon exposure to 0.8 p.p.b. and within 96 hours at 0.1 p.p.b. (table 3). They were especially sensitive to the toxicant at 22°:

MOST SENSITIVE

INTERMEDIATE

LEAST SENSITIVE




Figure 5.--The order of sensitivity of 11 families of fish to antimycin A in the laboratory and field.

concentrations of 0.04 p.p.b. caused complete kills within 24 hours, and partial kills occurred at 0.02 p.p.b. or more. It was noted that a narrow range of concentrations yielded all-or-none survival, particularly at the higher temperature and longer exposure.

Perches

The Iowa darter, yellow perch, and walleye were also very sensitive to antimycin A (table 4). All specimens in 0.08 p.p.b. at 12° for 24 hours survived, but those in 0.66 p.p.b. died. The narrow range in concentrations which caused all-or-none survival was more apparent at 22° and 96-hour exposures. Yellow perch, for example, survived 0.02 p.p.b. for 24 hours and 0.01 p.p.b. for 96 hours; they

died at 0.08 p.p.b. within 24 hours and at 0.06 p.p.b. within 96 hours.

Pikes

The fry and fingerlings of northern pike were difficult to use in bioassays because of cannibalism and rapid growth. Nevertheless, they exhibited great susceptibility to antimycin A. Complete kills were obtained in 24

TABLE 3.--Concentrations of antimycin A which caused all-or-none survival among gizzard shad at selected water temperatures in 24 and 96 hours

Number of fish	Temperature (°C.)	Concentrations (p.p.b.) and survival			
		At 24 hours		At 96 hours	
		All	None	All	None
120	12	0.10	0.80	0.06	0.10
60	17	0.08	0.40	0.04	0.08
60	22	0.02	0.04	0.02	0.04

hours by 0.8 p.p.b. at 12<sup>o</sup>, 0.2 p.p.b. at 17<sup>o</sup>, and 0.1 p.p.b. at 22<sup>o</sup>. In contrast, all specimens survived 0.4 p.p.b. at 12<sup>o</sup>, 0.08 p.p.b. at 17<sup>o</sup>, and 0.06 p.p.b. at 22<sup>o</sup>. Greater toxicity was detected in 48-hour exposures, but the concentrations related to all-or-none survival were not defined.

### Suckers

The white sucker and bigmouth buffalo differed in their sensitivities to the toxicant, and the former was among the most susceptible fishes tested (table 5). Concentrations greater than 0.06 p.p.b. produced partial kills of white suckers at 12<sup>o</sup> in 96 hours, and 0.22 p.p.b. caused complete kills. Even greater sensitivity was observed at 22<sup>o</sup>. The bigmouth buffalo, on the other hand, required concentrations of antimycin A in excess of 0.4 p.p.b. for complete kills in 96 hours at 12<sup>o</sup>.

### Sunfishes

Green sunfish, pumpkinseed, bluegill, longear sunfish, largemouth bass, and white crappie were moderately sensitive to antimycin A

TABLE 4.--Concentrations of antimycin A which caused all-or-none survival among Iowa darters, yellow perch, and walleye at selected water temperatures in 24 and 96 hours

Species	Number of fish	Temperature (°C.)	Concentrations (p.p.b.) and survival			
			At 24 hours		At 96 hours	
			All	None	All	None
Iowa darters....	275	12	0.10	0.66	0.08	0.14
Yellow perch....	560	12	0.10	0.40	0.06	0.20
Do.....	504	17	0.08	0.40	0.02	0.08
Do.....	60	22	0.02	0.08	0.01	0.06
Walleye.....	20	12	0.08	0.60	--	--
Do.....	20	17	<0.08	0.10	--	--

TABLE 5.--Concentrations of antimycin A which caused all-or-none survival among white sucker and bigmouth buffalo at selected temperatures in 24 and 96 hours

Species	Number of fish	Temperature (°C.)	Concentrations (p.p.b.) and survival			
			At 24 hours		At 96 hours	
			All	None	All	None
White sucker....	810	12	0.22	0.64	0.06	0.22
Do.....	36	17	0.06	0.40	0.04	0.10
Do.....	72	22	<0.04	0.20	<0.06	0.10
Bigmouth buffalo	430	12	0.20	2.00	<0.10	0.40

(table 6). The concentrations required to cause complete kills of them at 12<sup>o</sup> ranged from 1 to 6 p.p.b. in 24 hours and from 0.2 to 0.8 p.p.b. in 96 hours. At 22<sup>o</sup>, killing concentrations ranged from 0.2 to 0.8 p.p.b. in 24 hours and from 0.08 to 0.4 p.p.b. in 96 hours.

The pumpkinseed and bluegill were the more sensitive of the six species, and they were followed in order of decreasing sensitivity by longear sunfish, largemouth bass, white crappie, and green sunfish.

### Sticklebacks

Brook sticklebacks were moderately sensitive to antimycin A at 12<sup>o</sup>. Concentrations of 5 p.p.b. killed all specimens within 24 hours, and partial kills occurred at concentrations greater than 0.5 p.p.b. The exposures beyond 24 hours failed to give consistent results. The condition of the fish was suspect because of difficulty in maintaining them without feeding during the longer test periods.

### Minnnows and carps

Stoneroller, goldfish, carp, golden shiner, and fathead minnow responded over a wide range of concentrations in an interesting pattern of susceptibility. In contrast to other families, the minnows exhibited greater variation in response between species as well as between individual specimens (table 7).

TABLE 6.--Concentrations of antimycin A which caused all-or-none survival among green sunfish, pumpkinseed, bluegill, longear sunfish, largemouth bass, and white crappie at selected water temperatures in 24 and 96 hours

Species	Number of fish	Temperature (°C)	Concentrations (p.p.b.) and survival			
			At 24 hours		At 96 hours	
			All	None	All	None
Green sunfish....	396	12	0.80	6.00	0.04	0.80
Do.....	216	17	0.20	2.00	0.08	0.60
Do.....	30	22	0.08	0.80	0.08	0.40
Pumpkinseed.....	480	12	0.40	2.00	0.08	0.20
Do.....	120	17	0.06	1.00	0.04	0.10
Do.....	180	22	0.01	0.20	0.01	0.08
Bluegill.....	1,053	12	0.20	1.00	0.06	0.40
Do.....	360	17	0.10	0.60	0.01	0.10
Do.....	200	22	0.04	0.20	0.02	0.08
Longear sunfish..	240	12	0.20	2.00	0.04	0.40
Do.....	48	17	0.08	0.40	0.04	0.20
Do.....	48	22	0.08	0.40	0.02	0.20
Largemouth bass..	800	12	0.20	<0.06	0.10	0.80
White crappie....	180	12	0.60	>2.00	--	--

TABLE 7.--Concentrations of antimycin A which caused all-or-none survival among stoneroller, goldfish, carp, golden shiner, and fathead minnow at selected water temperatures in 24 and 96 hours

Species	Number of fish	Temperature (° C)	Concentrations (p.p.b.) and survival			
			At 24 hours		At 96 hours	
			All	None	All	None
Stoneroller.....	531	12	0.08	1.00	--	--
Goldfish.....	1,469	12	4.00	100.00	0.40	2.00
Do.....	312	17	1.00	8.00	0.20	1.00
Do.....	200	22	0.30	4.00	0.08	0.60
Carp.....	240	12	0.60	2.00	0.08	0.60
Do.....	84	17	0.40	1.00	0.08	0.40
Golden shiner....	60	12	0.20	2.00	0.05	0.60
Do.....	60	17	0.10	0.40	--	--
Do.....	40	22	0.05	0.50	--	--
Fathead minnow...	816	12	0.10	2.00	0.08	0.40
Do.....	96	17	<0.80	2.00	<0.06	0.10
Do.....	78	22	<0.10	0.80	<0.10	0.10

An outstanding highlight of the screening program was the discovery that carp are vulnerable to small concentrations of antimycin A. This prolific exotic is widely considered a most undesirable species in gamefish waters and is difficult to control with existing means.

At 12° all test carp were killed by 2 p.p.b. of antimycin in 24 hours and by 0.6 p.p.b. in 96 hours; at 17° all were killed by 1 p.p.b. in 24 hours and by 0.4 p.p.b. in 96 hours. Temperatures had less effect on toxicity to carp than to most species. There were only slight differences due to temperature in 24-hour exposures and even less at 96 hours. All carp survived 0.08 p.p.b.

The results on goldfish contrasted sharply with those on carp. In fact, the goldfish was the most tolerant of the minnows tested against antimycin A. It required 100 p.p.b. for complete kills within 24 hours at 12°, but only 2 p.p.b. were needed for kills within 96 hours. Higher temperatures contributed to greater toxicities, and all goldfish perished within 96 hours when exposed to 1 p.p.b. at 17° and 0.6 p.p.b. at 22°.

Stonerollers were among the more sensitive minnows. Concentrations of toxicant as low as 1 p.p.b. killed all specimens within 24 hours at 12°, but variations in suscepti-

bility were observed; a concentration which killed on one occasion failed on the next.

The golden shiner and fathead minnow were somewhat similar to the stoneroller in sensitivity, but all-or-none effects were delineated within a narrow range of concentrations. The golden shiners succumbed to 0.6 p.p.b. within 96 hours at 12°, and survival was noted at 0.05 p.p.b. Fathead minnows died at 0.4 p.p.b. and survived at 0.08 p.p.b.

### Fresh-water catfishes

The catfishes were significantly less sensitive to antimycin A than other families (table 8). Channel catfish were more susceptible than bullheads. They survived 24-hour exposures at 12° to 2 p.p.b. but perished at 20 p.p.b. All specimens died at 6 p.p.b. in 96-hour tests at 22°.

The black bullhead was the more tolerant to the toxicant, and the yellow bullhead was only slightly less so. Concentrations of 120 and 100 p.p.b. respectively were required for complete kills in 24 hours at 12°. These concentrations are more than 100 times greater than those needed to kill fish of the most sensitive families.

The bullheads were affected by somewhat smaller quantities of chemical at 17°. Nevertheless, black bullheads tolerated 4 p.p.b. for 24 hours at 22°, and all died at 40 p.p.b.

TABLE 8.--Concentrations of antimycin A which caused all-or-none survival among black bullhead, yellow bullhead, and channel catfish at selected water temperatures in 24 hours and 96 hours

Species	Number of fish	Temperature (° C)	Concentrations (p.p.b.) and survival			
			At 24 hours		At 96 hours	
			All	None	All	None
Black bullhead...	848	12	10.0	120.0	10.0	80.0
Do.....	120	17	6.0	60.0	4.0	40.0
Do.....	120	22	4.0	40.0	2.0	40.0
Yellow bullhead..	192	12	20.0	80.0	20.0	80.0
Do.....	84	17	<10.0	60.0	--	--
Channel catfish..	120	12	2.0	20.0	2.0	20.0
Do.....	120	17	1.0	10.0	1.0	10.0
Do.....	180	22	0.8	6.0	0.8	6.0



## Other animals

Four hundred water fleas were used in trials with antimycin A. At 12° C., specimens survived 1 and 0.5 p.p.b., but died in 100 p.p.b. in 24 hours and in 10 p.p.b. in 48 hours. Their susceptibility increased with temperature. At 22°, they survived 0.1 p.p.b., but died in 10 p.p.b. in 24 hours and in 0.5 p.p.b. in 48 hours.

There were no mortalities among 20 crayfish exposed to 10 p.p.b. of toxicant at 12° for 96 hours.

Tests with 120 damselfly nymphs disclosed that the insects were relatively tolerant to antimycin A. At 12°, specimens survived 100 and 50 p.p.b. for 24 and 48 hours respectively, and 1,000 and 500 p.p.b. were required to kill them in the same time periods. At 22°, they survived 50 and 10 p.p.b., but died at 500 and 100 p.p.b. in 24 and 48 hours. The observations were not continued to 96 hours because high mortalities began to occur among controls.

Ninety-six tiger salamanders were exposed to antimycin A at 12°. Specimens survived 80 p.p.b. for 96 hours, but were killed by 600 p.p.b.

Among the 40 bullfrog tadpoles tested for 24 hours at 12°, the individuals exposed to 20 p.p.b. of toxicant survived whereas those subjected to 40 p.p.b. perished.

## **RESULTS OF FIELD STUDIES**

### Tests in wading pools

Results in 1962.--Some preliminary bioassays were conducted in 18 pools in July and October, to determine the utility of the pools as bioassay vessels and to yield information on the performance of antimycin A outdoors. A shortage of toxicant limited the scope of the trials, and a scarcity of fish of desirable species, sizes, and condition affected their validity. A number of the species were wild fish which later proved to be unsatisfactory test animals because of variable sizes, heavy parasitism, and poor condition.

The wading pools worked well as bioassay vessels. Fish, invertebrates, and plants did well in the test units and controls. There were some differences in the quantity of plankton and aquatic vegetation in the sand- and loam-bottom units because the latter were more fertile. The abundance of plants, we believe, contributed to increases in pH and alterations of alkalinity, and these in turn influenced the efficacy of antimycin A.

Goldfish, golden shiner, black bullhead, bluegill, largemouth bass, and yellow perch were exposed to 5 and 10 p.p.b. of toxicant in July. Most of them survived in the sand pools. The mortality was greater in the loam pools, especially at 10 p.p.b., but in no instance did it reach 100 percent. The black bullheads exhibited high tolerance to the toxicant in all pools.

Another series of tests was made in October with higher concentrations against rainbow trout, goldfish, golden shiner, bluntnose minnow, yellow bullhead, green sunfish, and yellow perch (table 9). The pH values in the pools at the time ranged from 7.5 to 9.9. Ninety to 100 percent of the trout, golden shiner, bluntnose minnow, green sunfish, and perch, and 60 percent of the goldfish were killed by 20 p.p.b. over sand and loam bottoms. At 40 p.p.b., there was very low survival among the trout, goldfish, and sunfish, but nearly complete survival of bullheads.

There appeared to be rapid degradation and detoxification of antimycin in the pools within 24 to 96 hours, depending on the initial concentration and the pH. Small numbers of goldfish, golden shiner, bluntnose minnow, bluegill, and largemouth bass were stocked later in pools in which antimycin A had been present for 24 to 72 hours. No more than half of the golden shiners and bluegills perished within the following 2 days.

Results in 1963.--The plants, plankton, and bottom fauna were permitted to develop in the pools for 2 months before toxicity trials. In July, acetone solutions of antimycin A were tested at 10, 20, 40, and 80 p.p.b. against eight species of fish of various sizes (tables 10, 11, 12, and 13). Golden shiners, bluegills, largemouth bass, and yellow perch were the more



TABLE 9.--Toxicity of antimycin A at 20 and 40 p.p.b. on adult and fingerling fish in sand- and loam-bottom pools

[Mortalities are cumulative by observation periods]

Species of fish	Type of bottom	Antimycin A at 20 p.p.b.					Antimycin A at 40 p.p.b.				
		Number of fish	Number dead in (hours)--				Number of fish	Number dead in (hours)--			
			24	48	96	336		24	48	96	336
<b>Adults:</b>											
Rainbow trout.....	sand	20	16	16	17	19	20	16	17	18	20
Do.....	loam	20	8	12	15	18	20	14	16	17	19
Yellow bullhead.....	sand	40	0	0	0	1	40	0	0	0	1
Do.....	loam	40	0	0	0	3	40	0	0	0	0
Green sunfish.....	sand	40	0	0	7	37	40	1	1	12	40
Do.....	loam	40	0	0	5	39	40	2	2	11	39
<b>Fingerlings:</b>											
Goldfish.....	sand	40	1	11	17	25	40	3	21	37	37
Do.....	loam	40	1	7	16	26	40	14	40	--	--
Golden shiner.....	sand	40	35	39	40	--	40	40	--	--	--
Do.....	loam	40	37	40	--	--	40	40	--	--	--
Bluntnose minnow.....	sand	40	33	39	40	--	40	40	--	--	--
Do.....	loam	40	31	39	40	--	40	40	--	--	--
Yellow perch.....	sand	40	40	--	--	--	40	40	--	--	--
Do.....	loam	40	40	--	--	--	40	40	--	--	--

TABLE 10.--Numbers and sizes of fish exposed to antimycin A in wading pools in July 1963

Species	Total number	Average weight (grams)
Goldfish.....	180	1.0
Carp.....	180	2.2
Golden shiner.....	108	1.0
Black bullhead.....	180	2.0
Do.....	180	18.0
Green sunfish.....	144	2.7
Bluegill.....	270	0.8
Do.....	180	22.0
Largemouth bass.....	144	1.5
Yellow perch.....	270	2.5

sensitive, and they were killed by 10 p.p.b. within 48 hours over sand and loam bottoms. Carp and green sunfish perished at 10 p.p.b. within 24 hours over loam bottoms and at 20 p.p.b. in the same time over sand soil. Some lots of goldfish died at 20 p.p.b., but all succumbed at 40 p.p.b. A concentration of 80 p.p.b. killed all fingerling and adult black bullheads within 48 hours over sand bottom but only one-half of them over loam.

The green sunfish and yellow perch appeared at the surface of pools within 2 hours

TABLE 11.--Toxicity of antimycin A at 10 and 20 p.p.b. on adult and fingerling fish in sand- and loam-bottom wading pools

[Mortalities are cumulative by observation period]

Species	Type of bottom	Antimycin A at 10 p.p.b.					Antimycin A at 20 p.p.b.				
		Number of fish	Number dead in (hours)--				Number of fish	Number dead in (hours)--			
			24	48	96	480		24	48	96	480
<b>Adults:</b>											
Black bullhead.....	sand	20	0	0	0	2	20	0	0	0	0
Do.....	loam	20	0	0	0	1	20	0	0	0	3
Bluegill.....	sand	20	17	20	--	--	20	20	--	--	--
Do.....	loam	20	20	--	--	--	20	20	--	--	--
<b>Fingerlings:</b>											
Goldfish.....	sand	20	6	6	6	6	20	20	--	--	--
Do.....	loam	20	7	7	7	7	20	10	10	10	10
Carp.....	sand	20	14	14	14	14	20	20	--	--	--
Do.....	loam	20	20	--	--	--	20	14	14	14	14
Golden shiner.....	sand	14	14	--	--	--	14	14	--	--	--
Do.....	loam	14	14	--	--	--	14	14	--	--	--
Black bullhead.....	sand	20	0	0	0	0	20	0	0	0	0
Do.....	loam	20	0	0	0	0	20	0	0	0	0
Green sunfish.....	sand	16	15	15	15	15	16	16	--	--	--
Do.....	loam	16	16	--	--	--	16	16	--	--	--
Bluegill.....	sand	40	40	--	--	--	40	40	--	--	--
Do.....	loam	40	40	--	--	--	40	40	--	--	--
Largemouth bass.....	sand	16	16	--	--	--	16	16	--	--	--
Do.....	loam	16	16	--	--	--	16	16	--	--	--
Yellow perch.....	sand	20	20	--	--	--	20	20	--	--	--
Do.....	loam	20	18	20	--	--	20	20	--	--	--

TABLE 12.--Toxicity of antimycin A at 40 and 80 p.p.b. on adult and fingerling fish in sand- and loam-bottom wading pools

[Mortalities are cumulative by observation period]

Species	Type of bottom	Antimycin A at 40 p.p.b.						Antimycin A at 80 p.p.b.					
		Number of fish	Number dead in (hours)--				Number of fish	Number dead in (hours)--					
			24	48	96	480		24	48	96	480		
<b>Adults:</b>													
Black bullhead.....	sand	20	0	0	0	2	20	6	20	--	--		
Do.....	loam	20	0	0	0	1	20	1	2	2	3		
Bluegill.....	sand	20	20	--	--	--	20	20	--	--	--		
Do.....	loam	20	20	--	--	--	20	20	--	--	--		
<b>Fingerlings:</b>													
Goldfish.....	sand	20	20	--	--	--	20	20	--	--	--		
Do.....	loam	20	20	--	--	--	20	20	--	--	--		
Carp.....	sand	20	20	--	--	--	20	20	--	--	--		
Do.....	loam	20	20	--	--	--	20	20	--	--	--		
Golden shiner.....	sand	14	14	--	--	--	14	14	--	--	--		
Do.....	loam	14	14	--	--	--	14	14	--	--	--		
Black bullhead.....	sand	20	0	0	0	0	20	20	--	--	--		
Do.....	loam	20	0	0	0	0	20	11	11	11	11		
Green sunfish.....	sand	16	16	--	--	--	16	16	--	--	--		
Do.....	loam	16	16	--	--	--	16	16	--	--	--		
Bluegill.....	sand	40	40	--	--	--	40	40	--	--	--		
Do.....	loam	40	40	--	--	--	40	40	--	--	--		
Largemouth bass.....	sand	16	16	--	--	--	16	16	--	--	--		
Do.....	loam	16	16	--	--	--	16	16	--	--	--		
Yellow perch.....	sand	20	20	--	--	--	20	20	--	--	--		
Do.....	loam	20	18	20	--	--	20	14	20	--	--		

TABLE 13.--Average values of analyses made on water from sand- and loam-bottom wading pools before and after applications of antimycin A in July 1963

Item	Unit of measurement	Sand		Loam	
		Before	After	Before	After
Temperature.....	°C	23	25	23	27
Resistivity.....	at 20°C	2803	2864	3037	3052
Dissolved oxygen.....	p.p.m.O <sub>2</sub>	8.7	9.1	9.7	9.7
Carbon dioxide.....	p.p.m.CO <sub>2</sub>	0.0	0.0	0.0	0.0
Hydrogen ion.....	pH	8.8	9.1	8.8	9.2
Total alkalinity..... (as phenolphthalein) (as methyl orange).....	p.p.m.CaCO <sub>3</sub>	204.4 (10.7) (193.7)	181.2 (11.5) (169.7)	198.2 (14.6) (183.6)	183.7 (14.3) (169.4)
Total hardness.....	p.p.m.CaCO <sub>3</sub>	211.8	176.0	210.6	182.0
Calcium hardness.....	p.p.m.CaCO <sub>3</sub>	53.6	47.9	60.0	53.6
Total iron.....	p.p.m.Fe <sup>0</sup>	0.0	0.0	0.0	0.0
Sulfate ion.....	p.p.m.SO <sub>4</sub>	25.8	13.4	18.3	11.1
Total phosphorus.....	p.p.m.PO <sub>4</sub>	0.099	0.071	0.082	0.106
Ammonia nitrogen.....	p.p.m.NH <sub>3</sub>	0.399	0.730	0.710	1.100
Nitrite nitrogen.....	p.p.m.NO <sub>2</sub>	0.006	0.013	0.005	0.018
Nitrate nitrogen.....	p.p.m.NO <sub>3</sub>	0.117	0.191	0.154	0.220
Chloride ion.....	p.p.m.CL	10.3	16.5	10.2	15.1

after exposure to the toxicant, and they exhibited a narcosislike condition. They showed little response to motion stimulus or handling with a dip net. Some of the larger bullheads behaved as if in distress and were subject to development of an unidentified funguslike condition on the body prior to death.

The trials in October included two formulations of antimycin. One was a solution in acetone, and the other an emulsifiable concentrate, applied to pools at 1, 5, 10, and 100 p.p.b. against 10 species of fish. The pH values at the time in all pools were about 10, and the antimycin A degraded so rapidly that most fish escaped toxic effects (table 14). The

TABLE 14.--Average values of analyses made on water from sand- and loam-bottom wading pools before and after applications of antimycin A in October 1963

Item	Unit of measurement	Sand		Loam	
		Before	After	Before	After
Temperature.....	°C	16	16	16	16
Resistivity.....	at 20°C	3561	3631	3439	3396
Dissolved oxygen.....	p.p.m.O <sub>2</sub>	10.0	9.6	10.0	9.1
Carbon dioxide.....	p.p.m.CO <sub>2</sub>	0.0	0.0	0.0	0.0
Hydrogen ion.....	pH	10.0	10.0	10.0	9.8
Total alkalinity..... (as phenolphthalein) (as methyl orange).....	p.p.m.CaCO <sub>3</sub>	116.0 (29.0) (87.0)	107.0 (27.5) (79.5)	127.0 (36.0) (91.0)	121.0 (36.0) (85.0)
Total hardness.....	p.p.m.CaCO <sub>3</sub>	143.0	148.0	157.0	156.0
Calcium hardness.....	p.p.m.CaCO <sub>3</sub>	27.4	31.0	38.0	33.0
Total iron.....	p.p.m.Fe <sup>0</sup>	0.025	0.026	0.036	0.028
Sulfate ion.....	p.p.m.SO <sub>4</sub>	17.8	17.3	16.0	14.0
Total phosphorus.....	p.p.m.PO <sub>4</sub>	0.090	0.084	0.061	0.070
Ammonia nitrogen.....	p.p.m.NH <sub>3</sub>	0.25	0.270	0.380	0.290
Nitrite nitrogen.....	p.p.m.NO <sub>2</sub>	0.0	0.0	0.0	0.0
Nitrate nitrogen.....	p.p.m.NO <sub>3</sub>	0.0	0.0	0.0	0.0
Chloride ion.....	p.p.m.CL	12.4	17.25	11.0	13.6

exceptions were those individuals exposed to 100 p.p.b. It appeared that the acetone solution of toxicant deteriorated sooner than the other preparation.

Of the 10 species of fish, 7 species succumbed totally to 100 p.p.b. of acetone-antimycin A, and 9 species to the emulsifiable formulation, within 24 hours over sand bottoms; only carp, fathead minnow, bluegill, longear sunfish, and yellow perch died over loam bottoms. The black bullhead was the sole survivor of 100 p.p.b. over both bottom types. Neither preparation of toxicant caused 100-percent kills of any species within 96 hours at 5 or 10 p.p.b.

In general, most of the susceptible fish showed signs of distress within a short time after exposure, and many came to the surface of the pools. The length of time which elapsed before death varied with the species and water temperature, and ranged from a few hours to several days. It is significant that all specimens which displayed symptoms of distress eventually died. This suggests that the action of the toxicant on fish is irreversible.

There were no grossly toxic effects by antimycin A on the plankton, bottom fauna, or aquatic plants during the course of the July and October trials. For example, in the four pools which received 20 p.p.b. of antimycin A in July, the average quantity of plankton was 0.0036 cc./l. (range: 0.0020 to 0.0044) before treatment and 0.0040 cc./l. (range: 0.0033 to 0.0061) at 20 days after treatment. The quantities in two control pools were 0.0047 and 0.0089 cc./l. during pretreatment sampling and 0.0022 and 0.0044 cc./l. during post-treatment sampling.

### Tests in hatchery ponds

There appeared to be a more rapid response of fish to the antimycin A which was formulated with an emulsifiable concentrate than with acetone. With the former preparation in pond No. 2, fish surfaced within 4 to 6 hours after application, whereas in pond No. 5 there were no comparable effects for another 10 hours. By the end of the first full

day, we saw no significant differences in the effects produced by the two formulations. Table 15 gives before and after water analyses for the two ponds.

Northern pike were the first fish to exhibit distress. They surfaced and appeared to be in a state of narcosis which was followed by complete locomotor ataxia. The rainbow trout, white suckers, carp, walleye, and sunfishes followed in order with similar symptoms. The great majority of specimens were dead within 48 hours (tables 16 and 17). It is noteworthy that goldfish--a species which was relatively

TABLE 15.--Analyses of water from ponds No. 2 and No. 5 at Delafield Warmwater Fisheries Research Station before and after applications of antimycin A in September 1963

Item	Unit of measurement	Pond No. 2		Pond No. 5	
		Before	After	Before	After
Temperature.....	°C	21	15	21	17
Resistivity.....	at 20°C	2550	2600	2525	2600
Dissolved oxygen.....	p.p.m.O <sub>2</sub>	6.7	6.9	7.5	8.2
Carbon dioxide.....	p.p.m.CO <sub>2</sub>	3.4	0.0	2.0	0.0
Hydrogen ion.....	pH	8.0	8.4	8.9	8.5
Total alkalinity.....	p.p.m.CaCO <sub>3</sub>	210.0	202.0	201.1	189.5
(as phenolphthalein).....		( 0.0)	( 0.0)	( 8.8)	( 0.0)
(as methyl orange).....		(210.0)	(202.0)	(192.3)	(189.5)
Total hardness.....	p.p.m.CaCO <sub>3</sub>	213.0	220.0	202.0	208.0
Calcium hardness.....	p.p.m.CaCO <sub>3</sub>	77.0	82.0	80.0	75.0
Manganese.....	p.p.m.Mn <sup>0</sup>	0.0	0.0	0.0	0.0
Total iron.....	p.p.m.Fe <sup>0</sup>	0.00	0.05	0.00	0.13
Sulfate ion.....	p.p.m.SO <sub>4</sub>	44.3	38.0	39.0	35.0
Total phosphorus.....	p.p.m.PO <sub>4</sub>	1.40	0.10	0.50	0.15
Ammonia nitrogen.....	p.p.m.NH <sub>3</sub>	0.20	0.19	0.18	0.38
Nitrite nitrogen.....	p.p.m.NO <sub>2</sub>	0.0	0.0	0.0	0.0
Nitrate nitrogen.....	p.p.m.NO <sub>3</sub>	0.07	0.50	0.07	0.43
Chloride ion.....	p.p.m.CL	--	14.5	--	15.0

TABLE 16.--Effects of 10 p.p.b. of antimycin A in emulsifiable concentrate on 18 species of fish in pond No. 2

Species	Total fish stocked	Average length (inches)	Average weight (grams)	Number of fish dead at (hours)--			
				24	48	96	480
Longnose gar.....	3	25.6	658	0	0	0	0
Bowfin.....	1	16.8	545	0	0	0	0
Rainbow trout.....	312	4.0	82	312	--	--	--
Northern pike.....	7	17.8	713	5	5	5	7
Goldfish.....	740	2.4	9	740	--	--	--
Carp.....	18	15.3	1,126	17	18	--	--
White sucker.....	4	15.1	554	3	3	3	4
Black bullhead....	600	3.8	18	0	0	0	0
Yellow bullhead...	4	8.3	168	0	0	0	0
Brown bullhead....	1	4.2	50	0	0	0	0
Rock bass.....	1	8.0	136	1	--	--	--
Green sunfish.....	3	3.8	14	0	0	1	3
Pumpkinseed.....	13	4.6	41	11	12	13	--
Bluegill.....	27	6.1	68	21	21	22	27
Black crappie.....	7	8.3	95	5	5	5	7
Largemouth bass...	4	15.4	795	1	4	--	--
Hybrid sunfish....	1,400	1.7	9	1,400	--	--	--
Walleye.....	1	13.5	318	1	--	--	--

TABLE 17.--Effects of 10 p.p.b. of antimycin A in acetone solution on 19 species of fish in pond No. 5

Species	Total fish stocked	Average length (inches)	Average weight (grams)	Number of fish dead at (hours)--			
				24	48	96	480
Longnose gar.....	3	24.6	395	0	0	2	2
Bowfin.....	1	21.8	1,771	0	0	0	0
Rainbow trout.....	470	4.1	86	470	--	--	--
Northern pike.....	8	19.1	976	6	6	8	--
Goldfish.....	1,400	2.7	9	1,400	--	--	--
Carp.....	27	15.3	1,112	26	27	--	--
White sucker.....	6	15.7	636	5	5	5	6
Black bullhead....	875	3.7	14	0	2	3	157
Yellow bullhead...	1	5.7	59	0	0	0	0
Brown bullhead....	6	11.4	377	0	0	0	1
Rock bass.....	1	8.2	136	0	0	0	1
Green sunfish.....	4	4.2	23	0	1	2	4
Pumpkinseed.....	24	4.6	32	2	12	12	24
Bluegill.....	43	6.4	86	19	28	36	43
Black crappie.....	9	8.8	136	1	4	4	9
Largemouth bass...	5	12.8	477	0	2	4	5
Hybrid sunfish....	2,055	1.7	9	2,055	--	--	--
Walleye.....	2	13.0	386	2	--	--	--
Drum.....	1	11.6	272	0	0	1	--

tolerant to antimycin A in the laboratory--died in both ponds within 24 hours.

The longnose gar, bowfin, black bullhead, yellow bullhead, and brown bullhead were the only species which were not affected greatly by the toxicant at 10 p.p.b. Seventy percent of them were recaptured alive when the ponds were drained after 20 days.

The detoxification of antimycin A was monitored throughout the first 96 hours. It occurred within 72 hours after application, and fish placed in live cages after this time survived until the ponds were drained.

Plankton was sampled in both ponds during the experimental period. In pond No. 2, the pretreatment quantity was 0.018 cc./l and the posttreatment quantity was 0.047 cc./l. Pond No. 5 had pretreatment and posttreatment quantities of 0.0035 and 0.039 cc./l. None of the relatively minor changes was attributed to the toxicant. Also, there were no observable changes in the aquatic plants in the ponds.

Pretreatment and posttreatment samples of bottom fauna were taken. We concluded that antimycin A was nontoxic to the 15 taxonomic groups which were represented in both ponds because there were no significant changes in species composition or numerical abundance (table 18). The midges were the more numerous in all samples, and they increased by 55 to 65 percent during the experimental period. The nymphs of mayflies, dragonflies, and

TABLE 18.--Abundance of bottom fauns in ponds No. 2 and No. 5 before and after treatment with 10 p.p.b. of antimycin A

[Each collection consisted of 16 one-square foot samples

Organism	Average number per square foot			
	Pond No. 2		Pond No. 5	
	Sept. 23	Oct. 15	Sept. 17	Oct. 14
Horsehair worm (Nematomorpha)....	10.7	0.7	1.0	1.0
Aquatic earth-worm (Oligochaeta).....	0.0	0.0	34.5	3.0
Leech (Hirundinea).....	0.0	1.3	5.2	1.0
Scuds (Amphipoda).....	4.7	2.0	17.5	39.0
Mayflies (Ephemeroptera)....	9.0	145.3	6.8	6.5
Damselflies (Zygoptera).....	1.7	2.0	6.8	7.8
Dragonflies (Anisoptera).....	0.0	0.7	0.5	0.5
Waterbugs (Hemiptera).....	0.0	2.7	1.0	1.5
Caddisflies (Trichoptera)....	0.7	2.0	0.2	0.0
Water beetles (Coleoptera).....	2.7	18.0	5.0	3.5
Mosquitoes (Culicidae).....	0.0	0.0	0.8	0.0
Midges (Tendipedidae)....	209.7	388.0	269.5	422.0
Biting midges (Ceratopogonidae)..	1.3	0.0	2.8	1.0
Soldierflies (Stratiomyidae)...	0.3	0.0	0.0	0.0
Snails (Gastropoda).....	4.0	30.7	72.8	30.5
Total.....	244.8	593.4	424.4	517.3

damselflies were also more abundant in the posttreatment samples.

Care was taken to note any gross effects of the toxicant on frogs, salamanders, and turtles, but there were none.

#### Discussion of field studies

There was a lack of consistency in the performance of antimycin A in sand- and loam-bottom pools in July and October, 1962 and 1963, and in the hatchery ponds. The cause, we believe, was the chemistry of the waters and particularly the presence of the hydroxyl ion.

An alkaline shift occurred in the wading pools as the plant biomass increased. The relatively hard, well water which was used to fill the pools was gradually softened because of the decrease in calcium. There was a shift from bicarbonate (methyl orange alkalinity) to free hydroxide (phenolphthalein alkalinity). The measure of the acid-base shift was pH which rose from 7.5 upward to 10 or more. Diurnal fluctuations of several pH units are not uncommon in ponds, and the pH in wading pools ranged accordingly between morning and afternoon.

The highest pH values were observed in late afternoon in the presence of abundant plants and sunshine. In this situation, the



hydroxyl ions appear, and often they are not checked by buffering salts. Magnesium prevails as calcium ions are removed from solution, and the result is the sort of alkaline shift observed in softer waters.

We assume that the relative success of the toxicity trials in hatchery ponds was due in large part to the fact that the water had high buffer capacity and little reserve alkalinity in the form of hydroxide. Thus, the antimycin A was not subject to immediate detoxification by action of free hydroxide, and the 10 p.p.b. were effective in killing fish.

In contrast, the poorer results obtained in the wading pools reflected the greater concentrations of free hydroxide present. In July 1963, the pools had approximately the same pH and total alkalinity as the hatchery ponds, but there was more free hydroxide present. Therefore, the degradation of the toxicant was more rapid, and 20 to 40 p.p.b. were needed to kill fish.

The contrast was heightened by results in October 1963. The water was much softer and lower in buffer capacity, and there was even more free hydroxide present. The pH ranged up to 10. Under these conditions, there was almost immediate detoxification of the antimycin, and only partial fish kills were obtained at 100 p.p.b.

## CONCLUSIONS

Antimycin A is a powerful toxicant to fresh-water fish. We observed the responses of many specimens to concentrations which ranged from 0.01 to 120 p.p.b. Among them, the carp--a most undesirable fish in many waters--proved vulnerable to small concentrations and short exposures at cool and warm temperatures. Other fishes which at times may be undesirable, such as goldfish, white suckers, green sunfish, and pumpkinseeds, were also killed.

The sensitivities to antimycin A varied among species, and they were correlated with temperature and duration of exposure. The tests in the laboratory at 12<sup>o</sup>, 17<sup>o</sup>, and 22<sup>o</sup> C.

indicated that smaller quantities of toxicant or shorter exposures produced kills of fish in warmer waters, but the results at 12<sup>o</sup> were nonetheless satisfactory.

There were three general degrees of sensitivity detected among the 24 species of fish in the laboratory and a similar order among the 25 species used in outdoor trials. Indicative of the extremes in response, gizzard shad perished at 0.04 p.p.b. of toxicant whereas black bullheads survived 100 p.p.b. There also appeared to be a tendency for sensitivities to follow family lines, but species in the nine families tested exhibited great variations in susceptibility. For example, fingerling carp in the laboratory died within 24 hours upon exposure to 0.6 p.p.b. at 12<sup>o</sup>, but up to 100 p.p.b. were required for complete kills of goldfish.

Observations in the laboratory and field demonstrated that antimycin A was less toxic to other animals. Water fleas were killed by 100 p.p.b. in 24 hours at 12<sup>o</sup>, but their susceptibility increased at warmer temperatures or longer exposures. Crayfish were not harmed by 10 p.p.b. over 96 hours, and damselfly nymphs survived 50 p.p.b. for 48 hours. Tiger salamanders survived 80 p.p.b. for 96 hours at 12<sup>o</sup>, and bullfrog tadpoles were unharmed by 20 p.p.b. for 24 hours.

The plankton in wading pools and hatchery ponds was not significantly affected during experiments, and there was no gross evidence of toxicity to filamentous algae, and submersed and emergent plants. No deleterious effects were detected on the composition, numbers, and growth of bottom fauna in hatchery ponds.

Antimycin A degraded rapidly in water, and detoxification was complete within 24 to 96 hours under field conditions. The rate of molecular breakdown was accelerated sharply in the presence of free hydroxide, and this suggests a possibility for artificial detoxification. Bioassays with fish following the degradation of the toxicant revealed an absence of harmful residues in water.

Further investigation on antimycin A as a fish toxicant is warranted in the laboratory



and field. Studies are contemplated or in progress at the Fish Control Laboratories on its performance against various life stages of fish from egg to adult; against additional species; on minimum killing concentrations and exposures; in waters of various chemistries; and at cold and warm temperatures. Appropriate formulations for standing and flowing waters are desirable. Further attention must also be given to the effects of the toxicant on other aquatic organisms. The factors in water which contribute to degradation of the toxicant deserve study, and the nature and fate of residues require definition. Also--and depending on adequate supplies of toxicant--many, and more comprehensive, trials in the field are needed for full and fair evaluation of this material which has potential value in fishery management and research.

## LITERATURE CITED

American Public Health Association, American Water Works Association, and Water Pollution Control Federation.

1960. Standard methods for the examination of water and waste-water. 11th, ed. American Public Health Association, New York, 626 p.

Derse, P. H., and F. M. Strong.

1963. Toxicity of antimycin to fish. *Nature*, vol. 200, No. 4906, p. 600-601.

Dickie, J. P., M. E. Loomans, T. M. Farley, and F. M. Strong.

1963. The chemistry of antimycin A. XI. N-substituted 3-formamidosalicylic amides. *Journal of Medicinal Chemistry*, vol. 6, p. 424-427.

Dunshee, B. R., C. Leben, G. W. Keitt, and F. M. Strong.

1949. Isolation and properties of antimycin A. *Journal of the American Chemical Society*, vol. 71, p. 2436-2437.

Gottlieb, D., and S. Ramachandran.

1961. Mode of action of antibiotics. 1. Site of action of ascocin. p. 391-396.

Harada, Y., K. Nakayama, and F. Okamoto.

1959. Antimycin A, an antibiotic substance useful in prevention and treatment of imochiby, a disease of rice. *Japan*, 2200, (Chemical Abstracts, vol. 53 (1959), p. 19286i).

Keitt, George W., Curt Leben, and Frank M. Strong. 1953. Antimycin and process for production. U.S. Patent Office. Patent No. 2,657,170, 10 p.

Lawrence, J. M., and R. D. Blackburn.

1962. Evaluating herbicidal activity of chemicals to aquatic plants and their toxicity to fish in the laboratory and in plastic pools. Auburn University, mimeo: p. 1-23, (*In press*, Proceedings of 16th Annual Conference of Southeastern Association of Game and Fish Commissioners, Columbia, S.C.).

Lennon, Robert E., and Charles R. Walker.

1964. Investigations in Fish Control; 1 Laboratories and methods for screening fish-control chemicals. Bureau of Sport Fisheries and Wildlife, Circular 185.

Liu, Wen-chih, and F. M. Strong.

1959. The chemistry of antimycin A. VI. Separation and properties of antimycin A subcomponents. *Journal of the American Chemical Society*, vol. 81, p. 4387-4390.

Liu, Wen-chih, E. E. van Tamelen, and F. M. Strong.

1960. The chemistry of antimycin A. VIII. Degradation of antimycin A. *Journal of the American Chemical Society*, vol. 82, p. 1652-1654.

Lockwood, J. L., C. Leben, and G. W. Keitt,

1954. Production and properties of antimycin A from a new *Streptomyces* isolate. *Phytopathology*, vol. 44, p. 438-446.

Nakayama, K., F. Okamoto, and Y. Harada.

1956. Antimycin A: Isolation from *Streptomyces kitazawaensis* and activity against rice plant blast fungi. *Journal of Antibiotics (Japan) Ser. A* 9, p. 63-66 (Chemical Abstracts, vol. 53 (1959), p. 19030h).

Strong, F. M.

1956. Topics in microbial chemistry. John Wiley and Sons, Inc., New York, 166 p.

Strong, F. M., J. P. Dickie, M. E. Loomans, E. E. van Tamelen, and R. S. Dewey.

1960. The chemistry of antimycin A. IX. Structure of the antimycins. *Journal of the American Chemical Society*, vol. 82, p. 1513.

Tener, G. M., F. Merlin Bumpus, Bryant R. Dunshee, and F. M. Strong.

1953. The chemistry of antimycin A. II. Degradation studies. *Journal of the American Chemical Society*, vol. 75, p. 1100-1104.

Tener, G. M., E. E. van Tamelen, and F. M. Strong.

1953. The chemistry of antimycin A. III. The structure of antimycic acid. *Journal of the American Chemical Society*, vol. 75, p. 3623-3625.

van Tamelen, E. E., F. M. Strong, and U. Carol Quarck, 1959. The chemistry of antimycin A. IV. Studies on the structure of antimycin lactone. *Journal of the American Chemical Society*, vol. 81, p. 750-751.

van Tamelen, E. E., J. P. Dickie, M. E. Loomans, R. S. Dewey, and F. M. Strong. 1961. The chemistry of antimycin A. X. Struc-

ture of the antimycins. *Journal of the American Chemical Society*, vol. 83, p. 1639-1646.

Watanabe, K., T. Tanaka, K. Fukuhara, N. Miyairi, H. Yonehara, and H. Umezawa. 1957. Blastmycin, a new antibiotic from *Streptomyces*. *Journal Antibiotics (Japan)*, vol. A10, p. 39-45. (Chemical Abstracts, vol. 53 (1959), p. 22221g).

#### LITERATURE CITED

Association of Public Health and Water Pollution Control Engineers, 1959. Standard methods for the examination of water and waste water, 11th ed. American Public Health Association, New York, 647 p.

Case, F. H., and F. M. Strong. 1955. Toxicity of aminoglycoside antibiotics to fish. *Health, vol. 200*, no. 400, p. 600-601.

Chase, J. P., M. E. Loomans, I. M. Parker, and F. M. Strong. 1963. The chemistry of antimycin A. XII. Antimycin B. *Journal of the American Chemical Society*, vol. 85, p. 2450-2457.

Fincher, B. R., C. J. Chen, F. W. Kemp, and F. M. Strong. 1962. Isolation and properties of antimycin W. *Journal of the American Chemical Society*, vol. 84, p. 2450-2457.

Guthrie, G., and S. Hansch. 1951. Mode of action of antibiotic. I. Study of action of actidione. p. 381-395.

Harada, Y., K. Hasegawa, and F. Umemoto. 1958. Antimycin A as antibiotic against swine disease of fish. *Japan, 1958*, (Chemical Abstracts, vol. 52 (1958), p. 12507).

In prevention and treatment of infectious disease of fish. *Japan, 1958*, (Chemical Abstracts, vol. 52 (1958), p. 12507).

International Conference on the Control of Fish Diseases, 1958. *Journal of the American Chemical Society*, vol. 80, p. 100-105.

1958. The chemistry of antimycin A. XI. The structure of antimycin A. *Journal of the American Chemical Society*, vol. 80, p. 100-105.

1958. The chemistry of antimycin A. X. Structure and properties of antimycin A. *Journal of the American Chemical Society*, vol. 80, p. 100-105.

1958. The chemistry of antimycin A. IX. Structure and properties of antimycin A. *Journal of the American Chemical Society*, vol. 80, p. 100-105.

1958. The chemistry of antimycin A. VIII. Structure and properties of antimycin A. *Journal of the American Chemical Society*, vol. 80, p. 100-105.

1958. The chemistry of antimycin A. VII. Structure and properties of antimycin A. *Journal of the American Chemical Society*, vol. 80, p. 100-105.

1958. The chemistry of antimycin A. VI. Structure and properties of antimycin A. *Journal of the American Chemical Society*, vol. 80, p. 100-105.

1958. The chemistry of antimycin A. V. Structure and properties of antimycin A. *Journal of the American Chemical Society*, vol. 80, p. 100-105.

1958. The chemistry of antimycin A. IV. Structure and properties of antimycin A. *Journal of the American Chemical Society*, vol. 80, p. 100-105.

1958. The chemistry of antimycin A. III. Structure and properties of antimycin A. *Journal of the American Chemical Society*, vol. 80, p. 100-105.

1958. The chemistry of antimycin A. II. Structure and properties of antimycin A. *Journal of the American Chemical Society*, vol. 80, p. 100-105.

1958. The chemistry of antimycin A. I. Structure and properties of antimycin A. *Journal of the American Chemical Society*, vol. 80, p. 100-105.