

NOAA Technical Report NMFS SSRF-711



# A List of the Marine Mammals of the World

Dale W. Rice

April 1977

U.S. DEPARTMENT OF COMMERCE  
National Oceanic and Atmospheric Administration  
National Marine Fisheries Service

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

# CONTENTS

Page

Introduction . . . . .	1
Carnivora . . . . .	2
Ursidae (1) <sup>1</sup> . . . . .	2
Odobenidae (1) . . . . .	2
Otariidae (14) . . . . .	2
Mustelidae (1) . . . . .	3
Phocidae (19) . . . . .	4
Sirenia . . . . .	5
Dugongidae (2) . . . . .	5
Trichechidae (3) . . . . .	6
Mysticeti . . . . .	6
Eschrichtiidae (1) . . . . .	6
Balaenopteridae (6) . . . . .	6
Balaenidae (3) . . . . .	6
Odontoceti . . . . .	7
Platanistidae (5) . . . . .	7
Delphinidae (37) . . . . .	7
Monodontidae (2) . . . . .	10
Physeteridae (3) . . . . .	11
Ziphiidae (18) . . . . .	11
Synonyms . . . . .	12
Acknowledgments . . . . .	13
Literature Cited . . . . .	13

---

<sup>1</sup>Numbers in parentheses indicate number of species in each family.

# A List of the Marine Mammals of the World<sup>1</sup>

DALE W. RICE<sup>2</sup>

## ABSTRACT

Listed are the 116 species of Recent marine mammals, including freshwater species of the predominantly marine groups. The number of species are: Order Carnivora, 36 (polar bear, sea otter, and 34 pinnipeds); Order Sirenia, 5; Order Mysticeti, 10; and Order Odontoceti, 65. The geographic distribution of each species is indicated.

## INTRODUCTION

Listed here are the living and recently extinct marine mammals of the world: the sea otter, polar bear, pinnipeds, sirenians, and cetaceans. Living freshwater pinnipeds, sirenians, and cetaceans are included.

Attempts to classify marine mammals are difficult because they are poorly known. Some live on the high seas and others on remote oceanic islands or among polar ice fields. Some sirenians and smaller cetaceans live in tropical waters seldom visited by mammalogists. The carcasses of marine mammals are large, greasy, bloody, and often putrefied before they are brought to the attention of biologists. They are difficult and expensive to collect and to preserve for study. As a result, some kinds are represented in scientific collections by only a few skulls and their external appearance is poorly known. Thus, any list of the marine mammals, especially of the smaller cetaceans, can be regarded only as provisional.

The seals have usually been separated from the terrestrial carnivores as Order (or Suborder) Pinnipedia. It is now clear, however, that the pinnipeds had a common origin with the weasel, racoon, dog, and bear families, and together they constitute one of the two main subdivisions of living carnivores (Mitchell and Tedford 1973).

The cetaceans have usually been regarded as comprising a single order, but their origins are obscure, and it is questionable whether they are monophyletic. The Odontoceti (toothed whales) and the Mysticeti (baleen whales) have been separate since at least the late Eocene, and all authors have recognized them as valid taxa at either the subordinal or ordinal level. The differences between the two groups are as great as those between some of the universally recognized orders of mammals (Rice, in Anderson and Jones 1967). Therefore I follow Kleinenberg (1958) and some other authors in ranking them as separate orders.

In the Order Odontoceti, the species of the genera *Platanista*, *Sousa*, *Sotalia*, and *Tursiops* herein recog-

nized as valid have been changed to conform to the "List of Smaller Cetaceans Recognized," agreed upon at the special meeting on smaller cetaceans held by the Scientific Committee of the International Whaling Commission (1975). The present list now agrees with the IWC Scientific Committee's list. It appears that the specific classification of the Odontoceti is approaching a consensus. Future studies may indicate that certain closely related allopatric forms now listed as separate species should be regarded as conspecific; such cases are noted in the text. Recent studies on some species of small cetaceans (e.g., *Stenella longirostris* and *S. attenuata*) have revealed considerable geographic variation. Most species of cetaceans are still too poorly known for subspecies to be defined, but I have listed the proposed subspecies that appear to be valid.

In the few cases where the nomenclature or classification in the present list differs from that in the working list compiled by the U.S. Marine Mammal Commission (1976), the reasons are explained in the text or in the references cited therein.

The following names are senior synonyms of names used in this list; according to Article 23b of the International Code for Zoological Nomenclature (International Trust for Zoological Nomenclature 1964), these names are nomina oblita and cannot replace the names herein employed:

*Stenorhynchus* E. Geoffroy St.-Hilaire and F. Cuvier, 1826 (= *Hydrurga* Gistel, 1848)

*Susu* Lesson, 1828 (= *Platanista* Wagler, 1830)

*Nodus* Wagler, 1830 (= *Mesoplodon* Gervais, 1850)

*Tursiops nesarnack* (Lacépède, 1804) [= *T. truncatus* (Montagu, 1821)]

*Phoca vitulina stejnegeri* Allen, 1902 (= *Phoca vitulina kurilensis* Inukai, 1942)

Hyperoodontidae Gray, 1846 (= Ziphiidae Gray, 1865).

Synonyms commonly used in recent literature are listed on page 12; see Scheffer (1958) for synonymy of pinnipeds and Hershkovitz (1966) for synonymy of cetaceans.

Vernacular names are included for most species. In selecting vernacular names, I have been guided, but not bound, by the principles adopted by the American Fisheries Society's Committee on Names of Fishes

<sup>1</sup>A revision of two earlier lists under the same title (Scheffer and Rice 1963; Rice and Scheffer 1968).

<sup>2</sup>Marine Mammal Division Northwest Fisheries Center, National Marine Fisheries Service, NOAA, Seattle, WA 98115.

(Bailey et al. 1970) and the American Ornithologists' Union's Committee on Classification and Nomenclature (1973). Many small cetaceans lack distinctive vernacular names; hence some names listed here are "book" names. I am not attempting to "standardize" vernacular names—the time is not ripe for that. I hope that field workers will record or invent more appropriate names where needed.

The report deals only with the Recent marine mammals. Readers interested in fossil forms are referred to the following publications: Winge (1921); Miller (1923); Kellogg (1928, 1936); Slijper (1936); Simpson (1945); Piveteau (1958, 1961); Reinhart (1959); Matthes (1962); King (1964); and Romer (1966).

## Order CARNIVORA

The living carnivores comprise two superfamilies—Feloidea (mongooses, cats, hyenas, etc.) and Canoidea (dogs, bears, raccoons, weasels, seals, etc.) (Mitchell and Tedford 1973). Only the latter superfamily includes marine species.

### Family URSIDAE

There are five living genera of bears and giant pandas (Hendey 1972), but only one contains a marine species.

#### Genus URSUS Linnaeus, 1758

Only one of the four species of this genus is marine.

*Ursus maritimus* Phipps, 1774 (polar bear).

Ice-covered regions of Arctic Ocean and contiguous seas, and adjacent coasts and islands.

### Family ODOBENIDAE

Mitchell (1975) regarded the walruses as a subfamily of the Otariidae, whereas Repenning (1975) maintained them as a separate family. Pending a consensus, I follow most previous authors in listing them as a separate family.

#### Genus ODOBENUS Brisson, 1762

*Odobenus rosmarus* (Linnaeus, 1758) (walrus).

Shallow waters near ice in the Arctic Ocean and adjacent seas. Three subspecies are currently recognized: *O. r. rosmarus* in the Atlantic-Arctic; *O. r. divergens* (Illiger, 1815), in the Pacific-Arctic; and *O. r. laptevi* Chapskii, 1940, in the Laptev Sea. The Atlantic subspecies contains two breeding groups, one from the Kara Sea to eastern Greenland and another from western Greenland and eastern Canada. The latter population may be sub-specifically distinct. There are late Pleistocene to prehistoric records as far south as California, Michigan, North Carolina, and France (Repenning, pers. commun.).

### Family OTARIIDAE

#### Genus PHOCARCTOS Peters, 1866

*Phocarctos hookeri* (Gray, 1844) (Auckland sea lion; New Zealand sea lion).

Subantarctic islands south of New Zealand; breeds regularly only at Carnley Harbor and Enderby Island in the Auckland Islands, rarely at Campbell Island. Hauls out on Snares Islands, Macquarie Island, and South Island, N.Z. Occurred on North Island, N.Z., less than 1,000 yr ago. For discussion of the relationships of *Phocarctos*, *Neophoca*, and *Zalophus* see King (1960).

#### Genus OTARIA Péron, 1816

*Otaria flavescens* (Shaw, 1800) (South American sea lion).

Coastal waters from Recife das Tôrres, Brazil, and Zorritos, Peru, southward to Strait of Magellan and Falkland Islands. Some authors have used the specific name *byronia* de Blainville, 1820, but P. Hershkovitz pointed out (pers. commun.) that "the type of *flavescens* was a tangible specimen preserved in the old Leverian museum. It was adequately described and figured, is perfectly identifiable, and has a valid type locality. Its name has priority, usage and currency."

#### Genus ZALOPHUS Gill, 1866

*Zalophus californianus* (Lesson, 1828) (black sea lion; California, Japanese, and Galapagos sea lions).

One subspecies, *Z. c. californianus*, breeds from San Miguel Island, Calif., south to Punta Entrada, Baja California, and on islands in the upper Gulf of California, ranging at sea north to Vancouver Island, south to Cabo San Lucas and Mazatlán. A second subspecies of unconfirmed validity, *Z. c. japonicus* (Peters, 1866), was known from the Sea of Japan but was probably exterminated in the 1950's (International Union for Conservation of Nature and Natural Resources 1972; Nichiwaki 1973). A third subspecies, *Z. c. wollebaeki* Sivertsen, 1953, breeds on the Galápagos Islands.

#### Genus NEOPHOCA Gray, 1866

*Neophoca cinerea* (Péron, 1816) (Australian sea lion).

Coastal waters from Kangaroo Island, South Australia, to Houtman Rocks, Western Australia. There is a late Pleistocene record from Melbourne, Victoria.

#### Genus EUMETOPIAS Gill, 1866

*Eumetopias jubatus* (Schreber, 1776) (northern sea lion).

Breeds along west coast of North America from San Miguel Island, Calif., northwest to Prince William Sound and the Alaska Peninsula, throughout the Aleutian and Pribilof islands, along the east coast of Kamchatka, throughout the Kuril Islands, and on islands in the Okhotsk Sea. Some move north into the Bering Sea in summer, as far as St. Lawrence Island. Sometimes hauls out on ice. The spelling of *jubatus* follows a rule in International Trust for Zoological Nomenclature (1964:31): "a noun of variable gender . . . is to be treated as masculine . . ."

#### Genus CALLORHINUS Gray, 1859

*Calorhinus ursinus* (Linnaeus, 1758) (northern fur seal).

Breeds on Ostrov Tyuleniy (Robben Island) in the Okhotsk Sea, some of the Kuril Islands, the Pribilof and Commander islands in the Bering Sea, and San Miguel Island off southern California. Formerly bred on Ostrov Iony in the Okhotsk Sea 120 miles north of Sakhalin (Stejneger 1898), and possibly on Buldir Island in the Aleutians (Murie 1959) and the Farallon Islands off central California (Repenning et al. 1971). Bones, including those of young pups, from San Miguel Island, the Monterey area, and Año Nuevo Point, Calif., and from Bella Bella, British Columbia, suggest that it formerly bred all along the west coast from San Miguel Island to Alaska (Repenning, pers. commun.).

#### Genus ARCTOCEPHALUS E. Geoffroy Saint-Hilaire and F. Cuvier, 1826

The breeding ranges of the species of *Arctocephalus* are strictly allopatric; further studies on the relationships of these seals are much needed. The present classification follows Repenning et al. (1971).

*Arctocephalus pusillus* (Schreber, 1776) (Giant fur seal; Victorian, or Tasmanian, and South African, or Cape, fur seals).

There are two widely separated subspecies. *Arctocephalus p. pusillus* breeds in temperate coastal waters from Cape Cross, South West Africa, to Algoa Bay, South Africa; ranges north to Angola. *Arctocephalus p. doriferus* Wood Jones, 1925, breeds along the coast of southeastern Australia from Lady Julia Percy Island east to the Skerries, Victoria, including coasts of Tasmania and islands in Bass Strait; ranges east to Port Stephens, N.S.W.

*Arctocephalus gazella* (Peters, 1875) (Antarctic fur seal).

Islands south of the Antarctic Convergence; South Shetlands, South Orkneys, South Sandwich, South Georgia, Bouvet, Kerguelén (not breeding), Heard, and McDonald.

*Arctocephalus forsteri* (Lesson, 1828) (Antipodean fur seal; Western Australian and New Zealand fur seals).

Breeds around South Island, N.Z., on nearby subantarctic islands (Chatham, Bounty, Antipodes, Auckland, Campbell, Macquarie, Snares, Stewart, and Solander), and along coast of southwestern Australia from Eclipse Island, Western Australia, to Kangaroo Island, South Australia.

*Arctocephalus tropicalis* (Gray, 1872) (subantarctic fur seal).

Subantarctic islands of Atlantic Ocean and Indian Ocean, north of the Antarctic Convergence (Tristan da Cunha, Gough, Marion, Prince Edward, Crozet (not breeding), Amsterdam, and St. Paul). Rarely wanders to South Africa, New Zealand, and Macquarie Island.

*Arctocephalus australis* (Zimmerman, 1783) (South American fur seal).

Falkland Islands; coasts of South America from Tierra del Fuego north to Rio de Janeiro, Brazil, and Lima, Peru. The proposed subspecific separation of the Falkland Islands and mainland populations is not justifiable (Vaz Ferreira 1976).

*Arctocephalus galapagoensis* Heller, 1904 (Galapagos fur seal).

Confined to the Galápagos Islands.

*Arctocephalus philippii* (Peters, 1866) (Juan Fernandez fur seal).

Now known to breed only on the Islas Juan Fernandez, Chile. Probably bred formerly on Isla San Felix and Isla San Ambrosio, Chile.

*Arctocephalus townsendi* Merriam, 1897 (Guadalupe fur seal).

Now known to breed only on Isla Guadalupe, Baja California. Occasionally seen on other islands off southern California and Baja California (San Miguel, San Nicholas, and Cedros), where it may have bred formerly. An old sealer's report from Isla Socorro (Morrell 1832) and bones from an Indian midden on Monterey Bay (Repenning, pers. commun.) suggest that it originally ranged more widely.

#### Family MUSTELIDAE

This family includes the weasels, badgers, otters, and their allies. Only the otters (subfamily Lutrinae) include marine species and only the sea otter is usually regarded as marine, although several species of river otters of the genus *Lutra*—especially the chungungo, *L. felina* (Molina, 1782), of Chile and Peru—feed extensively in salt water.

#### Genus ENHYDRA Fleming, 1822

*Enhydra lutris* (Linnaeus, 1758) (sea otter).

Formerly ranged from Morro Hermoso, Baja California, north along the coast to Prince William Sound and the south shore of the Alaska Peninsula, throughout the Aleutian, Pribilof, and Commander islands, along the southeast coast of Kamchatka, and through the Kuril Islands to northern Hokkaido. Three subspecies are recognizable (Roest 1973; Davis and Lidicker 1975): *E. l. nereis* (Merriam, 1904) from Mexico to Prince William Sound, Alaska; *E. l. lutris* from the Aleutian and Commander islands; and *E. l. gracilis* (Bechstein, 1799) in Kamchatka and the Kuril Islands. (The Aleutian race has been introduced into Oregon, Washington, British Columbia, and southeastern Alaska, where the indigenous stock was extirpated.)

## Family PHOCIDAE

### Genus PHOCA Linnaeus, 1758

Scheffer (1958) raised *Pusa* (including *hispidus*, *caspicus*, and *sibiricus*), *Pagophilus* (*groenlandicus*), and *Histiophoca* (*fasciata*) to generic rank, but Burns and Fay (1970) have shown that they deserve only subgeneric rank.

*Phoca vitulina* Linnaeus, 1758 (harbor seal).

Shores of North America and Eurasia from Hokkaido, Baja California, North Carolina, and Spain north to the edge of arctic ice. Gives birth on land in May-August; the pup sheds its white coat in utero. The Atlantic subspecies, *P. v. vitulina*, is distinguishable from the Pacific subspecies by skull characters. Western North Atlantic seals are sometimes listed as a separate subspecies from those on the eastern side, but Doutt (1942) could find no differences between seals of the eastern and western Atlantic. *Phoca v. mellonae* Doutt, 1942, is said to be confined to the Seal Lakes complex of the Ungava Peninsula, though Mansfield (1967) doubted the validity of the subspecies. *Phoca v. richardii* (Gray, 1864) occurs in the eastern North Pacific west to the Aleutians, where it intergrades with *P. v. kurilensis* Inukai, 1942, of the western North Pacific (Shaughnessy 1974).

*Phoca largha* Pallas, 1811 (spotted seal; larga seal).

Chukchi, western Beaufort, northern Bering, Okhotsk, and Japan seas, southwest to the Shantung Peninsula, China. Reproductively isolated from *P. vitulina*, with which it is sympatric in the Kuril Islands and along the north shore of the Alaska Peninsula (Belkin 1964; McLaren 1966; Burns 1970). Gives birth on ice in late winter or spring and the pup retains its white coat for a week or more after birth.

*Phoca hispida* Schreber, 1775 (ringed seal).

Throughout the Arctic Ocean and adjacent seas,

chiefly in fast ice, and in several Finnish lakes. Four geographically isolated peripheral populations are fairly well-defined subspecies, one each from the Okhotsk Sea (*P. h. ochotensis* Pallas, 1811); the Baltic Sea (*P. h. botnica* Gmelin, 1788); Lake Ladoga (*P. h. ladogensis* Nordquist, 1899); and Lake Saimaa and adjacent lakes (*P. h. saimensis* Nordquist, 1899). Geographical variation in the Arctic Ocean and Bering Sea populations, tentatively referable to the nominate subspecies, requires much further study.

*Phoca sibirica* Gmelin, 1788 (Baikal seal).

Only in Lake Baikal, U.S.S.R., a freshwater body which freezes in winter.

*Phoca caspica* Gmelin, 1788 (Caspian seal).

Only in Caspian Sea, U.S.S.R., the northern end of which freezes in winter.

*Phoca groenlandica* Erxleben, 1777 (harp seal).

North Atlantic Ocean, in pack ice from northern shores of Europe to eastern Canada. Breeds on pack ice in three main areas: the White Sea, north of Jan Mayen, and Newfoundland. (The Newfoundland seals breed in two centers: the "Front" north of the island and the "Gulf" west of it.) Seals of the three areas differ in size, cranial features, and body coloration (Khuzin 1963, 1967; Yablokov and Sergeant 1963; Yablokov and Etin 1965). Subspecific names have been given to the Newfoundland stock, *P. g. groenlandica*, and that of the White Sea, *P. g. oceanica* Lepechin, 1778, but not to the Jan Mayen stock (Smirnov 1927).

*Phoca fasciata* Zimmermann, 1783 (ribbon seal).

North Pacific Ocean, chiefly in pack ice, from northern Hokkaido and the Okhotsk Sea to northwestern Alaska.

### Genus HALICHOERUS Nilsson, 1820

*Halichoerus grypus* (Fabricius, 1791) (gray seal).

Temperate coasts of the North Atlantic. There are three breeding populations: one in the western Atlantic from Newfoundland to Massachusetts, another in the eastern Atlantic from the British Isles (rarely France) and Iceland to the White Sea, and a third in the Baltic Sea. Seals of the western Atlantic and Baltic populations pup in February and March, seals of the eastern Atlantic from September to December.

### Genus ERIGNATHUS Gill, 1866

*Erignathus barbatus* (Erxleben, 1777) (bearded seal).

Circumboreal at edges of ice; along all coasts and islands of northern Eurasia and northern North America. No subspecies are recognizable (Kosygin and Potelov 1971).

Genus CYSTOPHORA Nilsson, 1820

*Cystophora cristata* (Erxleben, 1777) (hooded seal; bladdernose seal).

North Atlantic Ocean at edges of ice from Novaya Zemlya to eastern Canada. Jan Mayen, Newfoundland, and Davis Strait breeding stocks are perhaps distinct.

Genus MONACHUS Fleming, 1822

*Monachus monachus* (Hermann, 1779) (Mediterranean monk seal).

The original range included the southern and western coasts of the Black Sea, the coasts and islands of the Mediterranean Sea, the coast of northwestern Africa southwestward to Cape Blanc, Mauritania, and the Madeira and Canary islands. Now rare or extirpated throughout much of its range.

*Monachus tropicalis* (Gray, 1850) (Caribbean monk seal; West Indian monk seal).

Extinct (K. W. Kenyon, pers. commun.). In historic times its range included shores and islands of the western Caribbean Sea, the Greater Antilles, the northern Lesser Antilles, the Bahamas, the Yucatan Peninsula, and the Florida Keys. In prehistoric times ranged north to South Carolina.

*Monachus schauinslandi* Matschie, 1905 (Hawaiian monk seal).

Breeds on Leeward Chain of the Hawaiian Islands, from French Frigate Shoals to Kure Atoll; rarely wanders southeastward to Hawaii and south to Johnston Island.

Genus LOBODON Gray, 1844

*Lobodon carcinophagus* (Hombron and Jacquinot, 1842) (crabeater seal).

Crabeaters are circumpolar and abundant in pack ice of the Southern Ocean; they straggle to southern tips of New Zealand, Australia, Tasmania, South Africa, and South America.

Genus OMMATOPHOCA Gray, 1844

*Ommatophoca rossii* Gray, 1844 (bigeye seal; Ross seal).

Circumpolar in pack ice of Antarctic Ocean.

Genus HYDRURGA Gistel, 1848

*Hydrurga leptonyx* (de Blainville, 1820) (leopard seal).

Leopard seals are circumpolar in the Southern Ocean and are recorded from most subantarctic islands, as well as New Zealand, southern Australia, the Cook Islands, southern South America, and South Africa.

Genus LEPTONYCHOTES Gill, 1872

*Leptonychotes weddelli* (Lesson, 1826) (Weddell seal).

Circumpolar in fast ice around Antarctica, south to lat. 80°S in the Bay of Whales; straggling to subantarctic islands and as far north as Uruguay, lat. 35°S.

Genus MIROUNGA Gray, 1827

*Mirounga leonina* (Linnaeus, 1758) (southern elephant seal).

Circumpolar on subantarctic islands, south to edges of ice at lat. 78°S. The southern elephant seal breeds along a continental coast only at Argentina. Three subspecies have been proposed, one from the South American sector of the range, one from the southern Indian Ocean sector, and one from the New Zealand sector (Lydekker 1909). They may be valid, but further study is required before they can be accepted.

*Mirounga angustirostris* (Gill, 1866) (northern elephant seal).

Breeds from the Farallon Islands, Calif., south to Isla Guadalupe and Islas San Benito, Baja California. Formerly from Point Reyes, Calif., south to Cabo San Lázaro, Baja California. Ranges at sea north to southeastern Alaska.

## Order SIRENIA

### Family DUGONGIDAE

Genus DUGONG Lacépède, 1799

*Dugong dugon* (Müller, 1776) (dugong).

In tropical bays and estuaries of the Indian and western Pacific oceans from Lourenço Marques, Mozambique, and the Red Sea, east to the Ryukyu Islands (Amami Oshima), Palau, the Solomon Islands, the New Hebrides, New Caledonia, the Fiji Islands, and northern Australia. Carter et al. (1945) listed it from the Marshall Islands; if it ever occurred there it no longer does so. Now rare in all its range except along northern Australia.

Genus HYDRODAMALIS Retzius, 1794

*Hydrodamalis gigas* (Zimmermann, 1780) (great northern sea cow; Steller's sea cow).

Discovered in the Commander Islands in 1741, the sea cow was exterminated by Russian hunters about 1768. In historic time, it lived only on Bering and Copper islands and its total population probably did not exceed 1,000 or 2,000 animals. A rib was found on Attu, the westernmost Aleutian Island, in 1842 or 1843, by Iliia G. Wosnesenski. "There is no indisputable evidence of its ever having inhabited other coasts than those of the Commander Islands, as the find of a rib on Attu Island does not necessarily



prove that the animal once lived there, though that is not improbable" (Stejneger 1897). A skull fragment about 19,000 yr old was dredged from the sea floor off Monterey, Calif. (Jones 1967). Remains have also been recovered from Pleistocene deposits on Amchitka in the Aleutian Islands (Gard 1972). Recent rumors (Berzin et al. 1963) of living sea cows near Cape Navarin, Siberia, have been discredited.

#### Family TRICHECHIDAE

Genus TRICHECHUS Linnaeus, 1758

Three allopatric species are recognized (Hatt 1934), but their status needs confirmation.

*Trichechus manatus* Linnaeus, 1758 (Caribbean manatee; West Indian manatee).

Two doubtfully valid subspecies have been described: *T. m. manatus* from the sea coast, and lower reaches of rivers, from Bay of Campeche, Mexico, to northeastern South America, and in the Bahamas and the Greater Antilles; and *T. m. latirostris* (Harlan 1824) from the coast and coastal rivers of United States from Beaufort, N.C., to Florida Keys and coast of Gulf of Mexico, westward to mouth of Rio Grande.

*Trichechus senegalensis* Link, 1795 (West African manatee).

Coastal lagoons and the lower reaches of rivers from Sénégal to the Cuanza River, Angola, and in the Niger and Benue drainages of Nigeria.

*Trichechus inunguis* (Natterer, 1883) (Amazon manatee).

Rivers of northeastern South America, particularly the Amazon and Orinoco systems.

### Order MYSTICETI

#### Family ESCHRICHTIIDAE

Genus ESCHRICHTIUS Gray, 1864

*Eschrichtius robustus* (Lilljeborg, 1861) (gray whale).

Shallow coastal waters of the North Pacific. There are two stocks, one on the eastern side from the Gulf of California to the Chukchi and Beaufort seas, another on the western side from Korea and Japan to the Okhotsk Sea; the latter stock is nearly extinct. Formerly in the North Atlantic.

#### Family BALAENOPTERIDAE

Genus BALAENOPTERA Lacépède 1804

*Balaenoptera acutorostrata* Lacépède, 1804 (minke whale).

Widely distributed in all oceans. Three subspecies are recognizable: *B. a. acutorostrata* in the North

Atlantic, *B. a. davidsoni* Scammon, 1872, in the North Pacific, and *B. a. bonaerensis* Burmeister, 1867, in the Southern Hemisphere (Omura 1975). Specimens from Ceylon have also been described as a separate subspecies, *B. a. thalmaha* Deraniyagala, 1963, but its validity requires confirmation.

*Balaenoptera edeni* Anderson, 1878 (Bryde whale).

Tropical and warm temperate waters of the Atlantic, Indian, and Pacific oceans.

*Balaenoptera borealis* Lesson, 1828 (sei whale).

All oceans except tropical and polar seas. Two subspecies distinguished: a smaller one, *B. b. borealis*, in the Northern Hemisphere and a larger one, *B. b. schlegellii* Flower, 1865, in the Southern Hemisphere.

*Balaenoptera physalus* (Linnaeus, 1758) (fin whale).

All oceans, but rarely in tropical waters or among pack ice. Two subspecies are recognized: a smaller Northern Hemisphere form, *B. p. physalus*, and a larger Southern Hemisphere form, *B. p. quoyi* (Fischer, 1829).

*Balaenoptera musculus* (Linnaeus, 1758) (blue whale).

All oceans. Three subspecies are recognized: a small one, *B. m. musculus*, in the North Atlantic and North Pacific; a large one, *B. m. intermedia* Burmeister, 1871, that spends the summer in Antarctic waters; and a pygmy subspecies, *B. m. breviceauda* Ichihara, 1966, in the southern Indian Ocean. (The name *B. m. breviceauda*, published by Zemsky and Boronin 1964, is a nomen nudum according to van Bree, pers. commun.; the first valid publication of the name was by Ichihara 1966.) The taxonomic status of blue whales off the coasts of Chile and Peru and in the northern Indian Ocean is not settled.

Genus MEGAPTERA Gray, 1846

*Megaptera novaeangliae* (Borowski, 1781) (humpback whale).

Nearly worldwide; winters largely in tropical waters near islands or the coast, summers in temperate and subpolar waters. This species shows little or no geographical variation in size; the several discrete populations differ in the frequency of color variations.

#### Family BALAENIDAE

Genus BALAENA Linnaeus, 1758

Some authors place *glacialis* in a separate genus, *Eubalaena* Gray, 1864. Eschricht and Reinhardt (1861), the only authors who have made a detailed comparison between *mysticetus* and *glacialis*, regarded them as congeneric. Gray's generic name was ignored by virtually all subsequent authors until it was resurrected by Allen

(1908). Gray and Allen were two of the most notorious generic "splitters" in the history of mammalogy (cf. Simpson 1961:139). The differences between the two species are no greater than those separating, e.g., the various species of *Balaenoptera*.

*Balaena glacialis* Müller, 1776 (right whale; black right whale).

Temperate waters of the North Atlantic, the North Pacific, and the Southern Hemisphere. The southern populations are distinguishable as a separate subspecies (or species, according to some authors), *B. g. australis* Desmoulins, 1822, from *B. g. glacialis* of the North Atlantic (Muller 1954); North Pacific populations are apparently identical to those of the North Atlantic (Omura et al. 1969).

*Balaena mysticetus* Linnaeus, 1758 (bowhead whale; Arctic right whale).

Arctic waters. There are four geographically isolated populations: 1) From Spitzbergen west to east Greenland; 2) in Davis Strait, Baffin Bay, James Bay, and adjacent waters; 3) in the Bering, Chukchi, and Beaufort seas; and 4) in the Okhotsk Sea. The Alaskan Eskimo recognize two kinds: the larger "kairalik" or true bowhead, and the smaller "ingotok" (known as the "poggy" to the 19th century American whalers). I believe that the ingotok is most likely a young bowhead.

Genus CAPEREA Gray, 1864

*Caperea marginata* (Gray, 1846) (pygmy right whale).

Temperate waters of Southern Ocean; known mostly from strandings on New Zealand, Australia, South America, the Falkland Islands, and South Africa.

## Order ODONTOCETI

In recent years, several new family classifications of the odontocetes have been proposed (Fraser and Purves 1960; Nishiwaki 1963; Kasuya 1973). I believe that insufficient evidence has been published to support the validity of these classifications and that much more study is needed before any changes will be generally accepted. I therefore follow tradition in regarding the living odontocetes as divisible into five families.

Family PLATANISTIDAE

This family includes three well-marked subfamilies (which are sometimes accorded family rank): Platanistinae (*Platanista*), Iniinae (*Inia*, *Lipotes*), and Pontoporiinae (*Pontoporia*).

Genus INIA d'Orbigny, 1834

*Inia geoffrensis* (de Blainville, 1817) (boto; Amazon dolphin).

Amazon and Orinoco basins of South America. *Inia g. boliviensis* d'Orbigny, 1834, of the upper Madeira River system in Bolivia differs considerably from *I. g. geoffrensis* in the remainder of the Amazon basin and may be a distinct species (van Bree and Robineau 1973; van Bree, pers. commun.). The population in the Orinoco basin may also be sub-specifically distinct.

Genus LIPOTES Miller, 1918

*Lipotes vexillifer* Miller, 1918 (pei c'hi; whitefin dolphin).

Lower Yangtze River from its mouth (Shanghai) upstream to Tung-t'ing Hu (Lake) and its tributaries. The English name "whiteflag" dolphin is based on an erroneous interpretation of the Chinese (M. Nishiwaki, pers. commun.).

Genus PONTOPORIA Gray, 1846

*Pontoporia blainvillei* (Gervais and d'Orbigny, 1844) (franciscana; La Plata dolphin).

Coastal waters and estuaries of eastern South America, from Baia de Santos, Brazil, to Golfo San Matias, Argentina.

Genus PLATANISTA Wagler, 1830

The two allopatric forms of this genus are regarded as subspecifically distinct by Kasuya (1972) and as specifically distinct by Pilleri and Gehr (1971). I tentatively list them as separate species on the advice of van Bree (pers. commun.).

*Platanista gangetica* (Roxburgh, 1801) (Ganges susu; Ganges dolphin).

Ganges-Brahmaputra-Meghna river system of India, Bangladesh, and Nepal, from tidal limits to the foothills. It is presumably this species that formerly occurred in the Karnaphuli River.

*Platanista minor* Owen, 1853 (Indus susu; Indus dolphin).

Indus River system of Pakistan and India, from tidal limits to the foothills. (Most authors have used the name *P. indi* Blyth, 1859, but Owen's name has priority.)

Family DELPHINIDAE

Basic references on the classification of the Delphinidae are Flower (1883) and True (1889). The true porpoises (*Phocoena*, *Neophocaena*, and *Phocoenoides*) constitute a well-marked group that is sometimes accorded family rank (Phocoenidae). The remainder of the family has been variously subdivided or divided.

Genus STENO Gray, 1846

*Steno bredanensis* (Lesson, 1828) (rough-toothed dolphin).

All tropical and warm temperate seas.

Genus SOUSA Gray, 1866

*Sousa chinensis* (Osbeck, 1765) (Indo-Pacific humpback dolphin; white dolphin).

Coastal waters of the Indian and western Pacific oceans, from Port Elizabeth, South Africa, north to the Red Sea and east to southern China (including lower reaches of the Yangtze, Foochow, and Canton rivers), Borneo, and northeastern and eastern Australia. The taxonomy of the humpbacked dolphins is greatly in need of revision. Several nominal species have been described, but individual, sexual, age, and geographic variations have not been adequately studied.

*Sousa teuszii* (Kükenthal, 1892) (Atlantic humpback dolphin).

Coastal waters of west Africa from Mauritania to Cameroon. This form could perhaps be regarded as a subspecies of *S. chinensis*, from which it differs mainly in tooth count (26-29 versus 32-36).

Genus SOTALIA Gray, 1866

Further taxonomic studies of this genus are needed. Two forms are recognized—one chiefly in coastal and estuarine waters, the other in fresh water. The differences between them are slight; they are best regarded as subspecies of a single species. (The Old World species formerly placed in *Sotalia* are now placed in a separate genus, *Sousa*.)

*Sotalia fluviatilis* (Gervais, 1853) (tucuxi; tookashee).

*Sotalia f. fluviatilis* occurs in the Amazon River and its tributaries; *S. f. guianensis* (van Beneden, 1864) occurs in coastal waters and lower reaches of rivers of northeastern South America, from Lake Maracaibo, Venezuela, to Santos, Brazil.

Genus TURSIOPS Gervais, 1855

*Tursiops truncatus* (Montagu, 1821) (bottlenose dolphin).

Widely distributed in temperate and tropical waters, including the Black Sea. Occurs mostly close to shore, near islands, and over shallow banks. Ranges north to Japan, Hawaii, California, New Brunswick, and Norway; south to southern Australia, New Zealand, Chile, Argentina, and South Africa. There is considerable geographical variation; populations in warmer waters of the Indo-Pacific tend to average smaller in body size and greater in snout length and tooth count.

Genus STENELLA Gray, 1866

The species of this genus fall into three groups or superspecies which are sympatric in many areas: 1) The spinner dolphins with long snouts, about 50 teeth in each

jaw, and shallow palatal grooves. This group includes forms that differ markedly in body form and color pattern, but intermediate populations connect the extreme forms, so they are regarded as conspecific (Perrin 1972, 1975a, b). Current studies indicate, however, that there is probably another species in the Atlantic similar to, but sympatric with, the spinner dolphin (Perrin, pers. commun.). 2) The spotted and bridled dolphins with shorter snouts, about 37 teeth in each jaw, and no palatal grooves. This group apparently includes two species, the ranges of which overlap in the Atlantic (Perrin 1975b). 3) The striped dolphins, with about 44 teeth in each jaw, no palatal grooves, a dark stripe along the flank, and no spots. Only one species is recognized (Fraser and Noble 1970).

*Stenella longirostris* (Gray, 1828) (spinner dolphin).

Tropical Atlantic, Indian, and Pacific oceans. This species shows marked geographical variation. In the central and eastern Pacific, Perrin (1972, 1975a, b) has described four subspecies: "Costa Rican," "eastern," "whitebelly," and "Hawaiian"—the correct scientific names of which have not been determined. [Although most recent authors have used Gray's specific name, its applicability is questionable (Perrin 1975b).]

*Stenella attenuata* (Gray, 1846) (bridled dolphin; pantropical spotted dolphin).

Tropical Atlantic, Indian, and Pacific oceans. This species also shows considerable geographical variation. In the central and eastern Pacific, Perrin (1970, 1975a, b) has described three subspecies: "coastal" *S. a graffmani* (Lönnerberg, 1934), "offshore," and "Hawaiian"; the correct scientific names of the latter two have not been determined. [For the spotted dolphins I provisionally use the specific names employed by most recent authors, although their validity is dubious. Other available names are *S. dubia* (G. Cuvier, 1812) and *S. frontalis* (G. Cuvier, 1829).]

*Stenella plagiodon* (Cope, 1866) (spotted dolphin; Atlantic spotted dolphin).

Tropical and warm temperate waters of Atlantic Ocean. (See nomenclatorial note under preceding species).

*Stenella coeruleoalba* (Meyen, 1833) (striped dolphin).

Widely distributed in temperate and tropical waters around the world.

Genus DELPHINUS Linnaeus, 1758

*Delphinus delphis* Linnaeus, 1758 (saddleback dolphin).

Widely distributed in warm temperate and tropical waters of all oceans, including the Black Sea. There is marked geographical variation in snout length and

other features; the extremely longsnouted Arabian Sea form, *D. d. tropicalis* van Bree, 1971, may be a distinct species (van Bree and Purves 1972).

Genus LAGENODELPHIS Fraser, 1956

*Lagenodelphis hosei* Fraser, 1956 (shortsnouted whitebelly dolphin).

Tropical and warm temperate waters of Indian and Pacific oceans. Recorded from Natal, Japan, Taiwan, Philippines, Borneo, northeastern Australia, and the central and eastern tropical Pacific.

Genus LAGENORHYNCHUS Gray, 1846

This genus contains two well-defined species in the North Atlantic and another in the North Pacific. Bierman and Slijper (1947, 1948) regarded all Southern Hemisphere forms as conspecific, but Fraser (1966) showed that there are three distinct species. The species *electra* is now placed in a separate genus, *Peponocephala*.

*Lagenorhynchus albirostris* (Gray, 1846) (whitebeak dolphin).

North Atlantic from Davis Strait and Newfoundland east to the Barents Sea and North Sea (rarely to the southern British Isles).

*Lagenorhynchus acutus* (Gray, 1828) (Atlantic whiteside dolphin).

North Atlantic from Massachusetts and southern Greenland east to western Norway and the British Isles.

*Lagenorhynchus obliquidens* Gill, 1865 (Pacific whiteside dolphin; hookfin dolphin).

Waters off the coast of North America from southeastern Alaska to Baja California, and off the coast of Asia from the Kuril Islands to Japan.

*Lagenorhynchus obscurus* (Gray, 1828) (dusky dolphin).

Temperate waters off South America, South Africa, Kerguelen Island, southern Australia, and New Zealand. Primarily a coastal species.

*Lagenorhynchus australis* (Peale, 1848) (blackchin dolphin).

Temperate waters off southern South America and the Falkland Islands.

*Lagenorhynchus cruciger* (Quoy and Gaimard, 1824) (hourglass dolphin).

Temperate waters of the Southern Ocean. A pelagic species, found chiefly in waters immediately north of the Antarctic Convergence.

Genus CEPHALORHYNCHUS Gray, 1846

Four species are recognized (Harmer 1922).

*Cephalorhynchus commersonii* (Lacépède, 1804) (piebald dolphin; Jacobite).

Atlantic coast of South America from Golfo San Matias to Tierra del Fuego; Falkland Islands; South Georgia; and Kerguelen Island.

*Cephalorhynchus eutropia* (Gray, 1846) (black dolphin; Chilean dolphin).

Coast of Chile between lat. 37° and 55°S.

*Cephalorhynchus heavisidii* (Gray, 1828).

Coastal waters from Cape of Good Hope north to Cape Cross, South West Africa (P. B. Best, pers. commun.).

*Cephalorhynchus hectori* (van Beneden, 1881) (piebald dolphin; whitefront dolphin).

Coastal waters of New Zealand.

Genus LISSODELPHIS Gloger, 1841

The two species of this genus differ, as far as is known, mainly in color pattern, and they should perhaps be regarded as subspecies of a single species.

*Lissodelphis borealis* (Peale, 1848) (northern right-whale dolphin).

Temperate waters of the North Pacific from Japan and the Kurils to British Columbia and California. Individuals from Japan with a variant color pattern have been named *L. b. albiventris* Nishiwaki, 1972.

*Lissodelphis peronii* (Lacépède, 1804) (southern right-whale dolphin).

Temperate waters of the Southern Ocean.

Genus GRAMPUS Gray, 1828

*Grampus griseus* (G. Cuvier, 1812) (whitehead grampus; gray grampus).

All temperate and tropical seas.

Genus PEPONOCEPHALA Nishiwaki and Norris, 1966

*Peponocephala electra* (Gray, 1846) (little blackfish; many-toothed blackfish; melon-head blackfish).

Tropical Atlantic, Indian, and Pacific oceans.

Genus FERESA Gray, 1871

*Feresa attenuata* Gray, 1875 (pygmy killer whale).

Tropical and warm temperate waters of the Atlantic, Indian, and Pacific oceans.

Genus PSEUDORCA Reinhardt, 1862

*Pseudorca crassidens* (Owen, 1846) (false killer whale).

All temperate and tropical seas.

Genus GLOBICEPHALA Lesson, 1828

There appear to be two well-defined species, the ranges of which overlap off the middle Atlantic coast of the United States, off southern Europe, off South Africa, and perhaps elsewhere (van Bree 1971).

*Globicephala melaena* (Traill, 1809) (longfin pilot whale).

Nominate subspecies in the cool temperate North Atlantic Ocean; *G. m. edwardii* A. Smith, 1834, throughout cool temperate waters of the Southern Hemisphere. The validity of the subspecific distinction is questionable. Occurred in the North Pacific (near Japan) until at least the 10th century A.D. (Kasuya 1975).

*Globicephala macrorhynchus* Gray, 1846 (shortfin pilot whale).

Tropical and warm temperate waters of the Atlantic, Indian, and Pacific oceans. *Globicephala sieboldii* Gray, 1846, of the North Pacific is conspecific with *G. macrorhynchus*, although it may be recognizable at the subspecific level. (The specific name is usually spelled *macrorhyncha*, but it is a noun in apposition, not an adjective, so must retain its original gender.)

Genus ORCINUS Fitzinger, 1860

*Orcinus orca* (Linnaeus, 1758) (killer whale).

All oceans, chiefly in coastal waters and cooler regions.

Genus ORCAELLA Gray, 1866

*Orcaella brevirostris* (Gray, 1866) (Irrawaddy dolphin; lumbalumba).

Coastal waters from the Bay of Bengal east to New Guinea and northern Australia; ascends far up the Mekong, Irrawaddy, Ganges, and other rivers.

Genus PHOCOENA G. Cuvier, 1817

The genus includes four species (Norris and McFarland 1958; Noble and Fraser 1971).

*Phocoena phocoena* (Linnaeus, 1758) (harbor porpoise).

Coastal waters of the North Atlantic from Delaware and Sénégal north to Davis Strait, Iceland, and the White Sea; coastal waters of the North Pacific from Japan and Baja California north to Point Barrow, Alaska. An isolated population in the Black Sea has been named *P. p. relicta* Abel, 1905.

*Phocoena sinus* Norris and McFarland, 1958 (vaquita; cochito; Gulf of California porpoise).

Upper Gulf of California; sight records farther south are questionable.

*Phocoena dioptrica* Lahille, 1912 (spectacled porpoise).

Coast of Argentina and Uruguay; the Falkland Islands; and South Georgia.

*Phocoena spinipinnis* Burmeister, 1865 (black porpoise).

East coast of South America from Uruguay to Patagonia; west coast from Paita, Peru, to Valdivia, Chile.

Genus NEOPHOCAENA Palmer, 1899

*Neophocaena phocaenoides* (G. Cuvier, 1829) (finless porpoise).

Warm coastal waters and certain rivers from Pakistan east to Korea, Japan, Borneo, and Java. (The type-specimen allegedly came from the Cape of Good Hope, but Peter Best (pers. commun.) stated that there are no indisputable South African records.) Specimens from China and Japan differ from Indian Ocean specimens, and are best regarded as a subspecies, *N. p. asiaeorientalis* (Pilleri and Gühr, 1971), rather than a full species as originally described (van Bree 1973). More recently Pilleri and Gühr (1975) have differentiated three allopatric forms which they arbitrarily rank as species: *N. phocaenoides* from Pakistan to Borneo, *N. asiaeorientalis* from China, and *N. sunameri* Pilleri and Gühr, 1975, from Japan and Korea. Their published data are inadequate to reveal whether the Japanese and Chinese populations are sufficiently separable to warrant recognition of *sunameri* even as a subspecies.

Genus PHOCOENOIDES Andrews, 1911

*Phocoenoides dallii* (True, 1885) (Dall porpoise; whiteflank porpoise).

Immediate offshore waters of the North Pacific from Japan and southern California north to the southern Bering Sea. "True's porpoise" is a color phase localized in Japanese waters.

Family MONODONTIDAE

Some authors have included the narwhal and beluga in the family Delphinidae.

Genus DELPHINAPTERUS Lacépède, 1804

*Delphinapterus leucas* (Pallas, 1776) (beluga; belukha; white whale).

Arctic Ocean and adjacent seas; including Okhotsk and Bering Seas, and James Bay; isolated populations occur in Cook Inlet, Alaska, and the Gulf of St. Lawrence; ascends several hundred miles up larger rivers of Siberia and Alaska. Some authors recognize three subspecies: a large one, *D. l. dorofeevi* Barabash and Klumov, 1935, from the

Okhotsk Sea; a small one, *D. l. marisalbi* Ostroumov, 1935, in the Barents and White seas; and a medium-sized one, *D. l. leucas*, in the rest of the range. However, geographical variation is more complex than this classification suggests (Sergeant and Brodie 1969).

Genus MONODON Linnaeus, 1758

*Monodon monoceros* Linnaeus, 1758 (Narwhal).  
North polar seas, mainly in deep waters.

Family PHYSETERIDAE

The genus *Kogia* is sometimes placed in a separate family (Kogiidae).

Genus PHYSETER Linnaeus, 1758

*Physeter macrocephalus* Linnaeus, 1758 (sperm whale).

All oceans (except polar ice fields). (For use of this name instead of *P. catodon* Linnaeus, 1758, see Husson and Holthuis 1974.)

Genus KOGIA Gray, 1846

Handley (1966) has reviewed the distinguishing features of the two species in this genus.

*Kogia breviceps* (de Blainville, 1838) (pygmy sperm whale).

World-wide in tropical and warm temperate waters.

*Kogia simus* Owen, 1866 (dwarf sperm whale).

The seas adjacent to South Africa, India, Ceylon, Japan, Hawaii, California, Baja California, and eastern United States.

Family ZIPHIIDAE

See Moore (1968) for diagnoses of the genera.

Genus BERARDIUS Duvernoy, 1851

Two allopatric species are recognized. The North Pacific form differs from the Southern Hemisphere form chiefly by its much larger size. Possibly it should be regarded as only a subspecies of the Southern Hemisphere form.

*Berardius arnuxii* Duvernoy, 1851 (southern giant bottlenose whale).

Southern Ocean; known from South Australia, New Zealand, Argentina, Falkland Islands, South Georgia, South Shetlands, South Africa, and off the Antarctic Peninsula.

*Berardius bairdii* Stejneger, 1883 (North Pacific giant bottlenose whale).

North Pacific from Japan and southern California north to the Bering Sea.

Genus ZIPHIUS G. Cuvier, 1823

*Ziphius cavirostris* G. Cuvier, 1823 (goosebeak whale).  
All temperate and tropical seas.

Genus TASMASETUS Oliver, 1937

*Tasmacetus shepherdi* Oliver, 1937.

Known only from a few specimens stranded in New Zealand, Chile, and Argentina.

Genus INDOPACETUS Moore, 1968

The one species of this genus was formerly included in *Mesoplodon*.

*Indopacetus pacificus* (Longman, 1926) (Indo-Pacific beaked whale).

Known from only two specimens stranded at Mackay, Queensland, Australia, and Danane, Somalia.

Genus HYPEROODON Lacépède, 1804

Two well-defined species are recognized: one in the North Atlantic, the other in the Southern Hemisphere. The latter constitutes subgenus *Frasercetus* Moore, 1968. The occurrence of *Hyperoodon* in the North Pacific has never been verified, and most if not all published records of its occurrence there are based on misidentification of *Berardius*. Beaked whales possibly referable to *Hyperoodon* are taken by whalers off the Okhotsk Sea coast of Hokkaido, but to date none has been examined by a biologist (M. Nishiwaki, pers. commun.).

*Hyperoodon ampullatus* (Forster, 1770) (North Atlantic bottlenose whale).

North Atlantic from Davis Strait and Novaya Zemlya south to Rhode Island and the English Channel; doubtfully recorded from the Mediterranean Sea.

*Hyperoodon planifrons* Flower, 1882 (flathead bottlenose whale).

Southern Ocean; known from Australia, New Zealand, Argentina, the Falkland Islands, South Georgia, the South Orkney Islands, South Africa, and off the coast of Antarctica in the Pacific and Indian Ocean sectors.

Genus MESOPLONDON Gervais, 1850

Eleven species are currently recognized by Moore (1968). *Mesoplodon pacificus* is now placed in a separate genus, *Indopacetus*. *Mesoplodon layardii* is placed in the subgenus *Dolichodon* Gray, 1871; *M. densirostris* in the subgenus *Dioplodon* Gervais, 1850; and the remaining species are placed in subgenus *Mesoplodon*.

*Mesoplodon hectori* (Gray, 1871).

Known only from Tasmania, New Zealand, the Falkland Islands, and South Africa.

*Mesoplodon mirus* True, 1913.

North Atlantic from Florida and Nova Scotia east to the British Isles; an apparently isolated population in temperate waters off South Africa.

*Mesoplodon europaeus* (Gervais, 1855) (Antillean beaked whale; Gulf Stream beaked whale).

Western North Atlantic from Trinidad, Jamaica, and the Gulf of Mexico, to Long Island, N.Y.; one record from the English Channel.

*Mesoplodon ginkgodens* Nishiwaki and Kamiya, 1958 (ginkgo-tooth whale).

Recorded from Ceylon, Taiwan, Japan, and California.

*Mesoplodon grayi* von Haast, 1876 (scamperdown whale).

South Africa, South Australia, New Zealand, Chatham Islands, and Argentina; one record from the Netherlands.

*Mesoplodon carlhubbsi* Moore, 1963 (archbeak whale).

Temperate waters of the North Pacific from Japan east to British Columbia and California.

*Mesoplodon bowdoini* Andrews, 1908 (deepcrest whale).

Known only from New Zealand, Tasmania, Western Australia, Victoria, and Kerguelen Island.

*Mesoplodon stejnegeri* True, 1885 (sabertooth whale; Bering Sea beaked whale).

Subarctic waters of the North Pacific from the Bering Sea south to Japan and Oregon.

*Mesoplodon bidens* (Sowerby, 1804) (North Sea beaked whale).

Cool temperate waters of the North Atlantic from Newfoundland and Massachusetts east to southern Norway and the Bay of Biscay.

*Mesoplodon layardii* (Gray, 1865) (straptooth whale).

South Africa, southern Australia, New Zealand, and the Falkland Islands.

*Mesoplodon densirostris* (de Blainville, 1817) (dense-beak whale; tropical beaked whale).

Tropical and warm temperate waters of all oceans.

## SYNONYMS

Listed below are generic and specific synonyms frequently appearing in recent literature.

## In recent literature

*Arctocephalus doriferus*

*A. elegans*

*A. tasmanicus*

*Arctophoca*

*Balaena australis*

*B. japonica*

*B. sieboldii*

*Balaenoptera bonaerensis*

*B. brydei*

*B. davidsoni*

*B. huttoni*

*Callorhinus alascensis*

*C. curilensis*

*C. cynocephalus*

*C. mimicus*

*Cephalorhynchus*

*albifrons*

*C. albiventris*

*Delphinapterus dorofeevi*

*D. friemani*

*Delphinus bairdii*

*D. capensis*

*D. dussumieri*

*D. longirostris*

*D. roseiventris*

*D. tropicalis*

*Electra*

*Eschrichtius gibbosus*

*E. glaucus*

*Eubalaena*

*Eumetopias stellerii*

*Feresa intermedia*

*F. occulta*

*Globicephala brachyptera*

*G. edwardii*

*G. leucosagmaphora*

*G. scammonii*

*G. sieboldii*

*Grampidelphis*

*Grampus orca*

*G. rectipinna*

*Gypsophoca*

*Histriophoca*

*Hydrodamalis stelleri*

*Hyperoodon rostratus*

*Lagenorhynchus electra*

*L. fitzroyi*

*L. ognevi*

*L. superciliosus*

*L. thicola*

*L. wilsoni*

*Megaptera nodosa*

*Meomeris*

*Mesoplodon gervaisi*

*M. hotaula*

*M. pacificus*

## In present list

*Arctocephalus pusillus*

*A. tropicalis*

*A. pusillus*

*Arctocephalus*

*Balaena glacialis*

*B. glacialis*

*B. glacialis*

*Balaenoptera acutorostrata*

*B. edeni*

*B. acutorostrata*

*B. acutorostrata*

*Callorhinus ursinus*

*C. ursinus*

*C. ursinus*

*C. ursinus*

*Cephalorhynchus*

*hectori*

*C. eutropia*

*Delphinapterus leucas*

*D. leucas*

*Delphinus delphis*

*D. delphis*

*D. delphis*

*D. delphis*

*Stenella longirostris*

*Delphinus delphis*

*Peponocephala*

*Eschrichtius robustus*

*E. robustus*

*Balaena*

*Eumetopias jubatus*

*Feresa attenuata*

*F. attenuata*

*Globicephala*

*macrorhynchus*

*G. melaena*

*G. melaena*

*G. macrorhynchus*

*G. macrorhynchus*

*Grampus*

*Orcinus orca*

*O. orca*

*Arctocephalus*

*Phoca*

*Hydrodamalis gigas*

*Hyperoodon ampullatus*

*Peponocephala electra*

*Lagenorhynchus obscurus*

*L. obliquidens*

*L. obscurus*

*Lagenorhynchus* sp.?

*L. cruciger*

*Megaptera novaeangliae*

*Neophocaena*

*Mesoplodon europaeus*

*M. ginkgodens*

*Indopacetus pacificus*

*Monachus albiventer*  
*Neobalaena*  
*Neomeris*  
*Neophoca hookeri*  
*N. lobatus*  
*Nodus*  
*Odobenus divergens*  
*Orcaella fluminalis*  
*Orcella*  
*Orcinus rectipinna*  
*Otaria byronia*  
*Pagophilus*  
*Phoca insularis*  
*P. kurilensis*  
*P. richardii*  
*Phocoena vomerina*  
*Phocoenoides truei*  
*Physeter catodon*  
*Platanista indi*  
*Prodelphinus*  
*Pusa*  
*Rhachianectes*  
*Rhytina*  
*Sibbaldus*  
*Sotalia borneensis*  
*S. brasiliensis*  
*S. chinensis*  
*S. gadamu*  
*S. guianensis*  
*S. lentiginosa*  
*S. pallida*  
*S. plumbea*  
*S. sinensis*  
*S. teuszii*  
*S. tucuxi*  
*Sousa borneensis*  
*S. lentiginosa*  
*S. plumbea*  
*S. queenslandensis*  
*Stenella alope*  
*S. dubia*  
*S. euphrosyne*  
*S. frontalis*  
*S. graffmani*  
*S. malayana*  
  
*S. microps*  
*S. pernettensis*  
*S. roseiventris*  
*S. styx*  
*Steno rostratus*  
*Stenodelphis*  
*Stenorhynchus*  
*Stenorhynchus*  
*Susu*  
*Thalarectos*  
*Trichechus latirostris*  
*Tursiops abusalum*  
*T. aduncus*  
*T. catalania*

*Monachus monachus*  
*Caperea*  
*Neophocaena*  
*Phocarctos hookeri*  
*Neophoca cinerea*  
*Mesoplodon*  
*Odobenus rosmarus*  
*Orcaella brevirostris*  
*Orcaella*  
*Orcinus orca*  
*Otaria flavescens*  
*Phoca*  
*Phoca vitulina*  
*P. vitulina*  
*P. vitulina*  
*Phocoena phocoena*  
*Phocoenoides dallii*  
*Physeter macrocephalus*  
*Platanista minor*  
*Stenella*  
*Phoca*  
*Eschrichtius*  
*Hydrodamalis*  
*Balaenoptera*  
*Sousa chinensis*  
*Sotalia fluviatilis*  
*Sousa chinensis*  
*Tursiops truncatus*  
*Sotalia fluviatilis*  
*Sousa chinensis*  
*Sotalia fluviatilis*  
*Sousa chinensis*  
*S. chinensis*  
*S. teuszii*  
*Sotalia fluviatilis*  
*Sousa chinensis*  
*S. chinensis*  
*S. chinensis*  
*S. chinensis*  
*Stenella longirostris*  
*Stenella (?) attenuata*  
*S. coeruleoalba*  
*Stenella (?) attenuata*  
*S. attenuata*  
*Sousa chinensis* (fide  
van Bree, pers. commun.)  
*Stenella longirostris*  
*Stenella (?) plagiodon*  
*S. longirostris*  
*S. coeruleoalba*  
*Steno bredanensis*  
*Pontoporia*  
*Hydrurga*  
*Hydrurga*  
*Platanista*  
*Ursus*  
*Trichechus manatus*  
*Tursiops truncatus*  
*T. truncatus*  
*T. truncatus*

*T. gadamu*  
*T. gillii*  
*T. nesarnack*  
*T. nuuanu*  
*Zalophus cinereus*  
*Z. japonicus*  
*Z. lobatus*  
*Z. wollebaeki*  
  
*T. truncatus*  
*T. truncatus*  
*T. truncatus*  
*T. truncatus*  
*Neophoca cinerea*  
*Zalophus californianus*  
*Neophoca cinerea*  
*Zalophus californianus*

## ACKNOWLEDGMENTS

Victor B. Scheffer, coauthor of the previous two editions, reviewed the entire manuscript. P. J. H. van Bree and C. A. Repenning reviewed in detail the sections on cetaceans and pinnipeds, respectively. P. B. Best, W. F. Perrin, and P. D. Shaughnessy provided valuable comments on portions of the manuscript.

## LITERATURE CITED

- ALLEN, J. A.  
1908. The North Atlantic right whale and its near allies. Bull. Am. Mus. Nat. Hist. 24:277-329.
- AMERICAN ORNITHOLOGISTS' UNION, COMMITTEE ON CLASSIFICATION AND NOMENCLATURE.  
1973. Thirty-second supplement to the American Ornithologists' Union check-list of North American birds. Auk 90:411-419.
- ANDERSON, S., and J. K. JONES, JR. (editors).  
1967. Recent mammals of the world; a synopsis of families. Ronald Press, N.Y., 453 p.
- BAILEY, R. M., J. E. FITCH, E. S. HERALD, E. A. LACHNER, C. C. LINDSEY, C. R. ROBINS, and W. B. SCOTT.  
1970. A list of common and scientific names of fishes from the United States and Canada. Am. Fish. Soc. Spec. Publ. 6, 150 p.
- BELKIN, A. N.  
1964. A new species of seal, *Phoca insularis* sp. n., from the Kuril Islands. Dokl. Akad. Nauk SSSR 158:1217-1219. (In Russ. Transl. seen, Dokl. Biol. Sci., 1965, 158:756-758.)
- BERZIN, A. A., E. A. TIKHOMIROV, and V. I. TROININ.  
1963. Izchezla li Stellerova korova? (Is the Steller sea cow really extinct?) [In Russ.] Priroda 52(8):73-75.
- BIERMAN, W. H., and E. J. SLJPER.  
1947. Remarks upon the species of the genus *Lagenorhynchus*. I. Kon. Ned. Akad. Wetensch., Amsterdam, Afd. Natuur. 50:1353-1364.  
1948. Remarks upon the species of the genus *Lagenorhynchus*. II. Kon. Ned. Akad. Wetensch., Amsterdam, Afd. Natuur. 51:127-133.
- BURNS, J. J.  
1970. Remarks on the distribution and natural history of pagophilic pinnipeds in the Bering and Chukchi Seas. J. Mammal. 51:445-454.
- BURNS, J. J., and F. H. FAY.  
1970. Comparative morphology of the skull of the ribbon seal, *Histiophoca fasciata*, with remarks on systematics of Phocidae. J. Zool. 161:363-394.
- CARTER, T. D., J. E. HILL, and G. H. H. TATE.  
1945. Mammals of the Pacific world. Macmillan Co., N.Y., 227 p.
- DAVIS, J., and W. Z. LIDICKER, JR.  
1975. The taxonomic status of the southern sea otter. Proc. Calif. Acad. Sci., Ser. 4, 40:429-437.
- DOUTT, J. K.  
1942. A review of the genus *Phoca*. Ann. Carnegie Mus. 29:61-125.
- ESCHRICHT, D. F., and J. REINHARDT.  
1861. Om Nordhvalen (*Balaena mysticetus* L.). Vidensk. Selsk. Skr., 5 Raekke, Naturvidensk. Mathem. Afd. 5:433-592.
- FLOWER, W. H.  
1883. On the characters and divisions of the family Delphinidae. Proc. Zool. Soc. Lond. 1883:466-513.



- FRASER, F. C.  
1966. Comments on the Delphinoidea. In K. S. Norris (editor), Whales, dolphins, and porpoises, p. 7-31. Univ. Calif. Press, Berkeley and Los Ang.
- FRASER, F. C., and B. A. NOBLE.  
1970. Variation of pigmentation pattern in Meyen's dolphin, *Stenella coeruleoalba* (Meyen). In G. Pilleri (editor), Investigations on Cetacea, Vol. II, p. 147-163. Inst. Brain Anat., Berne, Switz.
- FRASER, F. C., and P. E. PURVES.  
1960. Hearing in cetaceans. Evolution of the accessory air sacs and the structure and function of the outer and middle ear in Recent cetaceans. Bull. Br. Mus. (Nat. Hist.) Zool. 7:1-140.
- GARD, L. M., JR., G. E. LEWIS, and F. C. WHITMORE, JR.  
1972. Steller's sea cow in Pleistocene interglacial beach deposits on Amchitka, Aleutian Islands. Geol. Soc. Am., Bull. 83:867-870.
- HANDLEY, C. O., JR.  
1966. A synopsis of the genus *Kogia* (pygmy sperm whales). In K. S. Norris (editor), Whales, dolphins, and porpoises, p. 62-69. Univ. Calif. Press, Berkeley and Los Ang.
- HARMER, S. F.  
1922. On Commerson's dolphin and other species of *Cephalorhynchus*. Proc. Zool. Soc. Lond. 1922:627-638.
- HATT, R. T.  
1934. A manatee collected by the American Museum Congo Expedition, with observations on the Recent manatees. Bull. Am. Mus. Nat. Hist. 66:533-566.
- HENDEY, Q. B.  
1972. A Pliocene ursid from South Africa. Ann. S. Afr. Mus. 59: 115-132.
- HERSHKOVITZ, P.  
1966. Catalog of living whales. U.S. Natl. Mus. Bull. 246, 259 p.
- HUSSON, A. M., and L. B. HOLTHUIS.  
1974. *Physeter macrocephalus* Linnaeus, 1758, the valid name for the sperm whale. Zool. Meded. 48:205-217.
- ICHIHARA, T.  
1966. The pygmy blue whale, *Balaenoptera musculus brevicauda*, a new subspecies from the Antarctic. In K. S. Norris (editor), Whales, dolphins, and porpoises, p. 79-113. Univ. Calif. Press, Berkeley and Los Ang.
- INTERNATIONAL TRUST FOR ZOOLOGICAL NOMENCLATURE.  
1964. International code of zoological nomenclature adopted by the XV International Congress of Zoology. [In Engl. and French.] Int. Trust Zool. Nomen., Lond., 176 p.
- INTERNATIONAL UNION FOR CONSERVATION OF NATURE AND NATURAL RESOURCES.  
1972. Red data book. Vol. 1: *Mammalia*. Int. Union Conserv. Nat. Natur. Resour., Morges, Switz.
- INTERNATIONAL WHALING COMMISSION, SUBCOMMITTEE ON SMALL CETACEANS.  
1975. Report of the meeting on smaller cetaceans: Montreal, April 1-11, 1974. J. Fish. Res. Board Can. 32:889-983.
- JONES, R. E.  
1967. A *Hydrodamalis* skull fragment from Monterey Bay, California. J. Mammal. 48:143-144.
- KASUYA, T.  
1972. Some informations on the growth of the Ganges dolphin with a comment on the Indus dolphin. Sci. Rep. Whales Res. Inst. 24: 87-108.  
1973. Systematic consideration of Recent toothed whales based on the morphology of tympano-periotic bone. Sci. Rep. Whales Res. Inst. 25:1-103.  
1975. Past occurrence of *Globicephala melaena* in the western North Pacific. Sci. Rep. Whales Res. Inst. 27:95-110.
- KELLOGG, R.  
1928. The history of whales—their adaptation to life in the water. Q. Rev. Biol. 3:29-76, 174-208.  
1936. A review of the Archaeoceti. Carnegie Inst. Wash. Publ. 482, 366 p.
- KHUZIN, R. SH.  
1963. Materialy po morfologicheskoi kharakteristike trekh stad grenlandskogo tyulenyia *Pagophilus groenlandicus* Erxl. 1777. Polyar. Nauchn.-Issled. Proekt. Inst. Morsk. Rybn. Khoz. Okeanogr., Murmansk, Materialy Rybokhoz. Issled. Severn. Basseina, Sbornik 1:51-54.  
1967. Izmehchivost'kranologicheskikh priznakov groendlanskogo tyulenyia (Variability of the craniological features of the harp seal *Pagophilus groenlandicus* (Erxleben)). Tr. Polyar. Nauchn.-Issled. Proekt. Inst. Morsk, Rybn. Khoz. Okeanogr. 21:27-50. (In Russ. Transl. Fish. Res. Board Can., Transl. Ser. 1306, 1969, 41 p.)
- KING, J. E.  
1960. Sea-lions of the genera *Neophoca* and *Phocarcetos*. Mammalia 24:445-456.  
1964. Seals of the world. Br. Mus. (Nat. Hist.), Lond., 154 p.
- KLEINENBERG, S. E.  
1958. On the origin of Cetacea. Proc. 15th Int. Congr. Zool., p. 445-447.
- KOSYGIN, G. M., and V. A. POTELOV.  
1971. Vozrastnaya, polovaya i populyatsionnaya izmenchivost'kranologicheskikh priznakov morskogo zaitsa (Age, sex and population variability of the craniological characters of bearded seals). Tr. Vses. Nauchn.-Issled. Inst. Rybn. Khoz. Okeanogr. 82:266-288. (In Russ. Transl. Fish. Res. Board Can., Transl. Ser. 2651, 1973, 29 p.)
- LYDEKKER, R.  
1909. On the skull-characters in the southern sea-elephant. Proc. Zool. Soc. Lond. 1909:600-606.
- McLAREN, I. A.  
1966. Taxonomy of harbor seals of the western North Pacific and evolution of certain other hair seals. J. Mammal. 47:466-473.
- MANSFIELD, A. W.  
1967. Distribution of the harbor seal, *Phoca vitulina* Linnaeus, in Canadian arctic waters. J. Mammal. 48:249-257.
- MATTHES, H. W.  
1962. Verbreitung der Säugetiere in der Vorzeit. In Handbuch der Zool., Band 8, Lief. 28, 11(1):1-198. Walter de Gruyter & Co., Berl.
- MILLER, G. S., JR.  
1923. The telescoping of the cetacean skull. Smithson. Misc. Collect. 76(5):1-72.
- MITCHELL, E. D.  
1975. Parallelism and convergence in the evolution of Otariidae and Phocidae. Rapp. P.-V. Réun. Cons. Int. Explor. Mer 169:12-26.
- MITCHELL, E., and R. H. TEDFORD.  
1973. The Enaliarctinae a new group of extinct aquatic Carnivora and a consideration of the origin of the Otariidae. Bull. Am. Mus. Nat. Hist. 151:201-284.
- MOORE, J. C.  
1968. Relationships among the living genera of beaked whales with classifications, diagnoses and keys. Fieldiana:Zool. 53:209-298.
- MORRELL, B.  
1832. A narrative of four voyages to the south sea, north and south Pacific Ocean, Chinese sea, Ethiopic and southern Atlantic Ocean, Indian and Antarctic Ocean. From the year 1822 to 1831. J. & J. Harper, N.Y., 492 p.
- MULLER, J.  
1954. Observations on the orbital region of the skull of the *Mystacoceti*. Zool. Meded. 32:279-290.
- MURIE, O. J.  
1959. Fauna of the Aleutian Islands and Alaska Peninsula. U.S. Fish Wildl. Serv., North Am. Fauna 61, p. 1-364.
- NISHIWAKI, M.  
1963. Taxonomical consideration on genera of *Delphinidae*. Sci. Rep. Whales Res. Inst. 17:93-104.  
1973. Status of the Japanese sea lion. Int. Union Conserv. Nat. Natur. Resour. IUCN Publ., New Ser., Suppl. Pap. 39:80-81.
- NOBLE, B. A., and F. C. FRASER.  
1971. Description of a skeleton and supplementary notes on the skull of a rare porpoise *Phocoena sinus* Norris & McFarland 1958. J. Nat. Hist. 5:447-464.
- NORRIS, K. S., and W. N. McFARLAND.  
1958. A new harbor porpoise of the genus *Phocoena* from the Gulf of California. J. Mammal. 39:22-39.

- OMURA, H.  
1975. Osteological study of the minke whale from the Antarctic. *Sci. Rep. Whales Res. Inst.* 27:1-36.
- OMURA, H., S. OHSUMI, T. NEMOTO, K. NASU, and T. KASUYA.  
1969. Black right whales in the North Pacific. *Sci. Rep. Whales Res. Inst.* 21:1-78.
- PERRIN, W. F.  
1970. Color pattern of the eastern Pacific spotted porpoise *Stenella graffmani* Lönnberg (Cetacea, Delphinidae). *Zoologica (N.Y.)* 54:135-149.  
1972. Color patterns of spinner porpoises (*Stenella* cf. *S. longirostris*) of the eastern Pacific and Hawaii, with comments on delphinid pigmentation. *Fish. Bull., U.S.* 70:983-1003.  
1975a. Distribution and differentiation of populations of dolphins of the genus *Stenella* in the eastern tropical Pacific. *J. Fish. Res. Board Can.* 32:1059-1067.  
1975b. Variation of spotted and spinner porpoise (Genus *Stenella*) in the eastern Pacific and Hawaii. *Bull. Scripps Inst. Oceanogr., Univ. Calif.* 21:1-206.
- PILLERI, G., and M. GIHR.  
1971. Differences observed in the skulls of *Plantanista indi* and *gangetica*. In G. Pilleri (editor), *Investigations on Cetacea*, Vol. 3, Part 1, p. 13-21. *Inst. Brain Anat., Berne, Switz.*  
1975. On the taxonomy and ecology of the finless black porpoise, *Neophocaena* (Cetacea, Delphinidae). *Mammalia* 39:657-673.
- PIVETEAU, J. (editor).  
1958. *Traité de paléontologie*. Tome 6, Vol. 2, 962 p. Masson, Paris.  
1961. *Traité de paléontologie*. Tome 6, Vol. 1, 1138 p. Masson, Paris.
- REINHART, R. H.  
1959. A review of the Sirenia and Desmostylia. *Univ. Calif. Publ. Geol. Sci.* 36:1-146.
- REPENNING, C. A.  
1975. Otarioid evolution. *Rapp. P.-V. Réunion. Cons. Int. Explor. Mer* 169:27-33.
- REPENNING, C. A., R. S. PETERSON, and C. L. HUBBS.  
1971. Contributions to the systematics of the southern fur seals, with particular reference to the Juan Fernández and Guadalupe species. *Antarct. Res. Ser.* 18:1-34.
- RICE, D. W., and V. B. SCHEFFER.  
1968. A list of the marine mammals of the world. *U.S. Fish Wildl. Serv., Spec. Sci. Rep. Fish.* 579, 16 p.
- ROEST, A. I.  
1973. Subspecies of the sea otter, *Enhydra lutris*. *Los Ang. Cty. Mus., Contrib. Sci.* 252:1-17.
- ROMER, A. S.  
1966. *Vertebrate paleontology*. 3d ed. Univ. Chicago Press, Chicago, 468 p.
- SCHEFFER, V. B.  
1958. Seals, sea lions, and walruses; a review of the Pinnipedia. *Stanford Univ. Press, Stanford*, 179 p.
- SCHEFFER, V. B., and D. W. RICE.  
1963. A list of the marine mammals of the world. *U.S. Fish Wildl. Serv., Spec. Sci. Rep. Fish.* 431, 12 p.
- SERGEANT, D. E., and P. F. BRODIE.  
1969. Body size in white whales, *Delphinapterus leucas*. *J. Fish. Res. Board Can.* 26:2561-2580.
- SHAUGHNESSY, P. D.  
1974. Biochemical identification of populations of the harbor seal, *Phoca vitulina*. Ph.D. Thesis, University of Alaska, Fairbanks, 179 p.
- SIMPSON, G. G.  
1945. The principles of classification and a classification of mammals. *Bull. Am. Mus. Nat. Hist.* 85:1-350.  
1961. *Principles of animal taxonomy*. Columbia Univ. Press, N.Y., 247 p.
- SLIJPER, E. J.  
1936. Die Cetaceen. *Capita Zool.* 7:1-590.
- SMIRNOV, N.  
1927. Diagnostic remarks about seals (Phocidae) of the Northern Hemisphere. *Tromsø. Mus. Årsh.* 48(5):1-23.
- STEJNEGER, L.  
1897. The Russian fur-seal islands. *U.S. Fish Comm. Bull.* for 1896, 16:1-148.  
1898. The Asiatic fur-seal islands and fur-seal industry. In D. S. Jordan (editor), *The fur seals and fur-seal islands of the North Pacific Ocean*, part 4, 384 p. Government Printing Office, Wash., D.C.
- TRUE, F. W.  
1889. Contributions to the natural history of the cetaceans, a review of the family Delphinidae. *Bull. U.S. Natl. Mus.* 36, 192 p.
- U.S. MARINE MAMMAL COMMISSION.  
1976. Marine mammal names used by the Marine Mammal Commission. Wash., D.C., 8 p.
- van BREE, P. J. H.  
1971. On *Globicephala sieboldii* Gray, 1846, and other species of pilot whales. *Beaufortia* 19:79-87.  
1973. *Neophocaena phocaenoides asiaorientalis* (Pilleri & Gühr, 1973), a synonym of the preoccupied name *Delphinus melas* Schlegel, 1841. *Beaufortia* 21:17-24.
- van BREE, P. J. H., and P. E. PURVES.  
1972. Remarks on the validity of *Delphinus bairdii* (Cetacea, Delphinidae). *J. Mammal.* 53:372-374.
- van BREE, P. J. H., and D. ROBINEAU.  
1973. Notes sur les holotypes de *Inia geoffrensis geoffrensis* (de Blainville, 1817) et de *Inia geoffrensis boliviensis* d'Orbigny, 1834 (Cetacea, Platanistidae). *Mammalia* 37:658-668.
- VAZ FERREIRA, R.  
1976. *Arctocephalus australis* (Zimmerman), South American fur seal. *Adv. Comm. Mar. Resour. Res., Sci. Consult. Mamm.* 49:1-13.
- WINGE, H.  
1921. A review of the interrelationships of the Cetacea. *Smithson. Misc. Collect.* 72(8):1-97.
- YABLOKOV, A. V., and V. YA. ETIN.  
1965. Analysis of population differences in the body coloring of mammals (as exemplified by the harp seal, *Phoca groenlandica*). [In Russ., Engl. summ.] *Zool. Zh.* 44:1103-1106.
- YABLOKOV, A. V., and D. E. SERGEANT.  
1963. Cranial variations in the harp seal (*Pagophilus groenlandicus* Erxleben, 1777). *Zool. Zh.* 42:1857-1865. (In Russ., *Can. State Bur. Transl. No.* 3908, 1964, 16 p.)
- ZEMSKY, V. A., and V. A. BORONIN.  
1964. On the question of the pygmy blue whale taxonomic position. *Nor. Hvalfangst-Tidsskr.* 53:306-311.