

Laboratory Guide to Early Life History Stages of Northeast Pacific Fishes

Ann C. Matarese, Arthur W. Kendall, Jr.,
Deborah M. Blood, and Beverly M. Vinter



NOAA TECHNICAL REPORT NMFS

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for their optimum use. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The NOAA Technical Report NMFS series was established in 1983 to replace two subcategories of the Technical Reports series: "Special Scientific Report—Fisheries" and "Circular." The series contains the following types of reports: Scientific investigations that document long-term continuing programs of NMFS; intensive scientific reports on studies of restricted scope; papers on applied fishery problems; technical reports of general interest intended to aid conservation and management; reports that review in considerable detail and at a high technical level certain broad areas of research; and technical papers originating in economics studies and from management investigations. Since this is a formal series, all submitted papers receive peer review and those accepted receive professional editing before publication.

Copies of NOAA Technical Reports NMFS are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained from: U.S. Department of Commerce, National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161. Although the contents have not been copyrighted and may be reprinted entirely, reference to source is appreciated.

48. Widow rockfish: Proceedings of a workshop, Tiburon, California, December 11–12, 1980, by William H. Lenarz and Donald R. Gunderson (editors). January 1987, 57 p.
49. Reproduction, movements, and population dynamics of the southern kingfish, *Menticirrhus americanus*, in the northwestern Gulf of Mexico, by Stephen M. Harding and Mark E. Chittenden, Jr. March 1987, 21 p.
50. Preparation of acetate peels of valves from the ocean quahog, *Arctica islandica*, for age determinations, by John W. Ropes. March 1987, 5 p.
51. Status, biology, and ecology of fur seals: Proceedings of an international workshop, Cambridge, England, 23–27 April 1984, by John P. Croxall and Roger L. Gentry (editors). June 1987, 212 p.
52. Limited access alternatives for the Pacific groundfish fishery, by Daniel D. Huppert (editor). May 1987, 45 p.
53. Ecology of east Florida sea turtles: Proceedings of the Cape Canaveral, Florida, sea turtle workshop, Miami, Florida, February 26–27, 1985, by Wayne N. Witzell (convener and editor). May 1987, 80 p.
54. Proximate and fatty acid composition of 40 southeastern U.S. finfish species, by Janet A. Gooch, Malcolm B. Hale, Thomas Brown, Jr., James C. Bonnet, Cheryl G. Brand, and Lloyd W. Reiger. June 1987, 23 p.
55. Proximate composition, energy, fatty acid, sodium, and cholesterol content of finfish, shellfish, and their products, by Judith Krzynowek and Jenny Murphy. July 1987, 53 p.
56. Some aspects of the ecology of the leatherback turtle *Dermochelys coriacea* at Laguna Jolova, Costa Rica, by Harold F. Hirth and Larry H. Ogren. July 1987, 14 p.
57. Food habits and dietary variability of pelagic nekton off Oregon and Washington, 1979–1984, by Richard D. Brodeur, Harriet V. Lorz, and William G. Pearcy. July 1987, 32 p.
58. Stock assessment of the Gulf menhaden, *Brevoortia patronus*, fishery, by Douglas S. Vaughan. September 1987, 18 p.
59. Atlantic menhaden, *Brevoortia tyrannus*, purse seine fishery, 1972–84, with a brief discussion of age and size composition of the landings, by Joseph W. Smith, William R. Nicholson, Douglas S. Vaughan, Donnie L. Dudley, and Ethel A. Hall. September 1987, 23 p.
60. Gulf menhaden, *Brevoortia patronus*, purse seine fishery, 1974–85, with a brief discussion of age and size composition of the landings, by Joseph W. Smith, Eldon J. Levi, Douglas S. Vaughan, and Ethen A. Hall. December 1987, 8 p.
61. Manual for starch gel electrophoresis: A method for the detection of genetic variation, by Paul B. Aebersold, Gary A. Winans, David J. Teel, George B. Milner, and Fred M. Utter. December 1987, 19 p.
62. Fishery publication index, 1980–85; Technical memorandum index, 1972–85, by Cynthia S. Martin, Shelley E. Arenas, Jacki A. Guffey, and Joni M. Packard. December 1987, 149 p.
63. Stock assessment of the Atlantic menhaden, *Brevoortia tyrannus*, fishery, by Douglas S. Vaughan and Joseph W. Smith. January 1988, 18 p.
64. Illustrated key to penaeoid shrimps of commerce in the Americas, by Isabel Pérez Farfante. April 1988, 32 p.
65. History of whaling in and near North Carolina, by Randall R. Reeves and Edward Mitchell. March 1988, 28 p.
66. Atlas and zoogeography of common fishes in the Bering Sea and northeastern Pacific, by M. James Allen and Gary B. Smith. April 1988, 151 p.
67. Index numbers and productivity measurement in multispecies fisheries: An application to the Pacific coast trawl fleet, by Dale Squires. July 1988, 34 p.
68. Annotated bibliography II of the hard clam *Mercenaria mercenaria*, by J.L. McHugh and Marjorie W. Sumner. September 1988, 59 p.
69. Environmental quality and aquaculture systems: Proceedings of the thirteenth U.S.–Japan meeting on aquaculture, Mie, Japan, October 24–25, 1984, edited by Carl J. Sindermann. October 1988, 50 p.
70. New and innovative advances in biology/engineering with potential for use in aquaculture: Proceedings of the fourteenth U.S.–Japan meeting on aquaculture, Woods Hole, Massachusetts, October 16–17, 1985, edited by Albert K. Sparks. November 1988, 69 p.
71. Greenland turbot *Reinhardtius hippoglossoides* of the eastern Bering Sea and Aleutian Islands region, by Miles S. Alton, Richard G. Bakkala, Gary E. Walters, and Peter T. Munro. December 1988, 31 p.
72. Age determination methods for northwest Atlantic species, edited by Judy Penttila and Louise M. Dery. December 1988, 135 p.
73. Marine flora and fauna of the Eastern United States. Mollusca: Cephalopoda, by Michael Vecchione, Clyde F.E. Roper, and Michael J. Sweeney. February 1989, 23 p.
74. Proximate composition and fatty acid and cholesterol content of 22 species of northwest Atlantic finfish, by Judith Krzynowek, Jenny Murphy, Richard S. Maney, and Laurie J. Panunzio. May 1989, 35 p.
75. Codend selection of winter flounder *Pseudopleuronectes americanus*, by David G. Simpson. March 1989, 10 p.
76. Analysis of fish diversion efficiency and survivorship in the fish return system at San Onofre Nuclear Generating Station, by Milton S. Love, Meenu Sandhu, Jeffrey Stein, Kevin T. Herbinson, Robert H. Moore, Michael Mullin, and John S. Stephens, Jr. April 1989, 16 p.
77. Illustrated key to the genera of free-living marine nematodes of the order Enopliida, by Edwin J. Keppner and Armen C. Tarjan. July 1989, 26 p.
78. Survey of fishes and water properties of south San Francisco Bay, California, 1973–82, by Donald E. Pearson. August 1989, 21 p.
79. Species composition, distribution, and relative abundance of fishes in the coastal habitat off the southeastern United States, by Charles A. Wenner and George R. Sedberry. July 1989, 49 p.
80. Laboratory guide to early life history stages of northeast Pacific fishes, by Ann C. Matarese, Arthur W. Kendall, Jr., Deborah M. Blood, and Beverly M. Vinter. October 1989, 651 p.

NOAA Technical Report NMFS 80

Laboratory Guide to Early Life History Stages of Northeast Pacific Fishes

Ann C. Matarese
Arthur W. Kendall, Jr.
Deborah M. Blood
Beverly M. Vinter

October 1989



U.S. DEPARTMENT OF COMMERCE
Robert Mosbacher, Secretary
National Oceanic and Atmospheric Administration
John A. Knauss, Under Secretary for Oceans and Atmosphere
National Marine Fisheries Service
James Brennan, Assistant Administrator for Fisheries

The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

Acknowledgments iv

Introduction 1

- Background 1
- Geographic coverage 2
- Information sources
 - Literature 2
 - Specimens 3

Species list 6

Identifying fish eggs and larvae 13

- Eggs 13
- Larvae 13
 - Meristic characters 13
 - Pigmentation 14
 - Morphology 14

Using this laboratory guide 28

- Format 28
- Meristics 28
- General life history 28
- Early life history 29

Taxon accounts

- Elopomorpha 31
- Clupeiformes 43
- Salmoniformes 51
- Stomiiformes 83
- Scopelomorpha 125
- Gadiformes 181
- Ophidiiformes 211
- Lophiiformes 221
- Gobiesociformes 229
- Atherinomorpha 235
- Lampriformes 243
- Beryciformes 249
- Zeiformes 257
- Gasterosteiformes 261
- Scorpaeniformes 267
- Perciformes 483
- Pleuronectiformes 565
- Tetraodontiformes 627

Citations 630

Index to scientific names 642

Index to common names 650

Acknowledgments

A project of this scope and magnitude always involves many people whose talents and skills were essential throughout its development and completion. This guide began as an idea in 1979 and has gradually evolved over the last 10 years with the assistance of numerous people. During the early stages, two major projects contracted with the University of Washington (UW) served as starting points. Kevin Howe, formerly of the UW, was asked to compile a checklist of fishes from Point Conception, California, north into the Arctic Ocean, and to set up a meristic database system. His assembled data form the basis of meristic sections on our taxonomic text pages and meristic tables. Kathryn Garrison, formerly of the UW, and Bruce Miller (UW) synthesized the available literature on the reproduction and early life history of Puget Sound fishes, and this enabled us to write many of our sections on life history.

During the several years we spent collecting data and examining material, a number of scientists donated specimens and/or unpublished data. We would like to thank the following people: Jeffrey Marliave (Vancouver Public Aquarium), David Misitano (NWAFC Mukilteo Laboratory), Albert Giorgi (NWAFC Coastal Zone and Estuarine Studies), Kevin Bailey (NWAFC Resource Assessment and Conservation Engineering), William Watson (Marine Ecological Consultants), Bruce Mundy (formerly Oregon State University), A. J. Paul (University of Alaska), Conrad Mahnken (NWAFC Manchester Field Station), Sally L. Richardson (deceased), and H. Geoffrey Moser (NMFS Southwest Fisheries Center). The complete files of the late E. H. Ahlstrom were kindly made available by H. Geoffrey Moser.

Throughout the planning, organizing, writing, and layout, many people contributed their special talents and we thank the following for their time: Theodore Pietsch, Kevin Howe, and Steven Leipertz, UW (meristic database); Kevin Howe and Steven Leipertz, UW, and Richard Bates, NWAFC, (literature database); Ralph Mintel, Richard Bates, and Michael McPhail, NWAFC (programming);

William Rugen, NWAFC (proofreading and verification); James Peacock, NWAFC (layout, typesetting); Jack McCormick and Nancy Peacock, NMFS Scientific Publications (format and editorial assistance); William Richards, formerly NMFS Scientific Editor (editorial assistance); and Michael Fahay, NMFS Northeast Fisheries Center, who has encouraged all of us along the way and who spent several days reviewing our very preliminary first draft.

We are grateful to the following, who in reviewing sections from an earlier draft made many suggestions and corrections and generously provided us with unpublished data from their personal data files: James Allen (ecology and life history), G. David Johnson (Perciformes), Wayne Laroche (Scorpaenidae, Agonidae, Carangidae), Douglas Markle (Gadiformes, Ophidiiformes), H. Geoffrey Moser (Myctophidae, Stomiiformes, Pleuronectiformes, Scorpaenidae), Bruce Mundy (meristic data), Barbara Sumida MacCall (Carangidae, Pleuronectidae, Stromateoidei), Betsy Washington (Scorpaeniformes). James Allen, Robert Lea, and Alex Peden reviewed the adult distributions and nomenclature.

The following shared unpublished figures with us: Kathryn Garrison and Jeffrey Marliave (stichaeids); Betsy Washington (cottids); William Watson (*Atherinops* and *Atherinopsis*); and Lucy Wold and Guillermo Moreno (*Sebastes mystinus*).

The following reviewed the entire draft and made many valuable suggestions, and we thank them sincerely for their time and efforts: Michael Fahay, Jeffrey Leis, Joanne Lyczkowski-Shultz, Douglas Markle, Jeffrey Marliave, Gerald McGowen, H. Geoffrey Moser, Bruce Mundy, Muneo Okiyama, William Richards, Betsy Washington, and William Watson.

Finally, we all express a special thanks to our friend and colleague Jean Dunn for his many contributions throughout the various stages of this guide. His years of experience with the early-life-history stages of North Pacific fishes and extensive knowledge of the literature were a valuable addition to this work.

Laboratory Guide to Early Life History Stages of Northeast Pacific Fishes

ANN C. MATARESE
ARTHUR W. KENDALL, JR.
DEBORAH M. BLOOD
BEVERLY M. VINTER

*Northwest and Alaska Fisheries Center
National Marine Fisheries Service, NOAA
7600 Sand Point Way N.E.
Seattle, Washington 98115*

ABSTRACT

This laboratory guide presents taxonomic information on eggs and larvae of fishes of the Northeast Pacific Ocean (north of California) and the eastern Bering Sea. Included are early-life-history series, illustrations, and comparative descriptions of 232 species expected to spawn here, out of a total 627 species known to occur in marine waters of this area. Meristic and general life-history data are included, as well as diagnostic characters to help identify eggs and larvae. Most of this information has been gleaned from literature, with the addition of 200 previously unpublished illustrations.

Introduction

Background

The importance of early-life-history studies to fisheries investigations and phylogenetic research has increased dramatically during the last decade. Early-life-history stages are now routinely used in fisheries studies to investigate the interannual variation in recruitment (e.g., Wooster 1983), and in studies of the phylogeny of fishes (e.g., Moser et al. 1984b). The fact that early-life-history stages of many species remain unknown in the northeastern Pacific Ocean limits their use in these disciplines, as well as in research on ecology, behavior, and the biological effects of pollution on fishes.

At the Northwest and Alaska Fisheries Center (NWAFC), progress in the identification and understanding of early-life-history stages of marine fishes has steadily increased. Our investigations beginning in 1965 were primarily aimed at determining spawning grounds and distribution of eggs and larvae for only one target species: the Pacific hake, *Merluccius productus*. Studies in the late sixties and early seventies concentrated on ascertaining abundance and horizontal distribution of the eggs and larvae of major taxa occurring off the northeast Pacific coast and in the Gulf of Alaska and Bering Sea. In addition to baseline studies on distribution and abundance, work has increasingly emphasized the use of egg and/or larval surveys (targeted on species such as walleye pollock, *Theragra chalcogramma*) to generate estimates of spawning stock biomass and to test hypotheses concerning the multitude of possible factors involved in the stock/recruitment relationship (e.g., feeding, predation, growth, and transport). These studies require the accurate identification of all early-life-history stages, from the egg stage to newly settled juveniles and adults. The ability to make such identifications is a direct result of taxonomic studies in our laboratory, especially for the family Gadidae. These studies have enabled identification of all gadid species occurring in the Northeast Pacific Ocean and have allowed additional research on development, osteology, and systematics (e.g., Matarese et al. 1981, Dunn and Vinter 1984, and Dunn and Matarese 1984).

The primary purpose of this laboratory guide is to allow expansion of the use of early-life-history studies in fisheries research by providing descriptive information necessary to insure accurate identification of eggs, larvae, and early juveniles of marine fishes in the northeastern Pacific Ocean and Bering Sea. The guide is designed to aid in species identification; thus, it includes little of the developmental information usually found in ontogenetic descriptions. We have placed emphasis on the accuracy and quality of illustrations, whether from the literature or originals drawn for this guide. In some cases, however, when specimens were not available, we reproduced substandard figures from the literature. Diagnostic characters are included, which enable identification of early-life-history stages of closely related and/or similar-looking taxa, as are comparative tables and figures which outline similarities and differences among taxa. Detailed information is included for species where considerable early-life-history data are available. For poorly known taxa, available information is given, sometimes limited to characters of adults of the species in the study area and a description of early-life-history stages of related species from other areas.




Figure 1
Study area showing geographic ranges used in species accounts.

Geographic coverage

The geographic area covered is the Pacific Ocean between approximately 38°N and 66°N and west to 180°; however, we have included only those taxa with northern limits of distribution between Oregon and the northern Bering Sea (Fig. 1). Taxa restricted to California waters or the Arctic are excluded. Because most ichthyoplankton surveys concentrate on coastal areas, usually within 200 miles (370 km) of shore, this guide emphasizes coastal, nearshore species rather than strictly oceanic species. Taxa that usually do not produce planktonic early-life-history stages are generally excluded (e.g., Embiotocidae) as are strictly freshwater and estuarine species or spawners. Sources useful in identifying early-life-history stages of freshwater and estuarine species found adjacent to our study area include Wang (1981, 1986) and Auer (1982). In general, we have based coverage on species' likelihood of spawning in the Northeast Pacific Ocean rather than on the occurrence of adults. The geographic distribution of spawning is generally more restricted than the overall range of a species. Some low-latitude oceanic species (e.g., most members of the family Scombridae) which occur off Oregon and Washington are not included because they usually spawn much further south and their eggs and larvae do not occur in our area.

Information sources

Literature In producing this guide, we first compiled meristic and life history information from the literature. This was accomplished largely through a computer-based meristic information file generated by Kevin Howe at the University of Washington, and a compendium of life history information produced under the direction of Bruce Miller, University of Washington (Garrison and Miller 1982). Information from these sources was augmented and updated as it was incorporated in this guide. The meristic database was compiled by examining specimens and original radiographs, and from the literature (original descriptions, revisions, or general sources). When possible, counts are from the left side of the body. Since Garrison and Miller (1982) is not generally available, original sources cited by them are referred to in this guide. Published early-life-history information and illustrations were used when available, supplemented when possible by data from our field collections. Among the most useful published sources were Fitch and Lavenberg (1968, 1971, 1975), Miller and Lea (1972), Hart (1973), Eschmeyer et al. (1983), Moser et al. (1984b), Ozawa (1986a), and the many papers authored or co-authored by Ahlstrom, Moser, and Richardson (see Citations). Okiyama (1988) was received too late to be fully included in this guide.

Specimens In addition to data gleaned from the literature, we have included some unpublished data and original illustrations obtained from specimens in our collections (Table 1). Specimens were from two major sources: (1) Investigations in the Kodiak Island region in the Gulf of Alaska conducted 1977-79, and (2) studies undertaken in cooperation with the U.S.S.R. beginning in 1980 to investigate the distribution and abundance of ichthyoplankton off northern California, Oregon, and Washington and in the Gulf of Alaska.

Some specimens were also derived from work in the Bering Sea during 1971-72, 1976-78, and 1979. Dunn (1986) provides a list of ichthyoplankton surveys conducted by the NWAFC from which were obtained many of the specimens examined or illustrated for this guide. Occasionally specimens were obtained from other research collections or from rearing experiments conducted primarily by Jeffrey Marliave (Vancouver Public Aquarium) and David Misitano (NWAFC).

Table 1
Collection data for original illustrations by Northwest and Alaska Fisheries Center (NWAFC).^a

Taxon	SL (mm)	Cruise	Station number	Gear ^b	Date	Location or N° W°
<i>Thalassenchelys coheni</i>	190.0				15 02 84	Puget Sound, WA ^c
<i>Clupea pallasii</i>	10.4	4MF81	G005A	6B5	20 05 81	58°19.6', 153°54.5'
	15.0	SF7703	P-4	N	05 04 77	Str. Juan de Fuca, WA
	19.0	SF7603	13	B	18 05 76	Str. Juan de Fuca, WA
	23.8			seine	14 06 72	
<i>Nansenia candida</i>	5.1	1PO84	G053A	6B5	22 03 84	44°40.1', 127°40.0'
<i>Bathylagus milleri</i>	15.0	2MF78	G074A	TT	02 07 78	57°07.0', 151°01.0'
<i>Bathylagus ochotensis</i>	7.9	1EQ83	G056A	B	03 05 83	44°00.0', 127°39.0'
<i>Bathylagus pacificus</i>	11.6	3MF79	S25A	6B5	17 06 79	54°05.2', 170°59.0'
	17.6			6B5		Gulf of Alaska
<i>Macropinna microstoma</i>	11.1	1PO84	G066A	6B5	24 03 84	43°40.0', 124°37.0'
<i>Danaphos oculatus</i>	22.4		M161		23 07 84	6 km W of Newport, OR
<i>Tactostoma macropus</i>	14.3	1PO80	G011A	6B5	03 08 80	47°18.7', 125°13.3'
<i>Lampanyctus regalis</i>	5.2	1PO80	G061A	6B5	12 08 80	38°00.0', 125°42.5'
<i>Diaphus theta</i>	4.6	K6703	50	1MN	08 05 67	42°00.0', 127°35.0'
	6.3	1PO80	G085A	6B5	20 08 80	40°20.0', 125°20.0'
	16.0	1PO80	G064A	6B5	13 08 80	42°22.5', 126°17.5'
<i>Stenobranchius leucopsarus</i>	4.9	2KE72	G50A	6B5	08 05 72	57°46.0', 149°21.0'
	6.3	1PO84	G112A	6B5	02 04 84	40°40.1', 126°47.0'
	18.0	4DI78	G051A	TT1	13 04 78	55°26.8', 153°53.7'
<i>Melanonus zugmayeri</i>	17.0	Ocean ACRE-9	9-11 N		19 03 70	32°03.0', 64°05.0'
<i>Boreogadus saida</i>	16.3	Glacier (OCSEAP)	11	6B5	11 08 76	70°47.0', 162°14.0'
<i>Gadus macrocephalus</i>	4.1	FOX86III (sample 111)	166	MOC	18 05 86	57°40.6', 155°09.8'
<i>Theragra chalcogramma</i>	egg	composite				Gulf of Alaska
	3.5				— 04 82	Gulf of Alaska ^d
Ophidiidae	15.6	1EQ83	G044A	6B5	30 04 83	45°20.0', 124°48.0'
	29.8	1EQ83	G073A	6B5	06 05 83	43°20.2', 127°57.0'
<i>Brosomphycis marginata</i>	10.6					Puget Sound ^e
<i>Gobiesox maeandricus</i>	7.0				12 02 76	Brit. Columbia ^f
<i>Cololabis saira</i>	6.7	1TK80	G091	N	08 05 80	42°00.0', 125°55.0'
	7.4	IDA81	G039A	N	02 11 81	45°37.0', 124°52.0'
<i>Trachipterus altivelis</i>	9.4	K6805	30	1MN	23 10 68	45°44.0', 124°38.0'
	24.0	1PO82	G46A	6B5	29 05 82	40°40.5', 126°48.4'
<i>Melamphaes</i> sp.	3.7	1EQ83	G034A	6B5	28 04 83	46°00.0', 128°31.0'
<i>Sebastes brevispinis</i>	4.6	SEI77-9	21	trawl	12 07 77	58°28.3', 139°30.0'
<i>Sebastes caurinus</i>	5.1				05 07 77	Reared by C. Mahnken ^g
<i>Sebastes jordani</i>	4.6	Marathon 85-1	229	trawl	25 05 85	48°09.6', 125°05.3'
<i>Sebastes melanops</i>	4.0				07 02 84	Newport, OR ^h
<i>Sebastes polyspinis</i>	6.1	Poseydon 85-1		trawl	— 07 85	Gulf of Alaska
<i>Sebastes rufus</i>	36.0	(Groundfish Comm. Invest.)	8	MWT	04 06 85	Farrallon Is., CA ⁱ
<i>Sebastes variegatus</i>	4.6	SEI77-9	21	trawl	12 07 77	58°28.3', 139°30.0'
<i>Sebastes zacentrus</i>	4.7	SEI77-9	33	trawl	16 07 77	59°40.1', 143°03.5'
<i>Sebastolobus</i> sp.	egg	1PO81	G065A	N	22 05 81	43°40.0', 125°01.4'

Table 1 (continued)

Taxon	SL (mm)	Cruise	Station number	Gear ^b	Date	Location or	
						N°	W°
<i>Anoplopoma fimbria</i>	5.6	1PO84	G019B	6B5	15 03 84	46°40.5'	124°59.0'
	8.8	MF77B-6	83(1)	N		55°40.7'	155°23.0'
	12.0	MF77B-6	2(4)	N	15 05 77	54°22.6'	166°42.5'
<i>Blepsias bilobus</i>	12.4	3MF79	V02A	N	02 06 79	56°03.5'	166°33.9'
	16.7	3MF79	S12A	N	06 06 79	56°35.7'	165°54.3'
	24.8	3MF79	S40A	N	21 06 79	56°31.6'	166°42.0'
<i>Blepsias cirrhosus</i>	11.5				— 04 80	Friday Harbor, WA	
	16.8				— 04 80	Friday Harbor, WA	
<i>Chitonotus pugetensis</i>	3.0					Puget Sound, WA ^j	
	4.9					Puget Sound, WA ^j	
<i>Gymnocanthus</i> A	9.6	MF76A	B07	6B5	04 05 76	56°49.7'	169°39.0'
	11.9	MF76A	B17	6B5	20 05 76	54°42.3'	165°25.9'
<i>Hemilepidotus jordani</i>	10.6	1SH81	066	N	19 03 81	57°03.0'	155°53.0'
	18.4	3MF79	V06A	N	03 06 79	56°02.8'	166°33.6'
<i>Myoxocephalus</i> B	9.1	1MD82	G135A	6B5	29 05 82	54°54.1'	158°39.0'
	12.2	1MD82	G135A	6B5	29 05 82	54°54.1'	158°39.0'
<i>Myoxocephalus</i> G	8.7	2KE72	G39A	6B5	06 05 72	57°33.0'	152°06.0'
<i>Nautichthys oculofasciatus</i>	8.3				— 04 80	Friday Harbor, WA	
<i>Psychrolutes paradoxus</i>	18.0					Mukilteo, WA ^j	
<i>Synchirus gilli</i>	10.5				15 05 79	Neah Bay, WA ^k	
	16.8			N	16 05 79	Neah Bay, WA ^k	
<i>Agonidae</i> A	4.7	1PO82	G026A	6B5	14 05 82	44°01.5'	124°33.8'
	10.0					Gulf of Alaska	
<i>Agonomalus mozinoi</i>	8.2	MF77B-5	3(2)	6B5	26 04 77	54°38.7'	167°14.0'
Cyclopteridae	4.0	3MF81	G058A	5B5	01 05 81	56°55.9'	154°55.8'
<i>Nectoliparis pelagicus</i>	7.9	MF77B-6	55(1)	TT	13 05 77	55°46.7'	169°25.1'
	20.5	MF77B-6	46(1)	6B5	13 05 77	55°44.4'	171°31.3'
<i>Paraliparis</i> sp.	28.5	4DI78	25	6B5	01 04 78	57°58.3'	150°02.2'
<i>Bathymaster</i> A	9.0	2MF78	G35A	6B5	25 06 78	56°02.0'	154°08.4'
	29.6	MF77B-5	2(2)	N	26 04 77	54°23.3'	166°44.0'
<i>Ronquilus jordani</i>	7.7		803E		— 05 72	Newport, OR	
	10.4		912E		— 05 72	Newport, OR	
<i>Anoplarchus purpureus</i>	6.1					Puget Sound, WA ^j	
	9.0					Puget Sound, WA ^j	
	12.0					Puget Sound, WA ^j	
	12.0					Puget Sound, WA ^j	
	12.0					Puget Sound, WA ^j	
<i>Bryozoichthys-Chirolophis</i>	16.5	MF76A	10	6B5		56°29.8'	171°34.1'
	29.0	MF77B-6	42	N	12 05 77	56°45.6'	171°30.8'
<i>Lumpenus sagitta</i>	17.3	1MD82	G028A	6B5	08 04 82	56°40.0'	155°27.0'
	35.1	4MF81	G028A	6B5	21 05 81	57°29.5'	155°43.0'
<i>Plectobranchnus evides</i>	9.2		1143E		24 04 73	Newport, OR	
	16.9		1107E		20 04 73	Newport, OR	
	31.3		933E		— 06 72	Newport, OR	
<i>Xiphister atropurpureus</i>	8.0				20 03 78	Brit. Columbia ^f	
<i>Delolepis gigantea</i>	17.5	1MF80	34A ?	N	27 03 80	57°52.4'	154°38.9'
<i>Lyconectes aleutensis</i>	16.0	MF77B-5	8(2)	N	25 04 77	54°37.0'	166°13.9'
<i>Apodichthys flavidus</i>	~15					Brit. Columbia ^f	
<i>Pholis</i> sp.	9.2					Brit. Columbia ^f	
	23.0	2MF78	G034A	6B5	25 06 78	56°33.3'	154°51.2'
<i>Anarhichas orientalis</i>	21.0	MF77B-5	8(2)	N	25 04 77	54°37.0'	166°13.9'
<i>Ammodytes hexapterus</i>	9.8	1SH81	177	6B5	22 04 81	55°54.0'	156°36.0'
	32.3	1CH83	S008A	N	18 05 83	59°03.2'	147°31.7'
<i>Clevelandia ios</i>	3.4			TT	04 05 82	Bohom Bay, OR	
	15.0				20 01 66	Yaquina Bay, OR	
<i>Coryphopterus nicholsi</i>	4.4				21 05 85	Dabob Bay, WA ^l	
<i>Lepidogobius lepidus</i>	3.5			TT (midwater)		Yaquina Bay, OR	
	20.8			6 ft. otter trawl	19 05 84	1 mi off Newport, OR	

Table 1 (continued)

Taxon	SL (mm)	Cruise	Station number	Gear ^b	Date	Location or N° W°
<i>Citharichthys sordidus</i>	4.5	1EQ83	G042A		30 04 83	45°20.0', 124°06.0'
	7.0	1EQ83	G042A		30 04 83	45°20.0', 124°06.0'
<i>Atheresthes stomias</i>	10.0	MF76A-III	36	6B5	25 05 76	55°29.0', 165°50.8'
	13.4	DE-4	14K#2	6B5	25 07 71	56°45.0', 168°05.0'
	25.6	DE-4	16J	6B5	02 08 71	56°15.0', 171°21.0'
<i>Glyptocephalus zachirus</i>	11.5	2MF78	G68A	TT	01 07 78	56°17.3', 152°55.8'
<i>Hippoglossoides elassodon</i>	5.0	2MF78	G66A	TT	30 06 78	55°59.7', 153°33.2'
	7.9	2MF78	G66A	TT	30 06 78	55°59.7', 153°33.2'
	15.0	3MF79	S33A	6B5	19 06 79	56°16.2', 166°29.6'
	18.0	SEI77-9	43	6B5	22 07 77	58°19.5', 150°53.0'
<i>Hippoglossus stenolepis</i>	14.4	4DI78	D45C	TT	14 04 78	56°14.5', 153°22.2'
	18.0	3MF81	G012A	6B5	27 04 81	57°57.0', 154°13.3'
<i>Lepidopsetta bilineata</i>	4.3	4DI78	G009A	6B5	30 03 78	58°22.0', 150°12.8'
	7.4	5TI79	G024A	6B5	19 05 79	56°23.7', 155°45.0'
	10.8	DE-4	DE4#2	6B5	26 07 71	57°30.0', 169°30.0'
	16.3	2MF78	G023A	6B5	24 06 78	57°19.5', 152°23.9'
<i>Lepidopsetta 2</i>	6.3	2MF78	G044A	TT	26 06 78	57°61.1', 151°17.4'
	9.7	2MF78	D01A	TT	28 06 78	56°42.3', 153°33.7'
	16.4	SEI77-9	43	6B5	22 07 77	58°19.5', 150°53.0'
<i>Microstomus pacificus</i>	15.0	MF84-6	B1	6B3	25 08 84	Bering Sea ^m
	26.0	1PO80	G095A	6B5	23 08 80	39°20.0', 124°22.0'
<i>Parophrys vetulus</i>	4.5	K6805	32	1MN	23 10 68	45°44.0', 124°07.0'
	10.0	6502	14F		14 05 65	46°10.0', 124°42.0'
	17.5	SF7702	2	6B3	23 02 77	Str. Juan de Fuca, WA
<i>Platichthys stellatus</i>	4.8	4MF81	G029A	6B5	21 05 81	57°24.8', 155°37.0'
	6.6				29 04 78	Str. Juan de Fuca, WA
	8.2				29 04 78	Str. Juan de Fuca, WA
	9.0				05 06 70	Puget Sound, WA
<i>Pleuronectes quadrituberculatus</i>	6.3	3MF79	V014A	6B5	05 06 79	57°02.9', 165°02.8'
	7.8	3MF79	V015A	6B5	05 06 79	57°03.8', 165°03.2'
<i>Psettichthys melanostictus</i>	2.5				15 02 84	Puget Sound, WA ^j
	6.9				07 03 84	Puget Sound, WA ^j
	8.1				27 03 84	Puget Sound, WA ^j
	13.9		591E		— 06 —	Newport, OR

^aSeveral illustrations originally drawn for this publication and included in this table have already been published (e.g., Moser et al. 1984b, Kendall and Matarese 1987).

^bGear

6B5 = 60-cm bongo net, 0.505-mm mesh 1MN = 1-m net
 6B3 = 60-cm bongo net, 0.333-mm mesh MWT = midwater trawl
 B = 60-cm bongo net N = neuston net
 MOC = MOCNESS net TT = 1-m Tucker trawl

^cCollected by Univ. Wash., Seattle, WA 98195.

^dReared by A.J. Paul, Univ. Alaska, Inst. Mar. Sci., Seward Mar. Cent., Seward, AK 99664.

^eReared by Steven Borton, formerly of Seattle Public Aquarium, Pier 59, Seattle, WA 98104.

^fReared by Jeffrey Marliave, Vancouver Public Aquarium, P.O. Box 3232, Vancouver, B.C., Canada V6B 3X8.

^gReared by Conrad Mahnken, NWAFC, Coastal Zone Estuarine Stud., Manchester Field Stn., P.O. Box 38, Manchester, WA 98353.

^hReared by George Boehlert, Southwest Fish. Cent., Honolulu Lab., Honolulu, HI 96822.

ⁱCollected by Wayne Samiere, Southwest Fish. Cent., Tiburon Lab., Tiburon, CA 94920.

^jReared by David Misitano, NWAFC, Environ. Conserv. Div., Mukilteo Field Stn., P.O. Box 21, Mukilteo, WA 98272.

^kCollected by Albert E. Giorgi, NWAFC, Coastal Zone Estuarine Stud., 2725 Montlake Blvd. E., Seattle, WA 98112.

^lCollected by Bruce Frost, Univ. Wash., School Oceanogr., Seattle, WA 98195.

^mCollected by Kevin Bailey, NWAFC, Resource Assess. Conserv. Eng., 7600 Sand Point Way N.E., Seattle, WA 98115-0070.

Species list

The following list of species found in the study area was compiled from the literature. The order of higher taxa generally follows J. Nelson (1984); genera and species are listed alphabetically within families. The list is annotated with page numbers indicating where taxa are described in the guide. Page numbers are out of sequence in instances (e.g., myctophids, cottids) where taxa are grouped according to larval similarities, rather than alphabetically. Letters in place of page numbers indicate that a species is not given individual treatment for the following reasons: A = anadromous, D = direct development, F = freshwater spawner, S = spawns south of study area, U = inadequate early-life-history information available, V = viviparous. In these cases, available meristic and ecological information is given in appropriate higher-category (e.g., order, family) accounts.

Elopomorpha	31	<i>Therobrommus callorhinus</i>	U
Notacanthiformes		Opisthoproctidae	
Notacanthidae	34	<i>Bathylychnops exilis</i>	70
<i>Notacanthus chemnitzii</i>	U	<i>Dolichopteryx</i> sp.	72
<i>Polyacanthonotus challengeri</i>	U	<i>Dolichopteryx longipes</i>	72
Anguilliformes	33	<i>Macropinna microstoma</i>	74
Xenocoelidae	34	Alepocephalidae	76
<i>Thalassenchelys coheni</i>	U	<i>Alepocephalus tenebrosus</i>	D
Nemichthyidae		<i>Bathylaco nigricans</i>	D
<i>Avocettina infans</i>	38	<i>Ericara salmoneum</i>	D
<i>Nemichthys larseni</i>	U	<i>Leptoichthys agassizi</i>	D
<i>Nemichthys scolopaceus</i>	40	<i>Narcetes stomias</i>	D
Cyematidae	34	<i>Talismania bifurcata</i>	D
<i>Cyema atrum</i>	U	Platyroctidae	77
Synphobranchidae	34	<i>Holtbyrnia innesi</i>	U
<i>Histiobranchus bathybius</i>	U	<i>Holtbyrnia latifrons</i>	U
Nettastomatidae	36	<i>Maulisia argipalla</i>	U
<i>Venefica</i> sp.	U	<i>Pellisulus eubranchnus</i>	U
Congridae (= or includes Muraenesocidae)	36	<i>Sagamichthys abei</i>	U
<i>Xenomystax atrarius</i>	U	Salmonoidei	
Serrivomeridae	36	Osmeridae	79
<i>Serrivomer jespersenii</i>	U	<i>Allosmerus elongatus</i>	U
Eurypharyngidae	36	<i>Hypomesus olidus</i>	U
<i>Eurypharynx pelecanooides</i>	U	<i>Hypomesus pretiosus</i>	U
Clupeiformes	43	<i>Mallotus villosus</i>	80
Clupeidae		<i>Osmerus mordax</i>	A/F
<i>Alosa sapidissima</i>	A	<i>Spirinchus starksi</i>	U
<i>Clupea pallasii</i>	44	<i>Spirinchus thaleichthys</i>	A
<i>Dorosoma petenense</i>	A	<i>Thaleichthys pacificus</i>	A
<i>Sardinops sagax</i>	46	Salmonidae	
Engraulidae		Coregoninae	
<i>Engraulis mordax</i>	48	<i>Coregonus laurettae</i>	F
Salmoniformes	51	<i>Coregonus nasus</i>	F
Argentinoidei		<i>Coregonus pidschian</i>	F
Argentinidae		<i>Coregonus sardinella</i>	F
<i>Argentina sialis</i>	52	<i>Stenodus leucichthys</i>	F
<i>Nansenia candida</i>	54	Salmoninae	
Bathylagidae		<i>Oncorhynchus gorbuscha</i>	A
<i>Bathylagus bericoides</i>	56	<i>Oncorhynchus keta</i>	A
<i>Bathylagus milleri</i>	58	<i>Oncorhynchus kisutch</i>	A
<i>Bathylagus ochotensis</i>	60	<i>Oncorhynchus nerka</i>	A
<i>Bathylagus pacificus</i>	62	<i>Oncorhynchus tshawytscha</i>	A
<i>Bathylagus wesethi</i>	64	<i>Parasalmo clarki</i>	A
<i>Leuroglossus schmidti</i>	66	<i>Parasalmo gairdneri</i>	A
<i>Leuroglossus stilbius</i>	68	<i>Parasalmo mykiss</i>	A
		<i>Parasalmo penshinensis</i>	A
		<i>Salvelinus alpinus</i>	A
		<i>Salvelinus leucomaenis</i>	A
		<i>Salvelinus malma</i>	A
		Stomiiformes	83
		Gonostomatoidei	86
		Gonostomatidae	87
		<i>Cyclothone acclinidens</i>	88
		<i>Cyclothone atraria</i>	88
		<i>Cyclothone pallida</i>	88
		<i>Cyclothone pseudopallida</i>	88
		<i>Cyclothone signata</i>	88
		<i>Cyclothone</i> sp.	U
		<i>Gonostoma atlanticum</i>	90
		<i>Gonostoma gracile</i>	90

Sternoptychidae	92	<i>Stenobranchius nannochir</i>	U
<i>Argyropelecus affinis</i>	94	<i>Symbolophorus californiensis</i>	154
<i>Argyropelecus hemigymnus</i>	96	<i>Taaningichthys bathyphilus</i>	174
<i>Argyropelecus lychnus</i>	98	<i>Tarletonbeania crenularis</i>	158
<i>Argyropelecus sladeni</i>	100	<i>Tarletonbeania taylori</i>	U
<i>Sternoptyx diaphana</i>	102	Gadiformes	181
<i>Sternoptyx pseudobscura</i>	102	Moridae	182
<i>Danaphos oculus</i>	104	<i>Antimora microlepis</i>	U
Stomioidei	107	<i>Laemonema longipes</i>	U
Chauliodontidae		Melanonidae	
<i>Chauliodus macouni</i>	108	<i>Melanonus zugmayeri</i>	184
Melanostomiidae		Merlucciidae	
<i>Bathophilus flemingi</i>	110	<i>Merluccius productus</i>	186
<i>Eustomias</i> sp.	112	Gadidae	189
<i>Opostomias mitsuui</i>	114	<i>Boreogadus saida</i>	190
<i>Tactostoma macropus</i>	116	<i>Eleginus gracilis</i>	192
Malacosteidae		<i>Gadus macrocephalus</i>	194
<i>Aristostomias scintillans</i>	118	<i>Microgadus proximus</i>	196
Idiacanthidae		<i>Theragra chalcogramma</i>	198
<i>Idiacanthus antrostomus</i>	120	Macrouridae	200
<i>Idiacanthus fasciola</i>	122	<i>Albatrossia pectoralis</i>	U
Scopelomorpha	125	<i>Coryphaenoides acrolepis</i>	202
Aulopiformes		<i>Coryphaenoides armatus</i>	U
Scopelarchidae		<i>Coryphaenoides cinereus</i>	U
<i>Benthalbella dentata</i>	126	<i>Coryphaenoides filifer</i>	204
<i>Benthalbella linguidens</i>	126	<i>Coryphaenoides leptolepis</i>	206
Notosudidae		<i>Coryphaenoides liocephalus</i>	U
<i>Scopelosaurus harryi</i>	128	<i>Coryphaenoides longifilis</i>	U
Bathysauridae		<i>Coryphaenoides yaquinae</i>	U
<i>Bathysaurus mollis</i>	130	<i>Nezumia stelgidolepis</i>	208
Synodontidae		Ophidiiformes	211
<i>Synodus lucioceps</i>	132	Ophidioidei	
Paralepididae		Ophidiidae	212
<i>Lestidiops ringens</i>	134	<i>Bassozetus</i> sp.	U
<i>Notolepis rissoi</i>	136	<i>Chilara taylori</i>	214
<i>Paralepis atlantica</i>	138	<i>Dicrolene filamentosa</i>	U
Anopteridae		<i>Holcomycteronus profundissimus</i>	U
<i>Anopterus pharao</i>	140	<i>Spectrunculus grandis</i>	U
Alepisauridae		<i>Spectrunculus radcliffei</i>	U
<i>Alepisaurus ferox</i>	142	Bythioidei	217
Myctophiformes		Aphyonidae	
Neoscopelidae		<i>Barathronus pacifica</i>	U
<i>Neoscopelus macrolepidotus</i>	144	<i>Sciadonus pedicellaris</i>	U
Myctophidae	146	Bythitidae	
<i>Ceratoscopelus townsendi</i>	160	<i>Brosomphycis marginata</i>	218
<i>Diaphus theta</i>	176	<i>Cataetyx rubrirostris</i>	U
<i>Dorsadena yaquinae</i>	U	Batrachoidiformes	
<i>Electrona rissoi</i>	148	Batrachoididae	
<i>Lampadena urophaos</i>	162	<i>Porichthys notatus</i>	D
<i>Lampanyctus fernae</i>	U	Lophiiformes	221
<i>Lampanyctus jordani</i>	U	Ceratioidei	222
<i>Lampanyctus regalis</i>	166	Ceratiidae	222
<i>Lampanyctus ritteri</i>	168	<i>Ceratias holboelli</i>	U
<i>Loweina rara</i>	156	<i>Ceratias</i> sp.	U
<i>Notoscopelus resplendens</i>	178	Oneirodidae	222
<i>Notoscopelus japonicus</i>	178	<i>Bertella idiomorpha</i>	U
<i>Parvilux ingens</i>	170	<i>Chaenophryne longiceps</i>	224
<i>Protomyctophum crockeri</i>	150	<i>Chaenophryne melanorhabdus</i>	U
<i>Protomyctophum thompsoni</i>	152	<i>Oneirodes bulbosus</i>	226
<i>Stenobranchius leucopsarus</i>	172	<i>Oneirodes thompsoni</i>	U

Gobiesociformes	229	<i>Sebastes elongatus</i>	302
Gobiesocidae		<i>Sebastes emphaeus</i>	304
<i>Gobiesox maeandricus</i>	230	<i>Sebastes entomelas</i>	306
<i>Rimicola muscarum</i>	232	<i>Sebastes flavidus</i>	308
Atherinomorpha	235	<i>Sebastes glaucus</i>	U
Beloniformes		<i>Sebastes goodei</i>	U
Scomberesocidae		<i>Sebastes helvomaculatus</i>	310
<i>Cololabis saira</i>	236	<i>Sebastes jordani</i>	312
Atheriniformes		<i>Sebastes maliger</i>	U
Atherinidae		<i>Sebastes melanops</i>	314
<i>Atherinops affinis</i>	238	<i>Sebastes melanostictus</i>	U
<i>Atherinopsis californiensis</i>	240	<i>Sebastes melanostomus</i>	316
Lampriformes	243	<i>Sebastes miniatus</i>	318
Lampridae		<i>Sebastes mystinus</i>	320
<i>Lampris guttatus</i>	244	<i>Sebastes nebulosus</i>	U
Trachipteridae		<i>Sebastes nigrocinctus</i>	U
<i>Trachipterus altivelis</i>	246	<i>Sebastes paucispinis</i>	322
Beryciformes	249	<i>Sebastes pinniger</i>	324
Anoplogastridae		<i>Sebastes polyspinis</i>	U
<i>Anoplogaster cornuta</i>	250	<i>Sebastes proriger</i>	326
Melamphaidae	252	<i>Sebastes rastrelliger</i>	U
<i>Melamphaes lugubris</i>	254	<i>Sebastes reedi</i>	328
<i>Melamphaes parvus</i>	U	<i>Sebastes ruberrimus</i>	U
<i>Poromitra crassiceps</i>	U	<i>Sebastes rufus</i>	330
<i>Scopeloberyx robustus</i>	U	<i>Sebastes saxicola</i>	332
<i>Scopelogadus bispinosus</i>	U	<i>Sebastes variegatus</i>	U
Cetomimoidae	256	<i>Sebastes wilsoni</i>	U
Barbourisiidae		<i>Sebastes zacentrus</i>	334
<i>Barbourisia rufa</i>	U	Sebastolobinae	336
Cetomimidae		<i>Sebastolobus alascanus</i>	336
<i>Cetomimus</i> sp.	U	<i>Sebastolobus altivelis</i>	336
<i>Cetostoma regani</i>	U	<i>Sebastolobus macrochir</i>	U
<i>Gyrinomimus</i> sp.	U	Anoplopomatoidei	
Rondeletiidae		Anoplopomatidae	
<i>Rondeletia loricata</i>	U	<i>Anoplopoma fimbria</i>	338
Zeiformes	257	<i>Erilepis zonifer</i>	U
Oreosomatidae		Hexagrammoidei	
<i>Alloctytus</i> sp.	258	Hexagrammidae	341
Gasterosteiformes	261	<i>Oxylebius pictus</i>	342
Aulorhynchidae		<i>Zaniolepis frenata</i>	344
<i>Aulorhynchus flavidus</i>	262	<i>Zaniolepis latipinnis</i>	344
Gasterosteidae		<i>Ophiodon elongatus</i>	346
<i>Gasterosteus aculeatus</i>	F	<i>Pleurogrammus monopterygius</i>	348
<i>Pungitius pungitius</i>	F	<i>Hexagrammos decagrammus</i>	352
Syngnathidae		<i>Hexagrammos lagocephalus</i>	354
<i>Syngnathus leptorhynchus</i>	264	<i>Hexagrammos octogrammus</i>	356
Scorpaeniformes	267	<i>Hexagrammos stelleri</i>	358
Scorpaenoidei		Cottoidei	
Scorpaenidae	268	Cottidae	360
Sebastinae	268	<i>Archaulis biseriatus</i>	U
<i>Sebastes aleutianus</i>	288	<i>Arteidiellichthys nigripinnis</i>	U
<i>Sebastes alutus</i>	U	<i>Arteidielliscus</i> sp.	U
<i>Sebastes auriculatus</i>	290	<i>Arteidiellus camchaticus</i>	U
<i>Sebastes aurora</i>	292	<i>Arteidiellus gomojunovi</i>	U
<i>Sebastes babcocki</i>	294	<i>Arteidiellus miacanthus</i>	U
<i>Sebastes borealis</i>	U	<i>Arteidiellus ochotensis</i>	U
<i>Sebastes brevispinis</i>	U	<i>Arteidiellus pacificus</i>	U
<i>Sebastes caurinus</i>	296	<i>Arteidiellus scaber</i>	U
<i>Sebastes chlorostictus</i>	U	<i>Arteidius corallinus</i>	420
<i>Sebastes ciliatus</i>	U	<i>Arteidius fenestralis</i>	414
<i>Sebastes crameri</i>	298	<i>Arteidius harringtoni</i>	416
<i>Sebastes diploproa</i>	300	<i>Arteidius lateralis</i>	418

<i>Artedius meanyi</i>	384	<i>Nautichthys oculo fasciatus</i>	454
<i>Artedius notospilotus</i>	420	<i>Nautichthys pribilovius</i>	U
<i>Ascelichthys rhodoros</i>	386	<i>Nautichthys robustus</i>	U
<i>Asemichthys taylori</i>	U	<i>Oligocottus maculosus</i>	430
<i>Blepsias bilobus</i>	448	<i>Oligocottus rimensis</i>	U
<i>Blepsias cirrhosus</i>	450	<i>Oligocottus snyderi</i>	432
<i>Chitonotus pugetensis</i>	388	<i>Paricelinus hopliticus</i>	404
<i>Clinocottus acuticeps</i>	422	<i>Phallocottus obtusus</i>	U
<i>Clinocottus embryum</i>	424	<i>Porocottus bradfordi</i>	U
<i>Clinocottus globiceps</i>	426	<i>Porocottus quadrifilis</i>	U
<i>Clinocottus recalvus</i>	428	<i>Psychrolutes paradoxus</i>	436
<i>Cottoid A</i>	438	<i>Psychrolutes phrictus</i>	U
<i>Cottus aleuticus</i>	U	<i>Radulinus asprellus</i>	406
<i>Cottus asper</i>	444	<i>Radulinus boleoides</i>	408
<i>Dasycottus setiger</i>	440	<i>Rhampnocottus richardsoni</i>	366
<i>Enophrys bison</i>	390	<i>Scorpaenichthys marmoratus</i>	382
<i>Enophrys diceraus</i>	U	<i>Sigmistes caulias</i>	U
<i>Enophrys lucasi</i>	U	<i>Sigmistes smithi</i>	U
<i>Eurymen gyrinus</i>	U	<i>Stelgistrum beringianum</i>	U
<i>Gilbertidia sigalutes</i>	434	<i>Stelgistrum concinnum</i>	U
<i>Gymnocanthus detriscus</i>	U	<i>Sternias xenostethus</i>	U
<i>Gymnocanthus galeatus</i>	U	<i>Stlegicottus xenogrammus</i>	U
<i>Gymnocanthus pistilliger</i>	U	<i>Synchirus gilli</i>	410
<i>Gymnocanthus tricuspis</i>	394	<i>Taurocottus brashnikovi</i>	U
<i>Hemilepidotus gilberti</i>	372	<i>Thecopterus aleuticus</i>	U
<i>Hemilepidotus hemilepidotus</i>	374	<i>Thyriscus anoplus</i>	U
<i>Hemilepidotus jordani</i>	376	<i>Triglops forficata</i>	U
<i>Hemilepidotus papilio</i>	U	<i>Triglops jordani</i>	U
<i>Hemilepidotus spinosus</i>	378	<i>Triglops macellus</i>	U
<i>Hemilepidotus zapus</i>	380	<i>Triglops metopias</i>	U
<i>Hemitripterus bolini</i>	U	<i>Triglops pingeli</i>	U
<i>Hemitripterus villosus</i>	452	<i>Triglops szepticus</i>	U
<i>Icelinus borealis</i>	U	<i>Zesticelus profundorum</i>	U
<i>Icelinus burchami</i>	U	Agonidae	456
<i>Icelinus filamentosus</i>	U	<i>Agonomalus mozinoi</i>	458
<i>Icelinus oculatus</i>	U	<i>Agonopsis vulsa</i>	U
<i>Icelinus tenuis</i>	U	<i>Agonus acipenserinus</i>	U
<i>Icelus canaliculatus</i>	U	<i>Agonus decagonus</i>	U
<i>Icelus euryops</i>	U	<i>Anoplagonus inermis</i>	U
<i>Icelus scutigera</i>	U	<i>Aspidophoroides bartoni</i>	U
<i>Icelus spatula</i>	U	<i>Aspidophoroides olriki</i>	U
<i>Icelus spiniger</i>	U	<i>Bathyagonus alascanus</i>	U
<i>Icelus uncinalis</i>	U	<i>Bathyagonus infraspinus</i>	U
<i>Jordania zonope</i>	U	<i>Bathyagonus nigripinnis</i>	U
<i>Leptocottus armatus</i>	446	<i>Bathyagonus pentacanthus</i>	U
<i>Malacocottus kincaidi</i>	U	<i>Bothragonus swani</i>	460
<i>Malacocottus sp.</i>	U	<i>Hypsagonus quadricornis</i>	U
<i>Malacocottus zonurus</i>	442	<i>Ocella dodecaedron</i>	U
<i>Megalocottus laticeps</i>	U	<i>Ocella impi</i>	U
<i>Megalocottus platycephalus</i>	U	<i>Ocella verrucosa</i>	U
<i>Microcottus sellaris</i>	U	<i>Odontopyxis trispinosa</i>	U
<i>Myoxocephalus axillaris</i>	U	<i>Pallasina barbata</i>	U
<i>Myoxocephalus brandtii</i>	U	<i>Percis japonicus</i>	U
<i>Myoxocephalus jaok</i>	U	<i>Sarritor frenatus</i>	U
<i>Myoxocephalus niger</i>	U	<i>Sarritor leptorhynchus</i>	U
<i>Myoxocephalus polyacanthocephalus</i>	U	<i>Stellerina xyosterna</i>	U
<i>Myoxocephalus quadricornis</i>	U	<i>Xeneretmus latifrons</i>	462
<i>Myoxocephalus scorpioides</i>	U	<i>Xeneretmus leiops</i>	U
<i>Myoxocephalus scorpius</i>	402	<i>Xeneretmus triacanthus</i>	U
<i>Myoxocephalus sp.</i>	U	Cyclopteridae	465
<i>Myoxocephalus stelleri</i>	U	Cyclopterinae	465
<i>Myoxocephalus verrucosus</i>	U	<i>Aptocyclus ventricosus</i>	466

<i>Cyclopteropsis phrynoides</i>	U	<i>Liparis pulchellus</i>	U
<i>Eumicrotremus andriashevi</i>	U	<i>Liparis rutteri</i>	U
<i>Eumicrotremus barbatus</i>	U	<i>Liparis tunicatus</i>	U
<i>Eumicrotremus birulai</i>	U	<i>Lipariscus nanus</i>	U
<i>Eumicrotremus derjugini</i>	U	<i>Nectoliparis pelagicus</i>	476
<i>Eumicrotremus gyrynops</i>	U	<i>Odontoliparis ferox</i>	U
<i>Eumicrotremus orbis</i>	468	<i>Osteodiscus cascadiae</i>	U
<i>Eumicrotremus soldatovi</i>	U	<i>Paraliparis cephalus</i>	U
<i>Eumicrotremus taranetzi</i>	U	<i>Paraliparis dactylosus</i>	U
<i>Lethotremus muticus</i>	U	<i>Paraliparis deani</i>	478
<i>Pelagocyclus vitiazi</i>	U	<i>Paraliparis holomelas</i>	U
Liparidinae	470	<i>Paraliparis latifrons</i>	U
<i>Acantholiparis caecus</i>	U	<i>Paraliparis megalopus</i>	U
<i>Acantholiparis opercularis</i>	U	<i>Paraliparis melanobranchus</i>	U
<i>Acantholiparis</i> sp.	U	<i>Paraliparis mento</i>	U
<i>Careproctus abbreviatus</i>	U	<i>Paraliparis paucidens</i>	U
<i>Careproctus attenuatus</i>	U	<i>Paraliparis pectoralis</i>	U
<i>Careproctus bowersianus</i>	U	<i>Paraliparis rosaceus</i>	U
<i>Careproctus cameliae</i>	U	<i>Paraliparis ulochir</i>	U
<i>Careproctus canus</i>	U	<i>Polypera beringiana</i>	U
<i>Careproctus colletti</i>	U	<i>Polypera greeni</i>	U
<i>Careproctus cypselurus</i>	U	<i>Rhinoliparis attenuatus</i>	U
<i>Careproctus ectenes</i>	U	<i>Rhinoliparis barbulifer</i>	480
<i>Careproctus filamentosus</i>	U	<i>Temnocora candida</i>	U
<i>Careproctus furcellus</i>	U	Perciformes	483
<i>Careproctus gilberti</i>	U	Percoidei	
<i>Careproctus longifilis</i>	U	Carangidae	
<i>Careproctus melanurus</i>	U	<i>Naucrates ductor</i>	484
<i>Careproctus microstomus</i>	U	<i>Seriola lalandi</i>	S
<i>Careproctus mollis</i>	U	<i>Trachurus symmetricus</i>	486
<i>Careproctus opisthotremus</i>	U	Bramidae	
<i>Careproctus oregonensis</i>	U	<i>Brama japonicus</i>	488
<i>Careproctus ostentum</i>	U	<i>Taractes asper</i>	U
<i>Careproctus ovigerum</i>	U	Caristiidae	
<i>Careproctus pellucidus</i>	U	<i>Caristius macropus</i>	490
<i>Careproctus phasma</i>	U	Sciaenidae	
<i>Careproctus pycnosoma</i>	U	<i>Atractoscion nobilis</i>	S
<i>Careproctus rastrinus</i>	U	<i>Genyonemus lineatus</i>	492
<i>Careproctus scottae</i>	U	<i>Seriphus politus</i>	S
<i>Careproctus simus</i>	U	Pentacerotidae	
<i>Careproctus</i> sp.	U	<i>Pseudopentaceros</i> sp.	494
<i>Careproctus spectrum</i>	U	Embiotocidae	
<i>Careproctus zachirus</i>	U	<i>Amphistichus koelzi</i>	V
<i>Crystallichthys cyclospilus</i>	U	<i>Amphistichus rhodoterus</i>	V
<i>Crystallichthys mirabilis</i>	U	<i>Brachyistius frenatus</i>	V
<i>Elassodiscus caudatus</i>	U	<i>Cymatogaster aggregata</i>	V
<i>Elassodiscus tremebundus</i>	U	<i>Embiotoca lateralis</i>	V
<i>Gyrinichthys minytremus</i>	U	<i>Hyperprosopon argenteum</i>	V
<i>Liparis bristolensis</i>	U	<i>Hyperprosopon ellipticum</i>	V
<i>Liparis callyodon</i>	U	<i>Hypocritichthys analis</i>	V
<i>Liparis catharus</i>	U	<i>Phanerodon furcatus</i>	V
<i>Liparis cyclopus</i>	U	<i>Rhacochilus vacca</i>	V
<i>Liparis dennyi</i>	U	Zoarcoidei	
<i>Liparis florae</i>	U	Bathymasteridae	496
<i>Liparis fucensis</i>	U	<i>Bathymaster caeruleofasciatus</i>	U
<i>Liparis gibbus</i>	U	<i>Bathymaster leurolepis</i>	U
<i>Liparis grebnitzki</i>	U	<i>Bathymaster signatus</i>	U
<i>Liparis mednius</i>	U	<i>Ronquilus jordani</i>	U
<i>Liparis megacephalus</i>	U	Zoarcidae	498
<i>Liparis micraspidophorus</i>	U	<i>Bothrocara brunneum</i>	U
<i>Liparis mucosus</i>	U	<i>Bothrocara hollandi</i>	U
<i>Liparis ochotensis</i>	U	<i>Bothrocara molle</i>	U

<i>Bothrocara pusillum</i>	U	<i>Anisarchus medius</i>	U
<i>Bothrocara remigerum</i>	U	<i>Lumpenella longirostris</i>	U
<i>Derepodichthys alepidotus</i>	U	<i>Lumpenus fabricii</i>	U
<i>Gymnelus bilabrus</i>	U	<i>Lumpenus maculatus</i>	U
<i>Gymnelus hemifasciatus</i>	U	<i>Lumpenus medius</i>	U
<i>Gymnelus popovi</i>	U	<i>Lumpenus sagitta</i>	U
<i>Gymnelus viridus</i>	U	<i>Poroclinus rothrocki</i>	U
<i>Krusensterniella pavlovskii</i>	U	Opisthocentrini	504
<i>Lycenchelys altus</i>	U	<i>Allolumpenus hypochromus</i>	U
<i>Lycenchelys camchaticus</i>	U	<i>Opisthocentrus ocellatus</i>	508
<i>Lycenchelys crotalinus</i>	U	<i>Plectobranthus evides</i>	510
<i>Lycenchelys hippopotamus</i>	U	Xiphisterinae	
<i>Lycenchelys jordani</i>	U	Alectrini	504
<i>Lycenchelys longirostris</i>	U	<i>Alectridium aurantiacum</i>	U
<i>Lycenchelys microporus</i>	U	<i>Anoplarchus insignis</i>	U
<i>Lycenchelys pliciferus</i>	U	<i>Anoplarchus purpurescens</i>	512
<i>Lycenchelys rassi</i>	U	Xiphisterini	504
<i>Lycenchelys ratmanovi</i>	U	<i>Cebidichthys violaceus</i>	U
<i>Lycenchelys roseus</i>	U	<i>Phytichthys chirus</i>	514
<i>Lycenchelys volki</i>	U	<i>Xiphister atropurpureus</i>	516
<i>Lycodapus derjugini</i>	U	<i>Xiphister mucosus</i>	518
<i>Lycodapus dermatinus</i>	U	Cryptacanthodidae	
<i>Lycodapus endemoscotus</i>	U	<i>Delolepis gigantea</i>	520
<i>Lycodapus fierasfer</i>	U	<i>Lyconectes aleutensis</i>	520
<i>Lycodapus leptus</i>	U	Pholididae	522
<i>Lycodapus mandibularis</i>	U	<i>Apodichthys flavidus</i>	U
<i>Lycodapus pachysoma</i>	U	<i>Pholis clemensi</i>	U
<i>Lycodapus parviceps</i>	U	<i>Pholis dolichogaster</i>	U
<i>Lycodapus poecilis</i>	U	<i>Pholis fasciata</i>	U
<i>Lycodapus psarosomatus</i>	U	<i>Pholis gilli</i>	U
<i>Lycodes brevipes</i>	U	<i>Pholis laeta</i>	U
<i>Lycodes concolor</i>	U	<i>Pholis ornata</i>	U
<i>Lycodes cortezianus</i>	U	<i>Pholis schultzi</i>	U
<i>Lycodes diapterus</i>	U	<i>Xererpes fucorum</i>	U
<i>Lycodes mucosus</i>	U	Anarhichantidae	
<i>Lycodes pacifica</i>	U	<i>Anarhichas orientalis</i>	524
<i>Lycodes palearis</i>	U	<i>Anarrhichthys ocellatus</i>	526
<i>Lycodes raridens</i>	U	Ptilichthyidae	
<i>Lycodes turneri</i>	U	<i>Ptilichthys goodei</i>	528
<i>Lyconema barbatum</i>	U	Zaproridae	
<i>Melanostigma pammelas</i>	U	<i>Zaprora silenus</i>	530
<i>Nalbantichthys elongatus</i>	U	Scytalinidae	
<i>Opaeophacus acrogeneius</i>	U	<i>Scytalina cerdale</i>	U
<i>Pachycara bulbiceps</i>	U	Trachinoidei	
<i>Puzanovia rubra</i>	U	Trichodontidae	
<i>Taranetzella lycoderma</i>	U	<i>Arctoscopus japonicus</i>	U
Stichaeidae	500	<i>Trichodon trichodon</i>	532
Stichaeinae		Blennioidei	
Stichaeini	500	Clinidae	535
<i>Eumesogrammus praecisus</i>	U	<i>Gibbonsia metzi</i>	U
<i>Stichaeus punctatus</i>	U	<i>Gibbonsia montereyensis</i>	U
Chirolophini	500	<i>Heterostichus rostratus</i>	536
<i>Bryzoichthys lysimus</i>	U	Icosteoidae	
<i>Bryzoichthys marjorius</i>	U	Icosteidae	
<i>Chirolophis decoratus</i>	U	<i>Icosteus aenigmaticus</i>	538
<i>Chirolophis nugator</i>	U	Ammodytoidei	
<i>Chirolophis snyderi</i>	U	Ammodytidae	
<i>Chirolophis tarsodes</i>	U	<i>Ammodytes hexapterus</i>	540
<i>Gymnoclinus cristulatus</i>	U	Gobioidei	
Lumpeninae		Gobiidae	543
Lumpenini	502	<i>Clevelandia ios</i>	544
<i>Acantholumpenus mackayi</i>	U	<i>Coryphopterus nicholsi</i>	546

<i>Lepidogobius lepidus</i>	548	<i>Atheresthes evermanni</i>	576
Scombroidei		<i>Atheresthes stomias</i>	578
Trichiuridae	551	<i>Clidoderma asperrimum</i>	U
<i>Aphanopus carbo</i>	U	<i>Embassichthys bathybius</i>	580
<i>Benthodesmus elongatus</i>	552	<i>Eopsetta jordani</i>	582
<i>Benthodesmus tenuis</i>	U	<i>Glyptocephalus stelleri</i>	584
<i>Lepidopus fitchi</i>	U	<i>Glyptocephalus zachirus</i>	586
Scombridae		<i>Hippoglossoides elassodon</i>	588
<i>Euthynnus pelamis</i>	S	<i>Hippoglossoides robustus</i>	590
<i>Sarda chiliensis</i>	S	<i>Hippoglossus stenolepis</i>	592
<i>Scomber japonicus</i>	554	<i>Inopsetta ischrya</i>	U
<i>Thunnus alalunga</i>	S	<i>Isopsetta isolepis</i>	594
<i>Thunnus albacares</i>	S	<i>Lepidopsetta bilineata</i>	596
<i>Thunnus obesus</i>	S	<i>Lepidopsetta 2</i>	599
<i>Thunnus thynnus</i>	S	<i>Limanda aspera</i>	602
Acanthuroidei		<i>Limanda proboscidea</i>	U
Luvaridae*		<i>Limanda sakhalinensis</i>	U
<i>Luvarus imperialis</i>	556	<i>Liopsetta glacialis</i>	604
Stromateoidei		<i>Lyopsetta exilis</i>	606
Centrolophidae		<i>Microstomus pacificus</i>	608
<i>Icichthys lockingtoni</i>	558	<i>Parophrys vetulus</i>	610
Tetragonuridae		<i>Platichthys stellatus</i>	612
<i>Tetragonurus cuvieri</i>	560	<i>Pleuronectes quadrituberculatus</i>	614
Stromateidae		<i>Pleuronichthys coenosus</i>	616
<i>Peprilus simillimus</i>	562	<i>Pleuronichthys decurrens</i>	618
Pleuronectiformes	565	<i>Psettichthys melanostictus</i>	620
Paralichthyidae		<i>Reinhardtius hippoglossoides</i>	622
<i>Citharichthys sordidus</i>	568	Cynoglossidae	
<i>Citharichthys stigmaeus</i>	570	<i>Symphurus atricauda</i>	624
<i>Paralichthys californicus</i>	S	Tetraodontiformes	627
Pleuronectidae	573	Molidae	
<i>Acanthopsetta nadeshnyi</i>	574	<i>Mola mola</i>	628

*Luvaridae is placed in the Scombroidei in J. Nelson (1984), but other studies (Leis and Richards 1984) have shown it belongs in the Acanthuroidei.

Identifying fish eggs and larvae

This guide is designed primarily to assist in identifying eggs and larvae of fishes collected by sampling plankton. This process is usually divided into two main steps: 1) Sorting fish material out of the total plankton, and 2) identifying fish eggs and larvae of interest. Sorting procedures such as outlined in Kramer et al. (1972) are usually used. Matarese and Sandknop (1984) discuss identification of fish eggs, and Powles and Markle (1984) discuss identification of fish larvae. Eggs should be fixed and maintained in 3-5% buffered formalin, whereas larvae are better preserved if transferred to 70% ethanol after fixation in 3-5% buffered formalin (see Lavenberg et al. 1984). Identification is usually accomplished using a dissecting microscope at magnifications of 7 to 70 \times , with lighting from above and below a clear or frosted glass stage. A calibrated ocular micrometer is needed for various measurements of eggs and larvae. Specimens are frequently examined as whole mounts in their preservation medium in dishes or watch glasses (\sim 50 mm diameter). Fine probes and forceps can be used carefully to manipulate the specimens. A piece of white paper inserted between the specimen dish and the stage makes pigment more readily visible.

Since the larvae, and certainly the eggs, do not look like their parents, different characters are required to identify them. In practice, it is often convenient to separate the eggs and larvae in a sample into similar-appearing types, before recording specific observations or trying to identify them. Generally, it is best to identify all specimens in a sample to the lowest taxonomic level possible, even if the collection was intended for certain target species; without careful examination, other species could be misidentified as those of a target species.

In addition to characters of the eggs and larvae themselves, clues to their identity are found in information such as geographic area, depth, and date of collection.

Eggs

For pelagic fish eggs, the following identifying characters are recorded as known in this guide (see Matarese and Sandknop 1984 for more details): **Shape** (most are spherical, a few ellipsoidal); **size** (most are \sim 1 mm in diameter, range \sim 0.5-4.0 mm for pelagic eggs covered in this guide); **oil globules** (present in many species, vary in number and size); **yolk** (homogeneous or segmented); **chorion** (smooth in most, but sculptured in some); **perivitelline space** (narrow [$<$ 0.1 egg diameter] in many, but reaches one-half egg diameter in some); and **embryonic characters** (shape, pigment, gut length, early fin development, and myomere count of the embryo in late-stage eggs). Few taxonomic generalizations can be made for fish eggs; closely related species may have eggs that are as dissimilar as species in different orders (Table 2). However, the yolk is usually segmented in primitive teleosts (e.g., clupeiforms, anguilliforms, and salmoniforms) and homogeneous in more advanced forms. Also, oil globules are absent in pleuronectids covered in this guide.

Larvae

Fish larvae possess a larger suite of characters than the eggs, and may undergo dramatic changes with development. To begin to identify a larval fish, the following should be noted: **Size** (NL or SL) and **stage of development** (e.g., preflexion, flexion, postflexion, transforming [Kendall et al. 1984]) as well as its **general appearance** (e.g., long and slender, short and deep bodied, lightly or

heavily pigmented). Figure 2 illustrates anatomical features and measurements used to describe larvae. With practice, it is often possible to group larvae according to order or family (Fig. 3).

Meristic characters Meristic characters are essential and should be determined, understanding that fin elements are gradually added during larval development. Myomeres are the first meristic character to stabilize, and the number usually reflects the number of adult vertebrae. The number of vertebrae varies among the fishes covered in this guide from $<$ 20 (molds) to $>$ 200 (e.g., most elopomorphs) (see Figure 4 where, for fishes occurring in the study area, vertebral counts are arranged by family in ascending order). Use of polarized light often facilitates counting myomeres. Myosepta are frequently more discernable than the myomeres, and if they are counted, one should be added to account for the myomeres anterior and posterior to the first and last myosepta.

The developing median fins contain several bits of taxonomic information. Dipping the larva in a potassium hydroxide solution and then in an Alizarin Red solution (see Hollister 1934 for solution recipes) for a few seconds, then rinsing and examining it in its preservative allows fin rays and head spines to be seen more clearly, since they stain red. However, this procedure may interfere with future attempts to clear and stain the larva. The principal caudal fin count is often an ordinal character (Table 2) and is very useful and relatively easy to determine in larvae, since it generally reaches its adult state shortly after flexion. The number, position, and order of development of the dorsal and anal fins, and their composition in terms of spines and soft rays, are important characters (Table 2). In several fishes (e.g., *Sebastes*), the final dorsal and anal spine develop first as soft rays, and change to spines during transformation. Thus, in late larvae, the median fins are different in the composition of spines and soft rays than in the adults. Fin rays that become elongate in larvae often develop precociously, otherwise second dorsal and anal soft rays generally develop concurrently when present. Spinous dorsal fins may develop before, concurrently, or after the second dorsal fin. When there is a gap between the spinous and soft-ray portions of the dorsal fin, spines generally develop after the soft rays.

In the sequence of paired fin development, the pectorals often develop early in the larval period, and the pelvics late. The pectoral fins form in the egg as larval pectoral fins, without fin rays. The fin rays generally develop much later in the larval stage. The length of the pectoral fin, as well as its number of rays, is a useful character. While the number of pectoral rays may vary within and among species in a genus, the pelvic fin position and formula are generally stable at a high level of classification (e.g., order; see Table 2). The pelvic fin is absent in anguilliforms and members of some other groups (e.g., zoarcids, ammodytids, and molds). In primitive fishes (e.g., clupeiforms, myctophiforms) it contains no spines, is abdominal, and generally contains more than five soft rays. In most perciforms and scorpaeniforms, the basic pelvic fin count is I,5, and it is thoracic in position. This count is reduced in some; notably in the northern Pacific, cottids often have fewer than five soft rays. The pelvic fin is modified into a sucking disc in gobioid forms and in some cyclopterids.

Other meristic characters (e.g., gill rakers, secondary caudal fin rays, scales) develop too late to be generally useful in identifying larvae, but may be essential when working with pelagic juveniles.

Pigmentation Pigmentation available as taxonomic characters on larvae is limited to melanophores, since other pigment cells (e.g., xanthophores) do not retain their color in currently-used fixatives and preservatives. Melanophore patterns are very useful for identifying larval fishes. The relative size, position, and sometimes the number of melanophores in series should be noted. In some cases, pigmentation consists of a group of melanophores in a specific area; in others the pigmentation consists of an individual melanophore. Pigmentation generally changes as larvae develop. Movement of individual melanophores is rather limited, but addition or loss of melanophores is common. Usually preflexion larvae are less pigmented than later larvae, and late in the larval period, as transformation occurs, the larval pigment pattern is overgrown by the largely superficial pattern of the juvenile. Between the preflexion and transformation stages in most fish there is a definite larval pigment pattern which is relatively stable, and unique to a species in many cases. Although the position of melanophores is a species characteristic, the degree of contradiction seems to be physiologically moderated. Thus, larvae of the same species could have a different overall pigmented appearance, lighter or darker. Several species in the Northeast Pacific have heavily pigmented larvae, which are readily recognized in samples but may be confused with one another (Table 3, Fig. 5).

Morphology Larval shape can vary from stout and robust to quite slender and elongate (Table 2). Several fishes in the study area have elongate larvae which may be confused with one another (Table 4, Fig. 6). The ratio of body depth at the pectoral fin to standard length is usually sufficient to characterize overall body shape. The size and shape of the head and eye may also be important. The length of the gut, measured as the ratio of preanal to standard length, is quite useful. As with other characters, larval shape characters vary with development, so the size and stage of development should be noted when comparing shape of an unknown larva to illustrations and descriptions of known specimens.

Head spines, when present, may be more numerous and accentuated in larvae than in adults. Among Northeast Pacific fishes, larval head spines are most prevalent in cottids and scorpaenids, although they also occur in some members of groups such as perciforms and pleuronectids (Table 2).

Once the above data on an unknown specimen or group of similar specimens are assembled, actual identification becomes largely a process of eliminating species whose characters do not match the unknown specimen. Keys are not presented in this guide and generally do not work well with fish larvae, because the larvae change so much with development, and the larvae of all species in a study area are rarely known. With larvae, particularly, the first attempt should be to identify the unknown to order or family, based on meristic values, shape, and general appearance. Table 2 and the figures of larval representatives of various orders (Fig. 4) should be of assistance in this. Once an idea of the appropriate order/suborder is established, more detailed information can be obtained in the material at the beginning of each ordinal account, including general life-history characters, species represented in the study area, and meristic and early-life-history characters. Illustrations, meristics, and other information given in the individual taxon accounts in this guide should then be compared with the unknown specimens to find the most likely species. Differences between the unknown and described larvae should be noted. If an unknown specimen does not match any larval descriptions given here, check meristic tables of the most likely taxa to see if the unknown specimen fits a species whose larvae have not yet been described.





Figure 2

Examples of features used to describe early stages of fishes: A, egg; B, preflexion larva; C, late larva showing base points for measurements; D, late larva showing morphological features (B-D, after Fahay 1983).



Elopomorpha




Clupeiformes



Argentinoidei



Salmonoidei




Stomiiformes



Stomiiformes



Aulopiformes




Myctophiformes




Gadiformes




Ophidiiformes



Lophiiformes




Gobiesociformes




Beloniformes




Atheriniformes



Lampriformes



Beryciformes



Zeiformes


Figure 3
Representative postflexion larvae of higher categories of fishes of the Northeast Pacific Ocean.



Gasterosteiformes



Gasterosteiformes




Scorpaenoidei




Anoplopomatoidei



Hexagrammoidei



Cottoidei



Percoidei




Zoarcoidei



Trachinoidei




Blennioidei




Icosteoidi




Ammodytoidei




Gobioidei




Scombroidei



Stromateoidi



Pleuronectiformes



Tetraodontiformes

Figure 3 (continued)

Table 2
Ordinal/subordinal egg, larval, and meristic characters of Northeast Pacific fishes (Ahlstrom and Moser 1976, Fahay 1983, Leis and Rennis 1983, in part).

Taxon	Families in study area	Eggs													Larvae										Meristics																			
		Development site		Shape	Chorion	Yolk	Perivitelline space	Oil globule		Diameter (mm)	Preanal length (%SL)		Gut shape	Eye shape	Head spines	Transformation	Special juvenile	Fin characters		Vertebrae		Principal caudal fin rays																						
		Planktonic	Demersal	Spherical	Ellipsoidal	Smooth	Sculptured	Segmented	Homogeneous	Wide	Moderate	Narrow	Absent	1 Present	>1	<0.9	1.0-2.0	>2.0	Elongate	Stocky	<50	50-75	>75	Straight	Coiled	Trailing	Round	Narrowed	Absent	Present	Marked	Gradual	Present	Present	Absent	Absent	Abdominal	Thoracic	Jugular	No. dorsal fins	Adipose fin	Upper	Lower	Total
Notacanthiformes/Anguilliformes	9	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	68-750	0-9	0-7	0-16			
Clupeiformes	2	X X ^a	X X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	43-57	10	9	19			
Salmoniformes																																												
Argentinoidei	5	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	34-84	10	9	19		
Salmonoidei	2	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	54-75	10	9	19		
Stomiiformes	6	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	35-83	10	9	19		
Aulopiformes	7	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	48-87	10	9	19		
Myctophiformes	2	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	29-42	10	9	19		
Gadiformes	5	X X	X	X X ^b	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	48-64;84-86	4-6 ^d	2-4 ^d	6-10 ^d		
Ophidiiformes	3	Unknown										X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	60-81	4	5	9				
Lophiiformes	2	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	19-21	4	4-5	8-9	
Gobiesociformes	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	32-36	?	?	14	
Belontiiformes ^f	1	X X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	62-69	7	8	15	
Atheriniformes	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	44-52	9	8	17	
Lampriformes	2	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	43-46;90-94	?	?	?	
Beryciformes	5	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	23-52	10	9	19	
Zeiformes	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	39	6	7	13	
Gasterosteiformes	3	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	30-64	6-7	6	12-13	
Scorpaeniformes																																												
Scorpaenoidei	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	26-31	7	7	14	
Anoplopomatoidei	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	61-66	7	7	14	
Hexagrammoidei	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	36-63	7-8	6-11	13-19	
Cottoidei	3 ^h	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	25-71	6	5-7	11-13	
Perciformes																																												
Percoidae	6	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	23-41	9	8	17	
Zoarcoidei	9	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	49-150;221-250	?	?	12-15	
Trachinoidei	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	44-47	?	?	13	
Blennioidei	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	55-58	9	8	17	
Icostoidei	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	66-68	9	8	17	
Ammodytoidei	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	65-70	9	8	17	
Gobioidei	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	26-38	?	?	17	
Scombroidei	2	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	22-23;30-32;148-150	9	8	17	
Acanthuroidei	1	Unknown										X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	22-23	?	?	16				
Stromateoidei	3	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	28-31;52-62	9	8	17	
Pleuronectiformes	3	X X ^m	X	X X ⁿ	X	X ^o	X	X ^p	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	35-66	9-12 ^q	8-11 ^q	17-24 ^q	
Tetraodontiformes	1	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	17-18			15 ^r

^a *Clupea pallasii* eggs demersal.

^b Macrouridae eggs sculptured.

^c Three dorsal fins in Gadidae.

^d Macrouridae lack caudal fins.

^e Pelvic fin modified into a disc.

^f J. Nelson (1984) places scomberesocids within the Cyprinodontiformes.

^g Dorsal spines isolated from soft-rayed dorsal fin.

^h J. Nelson (1984) considers Psychrolutidae a separate family; we include it in Cottidae.

ⁱ Pelvic fin absent or modified into a disc in Cyclopteridae.

^j *Trachurus* yolk segmented.

^k Pelvic fin present only in larvae.

^l Pelvic fin absent in *Peprilus*.

^m *Lepidopsetta bilineata* only species with demersal egg.

ⁿ *Pleuronichthys* eggs have hexagonal sculpturing.

^o *Hippoglossoides elassodon* and *H. robustus* only species with wide perivitelline space.

^p *Pleuronectidae* eggs have no oil globule.

^q Total caudal fin rays.

^r *Mola mola* has a pseudocaudal fin (see Tyler 1980).




Figure 4
 Ranges of total vertebral counts for families in the study area.

Table 3
Distinguishing features of some commonly collected heavily pigmented larvae.*


Taxon	Distribution	Total vertebrae	Description of pigment	Other diagnostic features	Page reference
Myomeres 25-50					
<i>Rhamphocottus richardsoni</i>	SSC-Bering Sea	26-28	Small spots, densely distributed on head, gut, and 3/4 body; absent on peduncle and tail	Subterminal mouth, deep body; after flexion head spines and prickles by 9-10 mm	366
<i>Myoxocephalus</i> G	Wash.-Bering Sea	34-37	Spots concentrated dorsally on head and gut, and over anterior 1/2 body; absent over gut	Head spines develop during flexion	400
<i>Hemilepidotus</i> spp.	SSC-SE Alaska	35-39	Spots widely distributed on head, gut, and body	Head spines develop during flexion	372
<i>Scorpaenichthys marmoratus</i>	SSC-SE Alaska	35-37	Dense, uniformly distributed spots over head, gut, and 3/4 body; absent on tail	Head bumps and preopercular spines develop after flexion; preanal finfold	382
<i>Blepsias</i> spp.	Cent. Calif.-Bering Sea	37-39	Dense, large spots evenly concentrated over dorsal head, gut, and >3/4 body; absent on tail tip	Head spines and bumps develop during flexion	448
<i>Pleuromichthys</i> spp.	SSC-Bering Sea	37-41	Dense spots concentrated over head, gut, and >3/4 body; dorsal and anal finfold	Slender body with wide finfolds	616
<i>Radulinus</i> spp.	SSC-Gulf of Alaska	38-40	Widely distributed over lateral gut surface and almost 3/4 body	Preopercular spines not prominent, gut coiled	406
<i>Hemitripteris villosus</i>	Gulf of Alaska-Bering Sea	39-41	Spots evenly distributed to 3/4 body and into dorsal and anal finfold	Large size at development	452
<i>Nautichthys oculofasciatus</i>	Cent. Calif.-Bering Sea	40-41	Large, widely distributed spots to 3/4 body; dorsal and anal finfold pigment at midbody	Long, pigmented precocious pectorals; head spines, bumps, ridges develop after flexion	454
<hr/>					
Myomeres >50					
<i>Hexagrammos</i> spp./ <i>Pleurogrammus monopterygius</i>	S. Calif.-Bering Sea	50-63	Widely distributed on dorsal head and gut, above and below notochord	Gut length <50% SL	348
<i>Ophiodon elongatus</i>	SSC-Gulf of Alaska	56-59	Widely distributed on dorsal head and gut, more concentrated on dorso- and ventrolateral surface; chin and isthmus	Pointed snout and large terminal mouth	346
<i>Zaprora silenus</i>	Cent. Calif.-Bering Sea	61-62	Small spots densely concentrated over entire body, except ventrally on gut, dorsal finfold, and posterior edge of gill cover; lightly on caudal peduncle and anal finfold	Large size at development	530
<i>Anoplopoma fimbria</i>	SSC-Bering Sea	61-66	Widely distributed over >3/4 body	>15 mm, pectoral fins long, pigmented; gut length >50% SL; anus curves ventrad	338
<i>Cololabis saira</i>	SSC-Bering Sea	62-69	Small spots densely concentrated, ventral tailtip without pigment	Long preanal finfold	236
<i>Lyconectes aleutensis</i>	N. Calif.-Bering Sea	73-75	Heavily concentrated over >3/4 body, isthmus; lighter over gut and absent on peduncle	Large size at development	520
<i>Delolepis gigantea</i>	N. Calif.-Bering Sea	81-83	Widely distributed to 3/4 body, absent over gut and isthmus	Large size at development	520

*Other heavily pigmented larvae that may be less frequently encountered include the following species:


Anarhichas orientalis (p. 524)
Naucrates ductor (p. 484)
Peprilus simillimus (p. 562)
Oxylebius pictus (p. 342)
Zaniolepis (p. 344)

HEAVILY PIGMENTED LARVAE (MYOMERES 25–50)


A *Rhamphocottus richardsoni*



B *Myoxocephalus* G



C *Hemilepidotus* spp. (*Hemilepidotus spinosus*)




Figures A, C, Richardson and Washington 1980; B, NWAFC original (B. Vinter).

Figure 5
Commonly collected heavily pigmented larvae with 25-50 myomeres (A-I), and >50 myomeres (J-P).


HEAVILY PIGMENTED LARVAE (MYOMERES 25–50)

D *Scorpaenichthys marmoratus*




8.6 mm NL

E *Blepsias* spp. (*Blepsias cirrhosus*)



11.5 mm SL

F *Pleuronichthys* spp. (*Pleuronichthys decurrens*)




6.5 mm NL

Figure D, Richardson and Washington 1980; E, NWAFC original (B. Vinter); F, Sumida et al. 1979.

Figure 5 (continued)


HEAVILY PIGMENTED LARVAE (MYOMERES 25–50)

G *Radulinus* spp. (*Radulinus asprellus*)




9.6 mm NL

H *Hemitripteris villosus*



14.4 mm SL

I *Nautichthys oculo-fasciatus*




8.3 mm SL

Figure G, Richardson and Washington 1980; H, Okiyama and Sando 1976; I, NWAFC original (B. Vinter).

Figure 5 (continued)


HEAVILY PIGMENTED LARVAE (MYOMERES > 50)

J *Hexagrammos* spp. (*Hexagrammos octogrammus*)




15.0 mm SL

K *Ophiodon elongatus*




12.2 mm SL

L *Zaprora silenus*



17.3 mm BL

M *Anoplopoma fimbria*




12.0 mm SL

Figures J–K, Kendall and Vinter 1984; L, Haryu and Nishiyama 1981; M, NWAFC original (B. Vinter).


Figure 5 (continued)

HEAVILY PIGMENTED LARVAE (MYOMERES > 50)


N *Cololabis saira*



O *Lyconectes aleutensis*



P *Delolepis gigantea*



Figures N–P, NWAFC originals (B. Vinter).

Figure 5 (continued)

Table 4
Comparison of selected diagnostic characters of some commonly collected families with similar-looking elongate larvae.

Taxon	Myomeres			Number between dorsal fin insertion and anal fin origin ^a (flexion stage)	Preanal length	Presence of adipose fin	Diagnostic pigment	Page reference
	Preanal	Postanal	Total					
Short dorsal and anal fin bases								
Clupeidae	28-32	19-23	46-57	6-8	80%	None	Gut	44
Engraulididae	24-26	19-21	43-47	0-2	67-75%	None	Isthmus, gut	48
Osmeridae ^b	—	—	54-73	9-11	80%	Yes	Single row pvm; ^c ventral gut midline	79
Long dorsal and anal fin bases								
Bathymasteridae	13-16	34-39	49-55	Overlaps	<50%	None	Urostyle or slash-like pigment along posterior hypaxial and epaxial myomeres	496
Stichaeidae	14-31	34-59	50-83	Overlaps	<50%	None	Gut, anus, pvm ^c	500
Pholididae	—	—	80-102	Overlaps	>50%	None	Gut, pvm ^c	522
Ammodytidae	40-47	23-25	65-74	Overlaps	60%	None	Double row pvm ^c	540


^aThe number of myomeres between dorsal and anal fins has been used as a taxonomic character in clupeiform larvae of certain size classes. During transformation the position of the gut and median fins shifts forward relative to myomere number (for more details, see McGowan and Berry 1984).

^bBased on larvae of *Mallotus villosus* (except for total myomere count); other osmerids, as yet unidentified to species, are similar.

^cpvm = postanal ventral midline melanophores.


ELONGATE LARVAE
Short dorsal and anal fin bases

A Clupeidae (*Clupea pallasii*)



19.0 mm SL

B Engraulididae (*Engraulis mordax*)



11.5 mm


C Osmeridae (*Mallotus villosus*)



16.0 mm


Long dorsal and anal fin bases

D Bathymasteridae (*Ronquilus jordani*)




10.4 mm SL

E Stichaeidae (*Anoplarchus purpurescens*)



12.0 mm SL

F Pholididae (*Pholis* sp.)



23.0 mm SL

G Ammodytidae (*Ammodytes hexapterus*)



16.0 mm

Figures A, D–F, NWAFC originals (B. Vinter); B, Kramer and Ahlstrom 1968; C, Fahay 1983 (after Templeman 1948); G, Stevens et al. 1984.

Figure 6
Commonly collected elongate larvae.

Format

This laboratory guide has been designed to be practical and easy to use. Only information deemed necessary for accurate and timely identification has been included. A two-page format is provided for each taxon where sufficient early-life-history information exists. The left page includes pertinent information for identification and the right page includes illustrations, usually with notations indicating important diagnostic features. The left page is divided into two columns: Information on meristic characters and life history features is presented on the left, and developmental information on the right. Blanks within the format indicate that particular information was not available (e.g., egg size, fecundity), to point out gaps in knowledge, and allow researchers to insert new information as it becomes available. The family name appears at the top of each page for quick reference. The phylogenetic sequence generally follows J. Nelson (1984) unless otherwise indicated. Species names along with their authorities appear at the upper right corner of the left page, and common names (from Robins et al. 1980, Hubbs et al. 1979, Shiino 1976, and others) are included at the upper left corner of the right page. Nomenclature generally follows Robins et al. (1980) and Steyskal (1980) unless a more recent revision is available (usually from Moser et al. 1984b). Exceptions are noted in the text.

For taxa treated at the species level, available illustrations are arranged on the right-hand page as follows: Late-stage egg is in the upper right-hand corner, and order of larvae is shown (top to bottom) as yolk sac, preflexion, flexion, postflexion, and either a transforming juvenile or special prejuvenile stage. Blank spaces indicate that stages were not available for illustration. When necessary for identification purposes, additional illustrations such as dorsal and ventral views are provided. Most illustrations were compiled from the literature; Moser et al. (1984b) provided over 100 illustrations. In a few cases, illustrations from the literature were redrawn, modified, or corrected; when this occurred, it is indicated. In addition to the published illustrations, original illustrations of 124 fish eggs and fish larvae by Beverly Vinter are included. Collection data are provided for original illustrations (Table 1).

Introductory sections are provided for each order and for taxa (usually genera or families) with difficult identification problems or for taxa that contain numerous similar species (e.g., Cottidae and *Sebastes*). Important diagnostic features are summarized, and, in some cases, tables are provided to aid in identification. In the Northeast Pacific Ocean, early-life-history stages of many species are either undescribed, incompletely described, or without adequate illustrations. For these taxa, identification material is provided at the lowest taxonomic level possible, usually family level (e.g., Osmeridae) or generic level (e.g., *Cyclothone*). Family level descriptions include a summary of available early and general life-history information, tables of meristic characters, and brief accounts of early-life-history characters from closely related taxa described from other geographic areas. For taxa where no early-life-history data are available (e.g., Cetomimoidae), life history summaries and meristic data are provided.

Meristics

Data summaries of meristic structures, except those for the caudal fin or from recently completed research, are from the NWAFC meristic database. The range (high and low value) and mode are presented for each entry. Ranges reported here generally represent the most extreme values ever recorded; in many cases, these values are not likely to be observed. These values may have been seen on specimens collected outside our study area. Thus, in using this guide more emphasis should be placed on the reported modal values. An "X" appears when data are unavailable. For fin ray counts, R = rays (soft rays) and S = spines; for gill raker counts, U = upper and L = lower. In addition to pelvic fin-ray counts, fin position is given. Position is indicated by the following descriptors: Abdominal, thoracic, jugular, absent, or modified (e.g., pelvic disc in cyclopterids). The total vertebral count given may not equal the sum of the precaudal and caudal vertebral counts in some cases, since these counts may have originated from different sources.

Data for caudal fin-ray counts have been gleaned from published material, as well as from original observations. Caudal fin-ray counts are reported in the following sequence: Upper secondary, upper principal + lower principal, lower secondary, with ranges for each when available. Total caudal fin rays or total principal caudal fin rays are reported if no other data are available. Blanks in the caudal field indicate that no data are available.

For some taxa, it was necessary to forego the standard meristic format in order to present the data or provide additional information (e.g., members of the family Gadidae have three dorsal fins and two anal fins). Departures from the standard format are either explained on the page where they occur or in introductory sections preceding species descriptions for certain taxa.

General life history

Life history data are provided as ancillary information which may aid in identification of eggs and larvae. These data were extracted from Garrison and Miller (1982) and supplemented by the general literature and original, unpublished material.

Geographic ranges are from the NWAFC meristic database and the literature. Allen and Smith (1988) provided a significant amount of new information. Range information is restricted primarily to the study area. Thus, the limits of the southern range beyond the California-Mexico border, the northern range beyond the Arctic to the north and east, and the western range beyond the Bering Sea are not specified.

The following general locations are used to approximate geographic range within the study area (abbreviations in parentheses are used hereafter in the text when necessary):

South of southern California (SSC)
Southern California, 32-34°N (S. Calif. or S. California)
Central California, 34-38°N (Cent. Calif. or Cent. California)
Northern California, 38-42°N (N. Calif. or N. California)
Oregon, 42-46°N
Washington, 46-48°30'N (Wash.)
British Columbia, 48°30'-55°N (Brit. Col.)
Southeastern Alaska, 55-59°N (SE Alaska)
Gulf of Alaska, 54-60°N
Aleutian Islands, 51-55°N (Aleutian Is.)
Bering Sea, 54-66°N
Chukchi Sea, north of 66°N
Arctic

In addition to geographic range, general ecological descriptors are incorporated in the NWAFC meristic database. The following descriptors are used.

Pelagic environment

Nearshore shelf pelagic: Extends from the shore seaward to include waters overlying an ocean bottom <200 m. Equivalent to the neritic province of Hedgpeth (1957) and other authors.

Oceanic: Waters overlying an ocean bottom >200 m. The following subdivisions based on water depth are used: Epipelagic 0-200 m; Mesopelagic 200-1000 m; Bathypelagic >1000 m.

Benthic environment

Intertidal, nearshore: Extends from high tide to low tide. Equivalent to the littoral province of many authors.

Nearshore shelf demersal: All bottom from low tide to 200 m (= epibenthic in text). Equivalent to the sublittoral zone of Hedgpeth (1957) and other authors.

Mesobenthic: Deep sea beyond the continental shelf at depths of 200-500 m. Lowest part of the shelf and upper part of continental slope (Fedorov 1973).

Bathybenthic: Deep sea along the continental slope at depths of 500-2500 m. Middle and lower sections of continental slope (Fedorov 1973).

Freshwater or anadromous: Generally the mouth and lower reaches of rivers and streams.


Other information under **Life History** includes reproductive mode (e.g., viviparous, ovoviviparous, oviparous) and indicates whether eggs and larvae are pelagic or demersal. Data on spawning are divided into four categories: Season, area, mode, and migration. Spawning often varies among geographic regions and populations, so care has been taken to provide as much specific geographic information as available. Fecundity values are given as counts of ripening eggs in individual females: Total ranges or (in a few cases) as a function of length in the form $F = aL^b$. Ages at first maturity and longevity values were extracted from the general literature. If available, age/length differences between sexes are noted.

Early life history

Egg and oil globule diameter measurements in millimeters are usually given as ranges (high and low values) with modal values as available in parentheses after the range. Precision varies among literature sources, but specimens used for original measurements were measured to the nearest 0.1 mm. Egg diameters on the illustrations are as they were given in the source. Incubation time, when available, is given in number of days to hatching for a specific temperature (°C). When describing embryonic pigment patterns, emphasis was placed on those characteristics which aid in identification. Diagnostic characters usually provide a summary of important features and comparisons with similar fish eggs.

Figure 2 provides examples of features used to describe and identify early stages of fishes. Original measurements of larvae are in millimeters and given in standard length (SL). Some measurements extracted from the literature were given as body length (BL), notochord length (NL), total length (TL), head length (HL), or percentages of these. Preanal length is usually expressed as a percentage of SL. If specific values are not available, preanal lengths are given as <50% SL, 50-75% SL, or >75% SL. Definitions for developmental stages are from Ahlstrom et al. (1976) and Kendall et al. (1984). Transformation is defined as acquisition of the adult complement of fin rays, and in some taxa this is accompanied by squamation. Sequence of fin development is usually described as the order in which fin rays accept alizarin stain, inferring ossification. For some taxa the sequence is determined by completion of the ossification of a fin element rather than the initiation of ossification (e.g., G.D. Johnson 1984); this is noted in the text. When three or more fins develop simultaneously, semicolons are used to separate the sequence of formation of one or more fins (e.g., dorsal; anal; caudal, pectoral, and pelvic). Otherwise, commas are used to separate fins developing individually in sequence (e.g., dorsal, anal, caudal, pectoral, and pelvic). Precocious fin development is usually noted. When describing pigment, those melanophores or patterns of melanophores important in identifying the taxon are stressed. Generally only melanophores are mentioned, since other pigment is not visible in formalin-preserved specimens. The general appearance, shape, and relative size of melanophores or groups of melanophores are indicated by imprecise descriptors such as spot, patch, or blotch. Diagnostic pigment characteristics are also indicated directly on the illustration page. The description of pigment is often brief and in telegraphic style and is not meant to be a substitute for more detailed discussions available in complete early-life-history descriptions. References are provided when more complete early-life-history descriptions are available. Under **Diagnostic Characters**, a brief summary of key features is provided which may help to distinguish a larval specimen from other similar larvae in either closely related taxa or from morphologically similar groups. When possible, comparative information is included in tables and is cross-referenced.

Data from the files of the late E.H. Ahlstrom are footnoted as "E.H. Ahlstrom notes." These files contain original lecture notes for classes conducted on ichthyoplankton taxonomy between 1971 and 1977, early-life-history data for most orders of fishes, and notes on teleost caudal fins. The files were made available to Kendall & Matarese by H. Geoffrey Moser (Southwest Fish. Cent., Natl. Mar. Fish. Serv., NOAA), as authors contributing to *Ontogeny and Systematics of Fishes* (Moser et al. 1984b).



Elopomorpha: Notacanthiformes Anguilliformes

The notacanthiform fishes (spiny eels) and anguilliform fishes (true eels) generally occur worldwide. The spiny eels are primarily a deep-sea group and are distinguished from the true eels by a number of characters including the absence or reduction of the caudal fin, the presence of well developed pelvic fins and fin spines, a short dorsal fin, and the presence in larvae of a thin postcaudal filament. According to Castle (1984), the Notacanthiformes consist of 3 families, 6 genera, and about 22 species; the Anguilliformes, a much larger order, include 21 families, 153 genera, and about 720 species.

The early life history of eels has been studied for many years primarily due to the presence of a distinctive leptocephalus larval phase, but many species remain inadequately known. Eggs are generally large, pelagic, possess segmented yolks, and have one or more oil globules. Although few larvae have been collected, at least 12 taxa from at least 9 families are thought to occur in the study area. Since early life histories are incomplete, this section (except for the nemichthyids) will be described by family.

Families in study area: **Notacanthidae**
Xencongridae
Nemichthyidae
Cyematidae
Synaphobranchidae
Nettastomatidae
Congridae
Serrivomeridae

ELOPOMORPHA



Key to elopomorph leptocephali in the Northeast Pacific (after Smith 1979 and Fahay 1983, in part).

Table 5
Meristic characters of superorder Elopomorpha.

Taxon	Distribution	Vertebrae		Fins				
		Precaudal	Caudal (Total)	Dorsal	Anal	Pectoral	Pelvic	Caudal
Notacanthiformes								
Notacanthidae								
		(234-244)						
<i>Notacanthus chemnitzii</i>	Cent. Calif. ^a -Oregon	47-55		VII-X or	XI-XXV,	10-17	I-IV,	Absent
		(234-239)		IX, 2-3	115-132		6-10	
<i>Polyacanthonotus challengeri</i>	Oregon-Bering Sea	(242-244)		XXXII-XXXV	161-162	12-13	I, 8	Reduced
Anguilliformes								
Xenocoelidae								
		(97-163)						
<i>Thalassenchelys coheni</i>		67-74	83-92	280-350	218-260	Absent	Absent	Absent
		(142-163)						
Nemichthyidae								
		(170-400+)						
<i>Avocettina infans</i>	SSC-Aleutian Is.		(181-201)	300-350	265-270		Absent	Reduced
<i>Nemichthys larseni</i>	SSC-Oregon	79-84				10-12	Absent	
		(400-750)						
<i>Nemichthys scolopaceus</i>	SSC-Aleutian Is.		(293-750)	307-450	312-454	10-14	Absent	Reduced
Cyematidae								
		(74-108)						
<i>Cyema atrum</i>	SSC-Oregon	38-43		79-83	72-86	12-15	Absent	5
		(74- 80)						
Synphobranchidae								
		(126-172)						
<i>Histiobranchus bathybius</i>	Bering Sea		(126-151)	265-302	188-203	15-17	Absent	18
Netastomatidae								
		(186-290)						
<i>Venefica</i> sp. A	Cent. Calif.-Wash.		(199-224)	310+	325		Absent	12
Congridae								
		(120-261)						
<i>Xenomystax atrarius</i>	SSC-Brit. Col.	50-57	107-123	253-292	189-214	11-14	Absent	7-8
		(141-219)						
Serrivomeridae								
		(137-170)						
<i>Serrivomer jespersenii</i>	Brit. Col.	89-125		141-161	127-161	6-7	Absent	Reduced (6 principal)
		(147-169)						
Eurypharyngidae								
		(97-125)						
<i>Eurypharynx pelecanoioides</i>	N. Calif. ^b		(97-125)	155-196	118-147	11	Absent	Absent in adult

^aLea and Rosenblatt (1987)

^bOne specimen collected off northern California.

Anguilliformes

Primary characters used in identifying anguilliform leptocephali are the following (Ref: E.H. Ahlstrom notes, Castle 1984, Fahay 1983, Smith 1979):

Body shape May vary from slender to deep-bodied, and the tail may be tapered or rounded.

Head characteristics Size of the head relative to the body may vary; also important are head shape (blunt, sharp, or elongate), snout shape, nasal organ (size and position), eyes (round, narrow, or telescopic), and teeth (presence/absence, fanglike if present).

Number of myomeres Usually 100-250; exceptions include *Cyema* (<100) and *Nemichthys* (>750). Preanal and predorsal counts are also useful.

Gut characteristics Gut may be a simple straight tube or more complex with loops or swellings. Relative length of gut can range from <50% SL to >90% SL, although most eels have gut lengths between 50-70% SL.

Position of vertical blood vessels Variation in the position of the last blood vessel.

Pigmentation General body pigment may be located above/below the gut, along body midline, above/below notochord internally, or along the dorsal body margin. Pigment may be in the form of fine stippling or large stellate melanophores (blotches). Head pigment also varies.

Size at transformation Maximum size before transformation can vary from <100 mm SL (most families) to ~400 mm SL (nemichthyids).

Fins Dorsal and anal fins are confluent with the caudal fin, caudal fin is usually markedly reduced with 10 or fewer rays (about 5-11 with highest numbers in synphobranchids); pectoral fin is moderate, reduced, or lacking, and fin rays usually form late, i.e., after the leptocephalus stage. Pelvic fin is absent.

ELOPOMORPHA






Notacanthidae (234-244 myomeres)* *Notacanthus chemnitzi* and *Polyacanthonotus challengerii* both occur in the study area but their early life histories are unknown, as is the case of most notacanthids. *Leptocephalus giganteus* (identity unknown) from outside the study area is presented for comparison only. Notacanthid larvae are easily separable from larvae of true eels. Among the most notable characters are the following: Greatly elongate shape, thin postcaudal filament, and pigment which occurs in a ventral series.

Xenocoelidae (97-163 myomeres) *Thalassenchelys coheni* (142-163) may or may not belong with the xenocoelids. These larvae are quite unusual, with a short, deep body, rounded tail, and lack of pigment. They appear to be widely distributed in the northeastern Pacific from Washington to south of southern California.

Cyematidae (74-80 myomeres, most 74-78) *Cyema atrum* larvae are identified by their deep body shape, pointed head and tail, 3-4 gut loops on posterior half of gut, and low myomere count. Pigment is scattered over the lateral body surface, on the snout and lower jaw, and along the gut, especially on the loops. Other general features include gut length ~67% SL and dorsal fin origin approximately above anus. Size at transformation is 60-70 mm SL.

Synphobranchidae (subfamily Synphobranchinae) (126-151 myomeres) Synphobranchinae are represented in the Northeast Pacific by *Histiobranchus bathybius* but their larvae are unknown. Larvae in other members of the subfamily are identified by their telescopic eye, general lack of pigment, and the broad white stripe formed by an opaque central area of myomeres around the notochord. The body is moderately elongate and the head is short and pointed. The dorsal fin origin is anterior to the anus. The gut is usually relatively simple, unpigmented, and about 75% SL. Some taxa have loops in the gut. Ventral pigment is lacking, and lateral pigment is restricted to the postanal body. Some genera have a prominent pigment spot laterally, near the level of the anus. Size at transformation is 130-170 mm SL.

*Vertebral range is given for taxa in the study area. When no data are available, the range of counts for family is given (Table 5).



Figures A, D, Castle 1984; B, NWAFC original (B. Vinter); C, E, Smith 1984.

ELOPOMORPHA

Nettastomatidae (186-290 myomeres) Nettastomatids are represented in the Northeast Pacific by *Venefica* sp., but their larvae are unknown. Larvae in the family are identified by their gut characteristics (presence of two loops and gut length <50% SL) and moderately long head. The body is deep to elongate and the tail is pointed. The dorsal fin origin is posterior to the head but well anterior, at myomeres 11-12. Ventral and lateral pigment is variable. Size at transformation is also variable but occurs between 120 and 200 mm SL.

Congridae (141-219 myomeres) Congrids are represented in the area by *Xenomystax atrarius*, a member of the subfamily Muraenesocinae (larvae unknown). *Xenomystax* is closely related to *Paraxenomystax* (Smith 1979). Members of this subfamily are sometimes considered part of a separate family. Their larvae have a moderately elongate body with the gut length about 75% SL. Pigment occurs on the head, widely spaced along the throat to anus, and below the lateral midline in the form of a row of large spots. Size at transformation is probably similar to the congrid with most at 100 mm SL but some up to 200-300 mm SL.


Serrivomeridae (147-169 myomeres) *Serrivomer jespersenii* larvae are identified by a head shape that is sharp and slightly concave, pointed tail, simple gut with a length of ~75% SL, and a small nasal organ near the eye. The dorsal fin origin is located anterior to the anus. The last blood vessel occurs between myomeres 30-37. Pigment may be located variously over the lateral body surface, but ventral pigment is lacking. Other pigment appears along the dorsal and anal fin bases and in a cluster on the orbit above the eye. Size at transformation is 60 mm SL.

Eurypharyngidae (97-125 myomeres) *Eurypharynx* larvae are identified by their short, deep head and body, gut about 50-67% SL with one loop posteriorly, and pointed tail. Ventral pigment is limited to the gut loop. Size at transformation is 30-40 mm SL.

NETTASTOMATIDAE

Saurenehelys sp.

A




Two gut loops

CONGRIDAE

Paraxenomystax sp.



B



115 mm SL

Large spots below lateral line

C




SERRIVOMERIDAE

Serrivomer jespersenii

D

Sharply pointed head




36 mm SL

EURYPHARYNGIDAE

Eurypharynx pelecانoides

E

Short head



19 mm SL

One gut loop

Figure A, Smith 1984; B–C, Smith 1979; D, Bauchot 1959; E, Fahay 1983 (after Smith 1979).

MERISTICS

Vertebrae	Total: 181-193-201 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	X-X-X
Caudal fin	Reduced
Pelvic fin	Absent
Dorsal fin	R: 279-339-432 D100: 130-164-210 ^a
Pectoral fin	R: 14-16-18
Anal fin	R: 240-299-372 A100: 103-138-176 ^a
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Aleutian Is., 51-55°N
Ecology	Meso- and bathypelagic, 510-4580 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length
Length at flexion
Length at transformation
Sequence of fin development Dorsal and anal (late)

Pigment


- Small spots on top of spinal cord, at least posteriorly
- Row of spots dorsally along gut length, ventral row anterior to stomach
- Several groups (usually three) of internal spots along body subaxially—about four spots in each group^b
- Dorsal and anal fin bases

Diagnostic characters

- Distinguished from *Nemichthys* spp. by
- Body less elongate
 - Caudal structure (tail more round, not filiform)
 - Position of liver and last blood vessel, 30th and 70-88th myomere, respectively
 - Number of myomeres (181-201)

^aThe total dorsal and anal fin ray counts and the number of lateral line pores and vertebrae are often of no value; they are not comparable because caudal parts are often missing and regeneration may have occurred. Nielsen and Smith (1978) introduced artificial lengths/counts with a greater comparative value which we employ here (see their Materials and Methods section for a complete discussion), e.g., D200, A200 = number of fin rays anterior to vertebrae no. 201.

^bAccording to Smith (1979), there are three blotches of internal lateral pigment spots which become less prominent with growth.



Figures A–D, Smith 1979.

MERISTICS

Vertebrae	Total: 300-300-300 Precaudal: 77-85-105 Caudal: X-X-X
Branchiostegal rays	7-X-15
Caudal fin	Reduced
Pelvic fin	Absent
Dorsal fin	R: 330-330-330 D200: 170-207-253 ^a
Pectoral fin	R: 10-11-14
Anal fin	R: 320-320-320 A200: 186-218-273 ^a
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epi-, meso-, and bathypelagic, 91-1829 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length
Length at flexion
Length at transformation ~200 mm SL
Sequence of fin development Dorsal and anal (late)

Pigment


- Three prominent spots laterally below midline in small larvae (lower on body than on *Avocettina infans*)
- Above gut along length
- Above spinal cord

Diagnostic characters

- Elongate with filiform tail
 - Number of myomeres (>300)
- Distinguished from *A. infans* by
- Position of liver (myomere 40) and last vertical blood vessel (80-100th myomere)

Nemichthys larseni larvae are not known but adults occur within the study area. See Table 5 for data on adults. Additional meristic characters from Nielsen and Smith (1978) are: D200, 173-205-222; A200, 164-200-208. See *A. infans* (p. 38, footnote a).

^aSee *Avocettina infans*.



Figures A–F, Smith 1979.



Clupeiformes

The Clupeiformes (herrings, anchovies) are generally small, coastal marine fishes that occur worldwide. Most species form schools and swim near the surface in nearshore waters, feeding on plankton. They have specialized gill rakers for straining large amounts of water. The order consists of 4 families, 78 genera, and about 317 species (McGowan and Berry 1984). Clupeids generally lay demersal eggs whereas engraulidids have pelagic eggs that are sometimes ellipsoidal in shape. Larvae are elongate and similar in appearance but may be distinguished by myomere counts and pigmentation characters.

Families in study area: **Clupeidae**
Engraulididae

MERISTICS

Vertebrae	Total: 46-52-57 ^a Precaudal: 29-31-32 Caudal: 19-22-22
Branchiostegal rays	8-X-9
Caudal fin	9, 10+9, 8-9
Pelvic fin	Abdominal R: 9-9-9
Dorsal fin	R: 15-18-21
Pectoral fin	R: 17-17-17
Anal fin	R: 13-16-20
Gill rakers	U: 20-20-20 L: 45-45-45 ^a

LIFE HISTORY

Range	South of southern California to Arctic, not specific
Ecology	Nearshore shelf pelagic, 0-137 m; ^b 475 m ^c
ELH pattern	Oviparous; demersal, adhesive attached eggs; pelagic larvae
Spawning	Season: Jan-Apr (California), Mar-June (Alaska) ^d Area: Demersal (usually on vegetation), nearshore ^e Mode: Schools ^e Migration: Inshore ^e
Fecundity	Range/function: 6300-41,000/ $F=0.000000436 \times L^{4.71}$, $L=SL \text{ mm}^f$
Age at first maturity	2-4 yr (Puget Sound) ^f 2-6 yr (Bering Sea) ^g
Longevity	>19 yr ^h

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.3-1.7 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Segmented
Envelope	Smooth, clear, thick
Hatch size	5.6-7.5 mm SL (7 mm); yolk absorbed 9-10 mm SL
Incubation time/temp.	14-15 d/8.5°C
Pigment	

Diagnostic characters

- Wide perivitelline space

LARVAE

Preanal length	72% SL
Length at flexion	18 mm SL
Length at transformation	26-35 mm SL
Sequence of fin development	Caudal, dorsal, anal, pelvics, pectorals

Pigment

- Isthmus, thoracic region
- Gut: Dorsal, midventral (midventral melanophores on intestine usually paired, sometimes slightly offset)
- Caudal, hypural

Diagnostic characters (see Table 4)

- Distinguished from *Engraulis mordax* (p. 48) by
- More myomeres (usually >50)
 - Longer gut (consistently >70% SL)
 - Placement of dorsal fin relative to anus (6-7 myomeres between dorsal and anal fins)
 - Less isthmus pigment
 - Prominent swimbladder in *E. mordax*
 - Dorsal spots at notochord tip usually not in *E. mordax*
 - Generally more pigmented
- Distinguished from *Sardinops sagax* by
- Presence of posteroventral pigment on gut
 - Dorsal spots at notochord tip usually not in *S. sagax*

See also *Mallotus villosus* (p. 80) and *Ammodytes hexapterus* (p. 540)

^a McGowan and Berry (1984) report total vertebral counts of 53-60 and lower gill raker counts of 37-52.

^b Alaska Department of Fish and Game 1985

^c Allen and Smith 1988

^d Scattergood et al. 1959

^e Schaefer 1937

^f Katz 1942

^g Rudomilov 1972

^h Fitch and Lavenberg 1975

Ref: Garrison and Miller 1982, Grant 1986, McGowan and Berry 1984.




Figure A, Uchida et al. 1958; B-E (D, ventral view), NWAFC originals (B. Vinter).

MERISTICS^a

Vertebrae	Total: 48-X-54 Precaudal: 28-29-30 Caudal: 22-22-23
Branchiostegal rays	6-X-10
Caudal fin	7-9, 10+9, 6-8
Pelvic fin	Abdominal R: 8-8-8
Dorsal fin	R: 17-X-20
Pectoral fin	R: 17-17-17
Anal fin	R: 17-X-20
Gill rakers	U: 21-X-23 L:44-X-45

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59°N
Ecology	Nearshore shelf pelagic, 0-80 m ^b
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Winter-summer ^c Area: Near surface in coastal and offshore waters ^b Mode: Pelagic, schools Migration:
Fecundity	Range/function: 30,000-65,000 ^c
Age at first maturity	1-2 yr ^b
Longevity	10 yr ^b , possibly 25 yr ^d

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.34-2.05 mm
No. of oil globules	One
Oil globule diameter	0.16 mm
Yolk	Irregularly segmented
Envelope	Smooth, thin
Hatch size	3.5-3.75 mm SL
Incubation time/temp.	2.5 d/17°C; 2.5-4 d/13-16°C ^e
Pigment	

Diagnostic characters

- Wide perivitelline space

LARVAE

Preanal length	>75% SL
Length at flexion	9.0-14.0 mm SL
Length at transformation	35 mm SL
Sequence of fin development	Caudal, dorsal, anal, pelvics, pectorals
Pigment	<ul style="list-style-type: none"> • Row of melanophores along dorsal surface of gut and along ventral midline • Melanophores over anterior gut appear dash-like, and those over posterior gut become larger and more intense with development

Diagnostic characters (see Table 4)

- See *Clupea pallasii* (p. 44)
- Dash-like melanophores along anterior surface of gut
- Distinguished from *C. pallasii* by
 - Lack of posteroventral pigment on gut
- Distinguished from *Engraulis mordax* (p. 48) by
 - Dorsal and anal fin placement (6-8 myomeres between fins)

^aMeristics include information from McGowan and Berry 1984.

^bFrey 1971

^cHart 1973

^dFitch and Lavenberg 1971

^eGarrison and Miller 1982




Figure A, Matarese and Sandknop 1984; B–F, Kramer 1970. Stippling in these figures is not pigment.

MERISTICS

Vertebrae	Total: 43-46-47 Precaudal: 24-25-26 Caudal: 19-20-21
Branchiostegal rays	14-14-14
Caudal fin	8-10, 10+9, 8-10
Pelvic fin	Abdominal R: 6-6-6
Dorsal fin	R: 14-X-19
Pectoral fin	R: 13-X-20
Anal fin	R: 19-X-26
Gill rakers	U: 28-X-41 L: 37-X-45

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epi- and mesopelagic, 0-300 m ^a
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: May ^b to mid-Aug ^c ; year-round (California) ^d Area: Pelagic Mode: Migration:
Fecundity	Range/function: 4025 ^e -30,000 ^f
Age at first maturity	1-2 yr ^g
Longevity	7 yr ^h

^a Pacific Fisheries Management Council 1978^b Kendall and Clark 1982^c Blackburn 1973^d Lasker and Smith 1977^e Frey 1971^f Baxter 1967^g Clark and Phillips 1952^h Hart 1973

Ref: McGowan and Berry 1984, Wang 1981.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.23-1.55 mm × 0.6-0.8 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Segmented
Envelope	Smooth, transparent
Hatch size	2.5-3.0 mm SL; yolk absorbed 3.5-4.0 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

- Ovoid
- Narrow perivitelline space

LARVAE

Preanal length	60-66% increasing to 75% SL, decreases in postflexion larvae
----------------	--

Length at flexion**Length at transformation**

Sequence of fin development	Caudal, dorsal, anal, pelvics, pectorals
-----------------------------	--

Pigment

- Isthmus
- Row of melanophores along dorsal surface of gut continuing along ventral body midline
- Ventral pigment on gut beginning about midway

Diagnostic characters (see Table 4)

- See *Clupea pallasii* (p. 44)
- Isthmus pigment present
- Dorsal and anal fin placement (0-2 myomeres between fins)
- Midventral melanophores on intestine usually staggered, not single as in osmerids, and usually not as evenly paired as in *C. pallasii*





Figure A, Bolin 1936; B–F, Kramer and Ahlstrom 1968. Stippling in these figures is not pigment.



Salmoniformes

The salmoniform fishes (salmons, smelts, deep sea smelts, and others) are primarily freshwater spawners except for members of the Argentinioidei, the largest taxon of salmoniforms with marine eggs and larvae in our area. There is much disagreement as to the composition of the order (see Fink 1984). In our area, it consists of 4 suborders, 15 families, 90 genera, and about 320 species (J. Nelson 1984). The argentinoids are mostly deep-sea fishes. Eggs are pelagic and have distinctive pustules on the inner surface of the egg membranes; larvae have a variety of forms. Larval characters include presence/absence of eyestalks, unique development of median fins, and distinctive pigment patterns. Pigment patterns for bathylagids are discussed according to the categories described by Ahlstrom et al. (1984b)—species with large, isolated melanophores and those with a linear series of small melanophores.

A summary of meristic characters is included for Alepocephalidae and Platytroctidae. Although salmon parr are occasionally collected in plankton tows, they are not treated here. Information on their early life history is reviewed in Kendall and Behnke (1984).

Families in study area: **Argentinidae**
Bathylagidae
Opisthoproctidae
Alepocephalidae
Platytroctidae
Osmeridae
Salmonidae

MERISTICS

Vertebrae	Total: 47-X-51
	Precaudal: 30-31-32
	Caudal: 16-18-18
Branchiostegal rays	5-5-5
Caudal fin	12, 10+9, 11
Pelvic fin	Abdominal
	R: 10-11-12
Dorsal fin	R: 10-X-13
Pectoral fin	R: 11-X-18
Anal fin	R: 12-X-15
Gill rakers	U: 7-8-9 L:14-17-20

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Epi- and mesopelagic, 11-274 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Jan-spring ^a
	Area:
	Mode:
	Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	>5 yr ^a

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.31-1.66 mm
No. of oil globules	One, at vegetal pole
Oil globule diameter	0.27-0.46 mm
Yolk	Segmented
Envelope	Pustules (pronounced raised bumps on inner surface)
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters

- Pustules more pronounced than in bathylagid eggs

LARVAE

Preanal length	76-84% SL
Length at flexion	>9 mm SL
Length at transformation	25-30 mm SL (prolonged) ^b
Sequence of fin development	Caudal, dorsal and anal, pectorals, pelvics
Pigment	

- Internal head pigment
- Series of ventral trunk blotches extending from pectoral fin to end of gut
- 1-2 large blotches postanal with a large caudal blotch
- Blotches expand with development (see figure)


Diagnostic characters

- Transverse rugae lining gut
 - Fin rays form in finfold, away from body margin
- Distinguished from *Nansenia candida* by
- Presence of internal head pigment
 - Caudal pigment on preflexion larvae
 - Fewer spots over gut

^aFitch and Lavenberg 1968

^bTransformation: Morphological changes (deepening of body, lengthening of snout, enlargement of eye) and folding of anterior gut to form stomach, along with masking of larval pigment, occur at 25-30 mm. Pelagic juveniles may occur at 50-100 mm.

Ref: Ahlstrom et al. 1984b.



Figures A-E, Ahlstrom et al. 1984b.

MERISTICS

Vertebrae	Total: 44-X-47	
	Precaudal: X-X-X	
	Caudal: X-X-X	
Branchiostegal rays	3-3-3	
Caudal fin	11, 10+9, 14	
Pelvic fin	Abdominal	
	R: 9-10-10	
Dorsal fin	R: 9-X-10 ^b	
Pectoral fin	R: 9-X-11 ^b	
Anal fin	R: 8-X-9	
Gill rakers	U: 12-12-12	L: 18-18-18

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesopelagic, 200-1000 m ^c
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.39-1.56 mm
No. of oil globules	One, at vegetal pole
Oil globule diameter	0.41-0.49 mm
Yolk	
Envelope	Pustules
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters

- Pustules

LARVAE

Preanal length	74-82% SL
Length at flexion	
Length at transformation	~15 mm SL
Sequence of fin development	Caudal, dorsal, anal, pectorals, pelvics

Pigment

- Embedded pigment above gut
- Superficial head pigment (>8.4 mm SL)
- With development, embedded line of melanophores running length of body
- Conspicuous caudal pigment
- Ventral pigment from isthmus along anterior 2/3 of gut

Diagnostic characters


- Transverse rugae lining gut only in posterior section, anteriorly an elongate s-shaped fold is present along with longitudinal rugae
- Fin rays form in finfold, away from body margin

^a Referred to by Ahlstrom as northern *Nansenia* (E.H. Ahlstrom notes).

^b Ahlstrom et al. 1984b

^c Cohen 1958

Ref: Ahlstrom et al. 1984b.



Figures A, C, Ahlstrom et al. 1984b; B, NWAFC original (B. Vinter).

MERISTICS

Vertebrae	Total: 48-X-53 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	2-2-2
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 9-X-10 ^a
Dorsal fin	R: 10-10-11
Pectoral fin	R: 10-10-12
Anal fin	R: 18-X-22 ^a
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Cent. California, 34-38°N, to Oregon, 42-46°N
Ecology	Mesopelagic, 200-1000 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**


Preanal length 84-89% SL
Length at flexion
Length at transformation^b
Sequence of fin Caudal, dorsal, anal,
development pectorals, pelvics
Pigment - Linear series of small melanophores
• Series of melanophores (up to 18) along lateral gut
• With development to late postflexion, additional
pigment appears on lower jaw, isthmus, opercle,
pectoral fin base, and lateral caudal peduncle

Diagnostic characters

- Anterior section of gut smaller in diameter compared with other species (see illustration)
 - Eye stalks longer and persist into later larval stages (65% HL)
 - Fin rays form in finfold, away from body margin
- Distinguished from other bathylagids with eyestalks by
- Linear series of small melanophores
 - *B. pacificus* has large, isolated spots, long eyestalks, and less intense pigment
 - *B. ochotensis* has shorter stalks and more intense pigment

^aAhlstrom et al. 1984b

^bTransformation: Direct, marked in all species; slender body becoming deeper with development, large head and eyes, gut coils and becomes covered by black peritoneal sheath.



Figures A–B, Ahlstrom et al. 1984b (North Atlantic specimen).

MERISTICS

Vertebrae	Total: 50-X-55
	Precaudal: 18-X-21
	Caudal: 30-X-34
Branchiostegal rays	2-2-2
Caudal fin	16-18, 10+9, 15-17
Pelvic fin	Abdominal
	R: 6-X-8
Dorsal fin	R: 6-X-9
Pectoral fin	R: 11-X-16
Anal fin	R: 20-X-28
Gill rakers	U: X-X-X L: 25-X-27 ^a

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesopelagic, 60-1000 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	Segmented
Envelope	Pustules on inner surface
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	59-61% SL
Length at flexion	
Length at transformation	Larval characters visible to 50 mm SL
Sequence of fin development	Caudal, dorsal, anal, pectorals, pelvics
Pigment - Large isolated melanophores	
• Preflexion larvae: Pigment on lower jaw, midgut, and tail	
• Flexion larvae	
—Opposing dorsal and ventral midline melanophores	
—Large melanophores on head and pectoral fin base	
—Large lateral blotch at base of caudal fin	

Diagnostic characters

- Gut shorter (usually only 50% SL) than in other species
- Eye rounder and larger than in other bathylagids (not stalked)
- Fin rays develop in finfold, away from body margin
- Distinguished from other bathylagids without eyestalks by
- Large isolated spots
- *B. wesethi* has a series of melanophores along hypaxial region, large eye, and spots posteriorly along dorsal midline
- *Leuroglossus* spp. larvae have smaller eyes and no dorsal spots

^aE.H. Ahlstrom notes

Ref: Ahlstrom et al. 1984b.




Figure A, NWAFC original (B. Vinter); B–C, Ahlstrom et al. 1984b.

MERISTICS

Vertebrae	Total: 47-X-49 Precaudal: 25-26-27 Caudal: 21-22-23
Branchiostegal rays	2-2-2
Caudal fin	13-14, 10+9, 15-16
Pelvic fin	Abdominal R: 9-10-10
Dorsal fin	R: 9-11-12
Pectoral fin	R: 9-10-11
Anal fin	R: 12-X-15
Gill rakers	U: X-X-X L: 28-28-28 ^a

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi- and mesopelagic, 49-900 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Jan-Feb ^b Area: Off continental slope ^c Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	0.92-1.1 mm
No. of oil globules	>10, usually coalesce to 1 in late stage ^d
Oil globule diameter	
Yolk	Segmented
Envelope	Pustules on inner surface
Hatch size	
Incubation time/temp.	
Pigment	• Unpigmented

Diagnostic characters

- Lack of pigmentation in late-stage eggs

LARVAE

Preanal length	81-90% SL
Length at flexion	
Length at transformation	
Sequence of fin development	Caudal, dorsal, anal, pectorals, pelvics
Pigment	- Linear series of small melanophores
	• Series of melanophores develops on hypaxial myomeres
	• Epaxial melanophores limited to posterior body
	• Series along posterior gut
	• May occur on urostyle at sizes <7.9 mm SL

Diagnostic characters


- Eye stalks
- Distinguished from other bathylagids with linear series of smaller melanophores by
- Posterior gut melanophores larger and fewer
 - Anterior region of gut lacks pigment
 - Epaxial myomere series limited to posterior region
 - Fin rays form in finfold, away from body margin
- Distinguished from *B. bericoides* by
- Small *B. bericoides* larvae are unavailable but presumably eye stalks are longer at comparable stages
- Distinguished from *B. pacificus* by
- Series of melanophores as opposed to isolated spots

^aE.H. Ahlstrom notes

^bWang 1981

^cAhlstrom 1965

^dNumerous globules at vegetal pole which coalesce to one clump at each equatorial pole.



Figures A, C–D, Ahlstrom et al. 1984b; B, NWAFC original (B. Vinter).

MERISTICS

Vertebrae	Total: 44-X-49 Precaudal: 18-X-25 Caudal: 21-X-28
Branchiostegal rays	2-2-2
Caudal fin	13, 10+9, 13-14
Pelvic fin	Abdominal R: 7-X-10
Dorsal fin	R: 8-X-9
Pectoral fin	R: 7-X-11
Anal fin	R: 15-X-22
Gill rakers	U: X-X-X L: 28-X-32 ^a

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesopelagic, 149-1000 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Feb-Mar; ^b spring ^c Area: Mode: Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	Segmented
Envelope	Pustules
Hatch size	
Incubation time/temp.	
Pigment	


Diagnostic characters**LARVAE**

Preanal length	76-85% SL
Length at flexion	
Length at transformation	
Sequence of fin development	Caudal, dorsal, anal, pectorals, pelvics
Pigment - Large isolated melanophores	
• Early larvae have large lateral blotch at midbody and another posteriad, becoming located on trunk with development	
• Third blotch forms midway between and below other two blotches	
• Fourth lateral trunk blotch forms in some late larval specimens between pectoral fin and large midbody blotch	
• Other melanophores form lateral to liver and terminal section of gut	
• Also isthmus/thoracic, dorsal gut, caudal, and two bands on body (early)	

Diagnostic characters

- Stalked eyes
- Fin rays form in finfold, away from body margin
- Only bathylagid with stalked eyes and pigment consisting of large isolated spots

^aE.H. Ahlstrom notes^bWang 1981^cFitch and Lavenberg 1968



Figures A–B, NWAFC original (B. Vinter); C, Ahlstrom et al. 1984b.

MERISTICS

Vertebrae	Total: 43-X-46 Precaudal: 23-23-23 Caudal: 22-22-22
Branchiostegal rays	2-2-2
Caudal fin	14-15, 10+9, 14-15
Pelvic fin	Abdominal R: 9-X-11
Dorsal fin	R: 11-12-13
Pectoral fin	R: 9-10-11
Anal fin	R: 14-X-16
Gill rakers	U: 8-8-8 L: 16-X-17

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Epi- and mesopelagic, 40-1000 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	0.90-1.10 mm
No. of oil globules	12-20, subequal ^a
Oil globule diameter	
Yolk	
Envelope	
Hatch size	
Incubation time/temp.	
Pigment	<ul style="list-style-type: none"> No pigment over oil globules

Diagnostic characters**LARVAE^b**

Preanal length	79-94% SL
Length at flexion	>6 mm, by 11 mm SL
Length at transformation	~25 mm SL
Sequence of fin development	Caudal, dorsal, anal, pectorals, pelvics
Pigment	Linear series of small melanophores <ul style="list-style-type: none"> Initially a series of paired spots dorsolaterally to gut extending from pectoral fin base to terminal section, becoming embedded with development (6-8 pairs developing to 7-8 pairs) Notochord tip (dorsal and ventral)
At flexion	<ul style="list-style-type: none"> Series of melanophores develops along hypaxial region, and soon after a series develops along epaxial More lateral spots are added Median finfold Increased head pigment

Diagnostic characters

- Eyes not stalked
- Pigment pattern unique among *Bathylagus* spp. (including *Leuroglossus* spp.)
- Fin rays form in finfold, away from body margin
- Only bathylagid without stalked eyes and pigment consisting of a linear series of small melanophores

^aOil globules may only partially coalesce.

^bData on preflexion and postflexion larvae are from E.H. Ahlstrom notes. Illustrations are unavailable.




Figure A, Ahlstrom 1969; B–C, Ahlstrom et al. 1984b.

MERISTICS

Vertebrae	Total: 47-X-52 Precaudal: 26-X-29 Caudal: X-X-X
Branchiostegal rays	2-2-2
Caudal fin	15-17, 10+9, 15 ^b
Pelvic fin	Abdominal R: 8-X-9
Dorsal fin	R: 10-X-11
Pectoral fin	R: 8-X-9
Anal fin	R: 11-X-14
Gill rakers	U: 8-X-9 L: 17-X-19

LIFE HISTORY

Range	Brit. Col., 48°30'-55°N, to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathypelagic, 0-1800 m ^c
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Fall-winter; ^b summer ^d Area: Off continental slope ^b Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.65-1.90 mm
No. of oil globules	<5 up to 9, usually coalescing to 1 ^e
Oil globule diameter	0.35-0.40 mm, 0.47 after fusion ^e
Yolk	Segmented
Envelope	Pustules
Hatch size	
Incubation time/temp.	
Pigment	<ul style="list-style-type: none"> • Tip of notochord

Diagnostic characters

- Egg diameter

LARVAE

Preanal length	72-78% SL
Length at flexion	13-18 mm SL
Length at transformation	31-35 mm SL
Sequence of fin development	Caudal, dorsal and anal, pectorals, pelvics
Pigment - Large isolated melanophores	<ul style="list-style-type: none"> • Series of 5-6 spots on posterior gut • Midtrunk patch • Lower trunk blotches • 1-2 postanal lateral blotches in larger larvae

Diagnostic characters

- Eye stalks short and only in early larvae
 - Fin rays form in finfold, away from body margin
 - More pigmented than *L. stilbius*
- Distinguished from *Bathylagus milleri* by
- Eye stalks short
 - Lack of dorsal spots

^aPlaced in *Bathylagus* by Ahlstrom et al. 1984b.

^bDunn 1983

^cFedorov 1973

^dAhlstrom 1969

^eSimilar migrations, see *L. stilbius*.




Figure A, Ahlstrom 1969; B–F, Dunn 1983.

MERISTICS

Vertebrae	Total: 38-X-42 Precaudal: 20-21-22 Caudal: 19-20-21
Branchiostegal rays	2-2-2
Caudal fin	12-16, 10+9, 13-15
Pelvic fin	Abdominal R: 8-X-10 ^b
Dorsal fin	R: 9-X-11
Pectoral fin	R: 8-X-11
Anal fin	R: 11-X-14
Gill rakers	U: 7-X-9 L: 18-X-20

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Epi- and mesopelagic, 0-690 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Winter-spring ^c Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.10-1.21 mm
No. of oil globules^d	
Oil globule diameter	
Yolk	Finely segmented
Envelope	Pustules
Hatch size	3 mm SL
Incubation time/temp.	
Pigment	
	<ul style="list-style-type: none"> • Yolk membrane above oil globule • Notochord tip

Diagnostic characters

- Egg diameter: Smaller than in *L. schmidti*

LARVAE^e

Preanal length	74-80% SL
Length at flexion	
Length at transformation	25-29 mm SL
Sequence of fin development	Caudal, dorsal and anal, pectorals, pelvics
Pigment	Large isolated melanophores
	<ul style="list-style-type: none"> • Midtrunk patch between pectoral fin base and anus in smaller larvae is similar to <i>L. schmidti</i> • Series of 5-6 melanophores on posterior gut

Diagnostic characters

- Eye stalks short and only in early larvae
 - Fin rays form in finfold, away from body margin
- Distinguished from *L. schmidti* by
- Lack of lower trunk and postanal blotches

^aPlaced in *Bathylagus* by Ahlstrom et al. 1984b.

^bAhlstrom et al. 1984b

^cAhlstrom 1965

^dAfter fertilization, 15-25 oil globules at vegetal pole; 2-5 oil globules during early stage which coalesce to 2 (sometimes 1) of equal size at opposite poles. These oil globules migrate toward and coalesce under the embryo prior to hatching.

^eData on preflexion and postflexion larvae are from E.H. Ahlstrom notes. Illustrations are unavailable.




Figure A, Ahlstrom 1969; B-C, Ahlstrom et al. 1984b.

MERISTICS

Vertebrae	Total: 81-X-84 ^a Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	2-2-2
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 7-7-8 ^b
Dorsal fin	R: 13-14-16 ^b
Pectoral fin	R: 10-12-13 ^b
Anal fin	R: 10-12-14 ^b
Gill rakers	U: 16-X-20 L: 28-X-32

LIFE HISTORY

Range	N. California, 38-42°N, to Brit. Col., 48°30'-55°N
Ecology	Epi- and mesopelagic
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	(Both sexes appear to mature at 400 mm SL) ^b
Longevity	5 yr ^c

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	2.2-2.6 (ripe ovarian eggs)
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	80-82% SL
Length at flexion	
Length at transformation	Up to 124 mm SL ^d
Sequence of fin development	Caudal, pectorals and pelvics, dorsal and anal

Pigment

- Dorsal blotches (six pairs), which extend into finfold, and ventral (eight pairs) lateral blotches; blotches alternate except for postanal ones which form a band
- Large caudal blotch
- Head heavily pigmented
- Lower gill arches heavily pigmented
- Pectoral and pelvic fin bases

Diagnostic characters

- Gut elongate
 - Sac-like stomach (elongate and pointed at tip)
- Distinguished from other opisthoproctids by
- Elongate snout with unique triangular flap at tip
 - Round eyes (anterodorsal)
 - Distinctive heavy pigment pattern
 - Number of myomeres (81-84)

^aAhlstrom et al. 1984b^bStein and Bond 1985^cFitch and Lavenberg 1968^dTransformation (family): Marked by deepening of body and attainment of melanistic integument and large scales.




Figure A–B (B, ventral view), Ahlstrom et al. 1984b; C, Cohen 1960.

MERISTICS

Vertebrae	Total: 40-X-44 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	2-2-2
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 8-X-9
Dorsal fin	R: 10-X-11
Pectoral fin	R: 13-13-13
Anal fin	R: 8-X-9
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Epi- and mesopelagic, 152-457 m
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	5 yr ^b

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length	74-75% SL
Length at flexion	
Length at transformation	See <i>Bathylchnops exilis</i>
Sequence of fin development	Pectorals and pelvics probably right after caudal, before dorsal and anal

Pigment - Genus

- Lateral series of melanophores above gut; some species develop serial melanophores on hypaxial myomeres
- Head pigment: Jaws, internal snout, gill arches

Diagnostic characters

- Gut elongate (sac-like stomach, elongate and pointed at tip)
- Tubular eyes
- Elongate pectoral and pelvic fin rays
- Number of myomeres (40-44), less than *B. exilis* (81-84) and more than *Macropinna microstoma* (34-37)

^a According to A.E. Peden (Brit. Col. Prov. Mus., Victoria, B.C., Canada V8V 1X4, pers. commun., 22 Jan. 1987), several species may be in the area with *D. longipes* having the more southerly distribution. Other species may occur north of Oregon and at least one form occurs off British Columbia.

^b Fitch and Lavenberg 1968




Figure A, Ahlstrom et al. 1984b (after Roule and Angel 1930, Mediterranean specimen).

MERISTICS

Vertebrae	Total: 34-X-37 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	3-3-3
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 9-X-10
Dorsal fin	R: 11-X-12
Pectoral fin	R: 17-X-19
Anal fin	R: 14-14-14
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi- and mesopelagic, 99-891 m
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length 59-64% SL
Length at flexion
Length at transformation See *Bathylchnops exilis*
Sequence of fin development Caudal, pectorals and pelvics, dorsal and anal

Pigment

- Series of melanophores on each hypaxial myomere
- Heavy embedded blotch at pelvic fin base; expands dorsad and ventrad
- Caudal blotch
- Above terminal section on gut ventral to liver
- Lower jaw

Diagnostic characters

- Deeper body and shorter gut than *B. exilis* and *Dolichopteryx longipes*
- Head with pronounced hump or bend at nape
- Tubular eyes directed dorsally
- Number of myomeres (34-37)




Figure A, NWAFC original (B. Vinter); B–C, Chapman 1939.

Members of the slickhead family are found worldwide in the deep sea, with six species in six genera found within the limits of this study area. Adults occur at depths of 45 to 5500 m but are primarily taken in hauls near the ocean bottom below 600 m. Juveniles occur in midwater (Fitch and Lavenberg 1968). Reproductive characteristics are unknown. Little information on alepocephalid early-life-history stages has appeared since Beebe (1933) in which they were found to hatch from large eggs (3-4 mm) and have direct development. There is no close relationship between alepocephalids and argentinoids because alepocephalids have large eggs, direct development, and share no specialized ontogenetic characters with argentinoids (Ahlstrom et al. 1984b).

Table 6
Meristic characters of family Alepocephalidae.

Taxon	Distribution	Vertebrae		Fins				Gill rakers		Branchiostegals
		Precaudal	Caudal (Total)	Dorsal	Anal	Pectoral	Pelvic	Upper	Lower	
<i>Alepocephalus tenebrosus</i>	Cent. Calif.-Bering Sea		(53-55)	17	17-18	10	6-7	7-8	17-18	6-7
<i>Bathylaco nigricans</i>	SSC-Oregon		(45-46)	17-22	11-12	6-11	6-9	3-5	8-13	9-10
<i>Ericara salmoneum</i>	S. Calif.-Bering Sea			17-19	4-28	12	6			6
<i>Leptochilichthys agassizi</i>	SSC-Oregon			14	13	11	10	8	19	13
<i>Narctes stomias</i>	SSC-Wash.			17-21	14-16	10-11	8-9	3-4	12-14	8
<i>Talismania bifurcata</i> *	SSC-Brit. Col.	16-17	27-28 (43-46)	22	21-23	10-12	6-7	7-8	15-18	7

*The NWAFC meristic database was updated by Sazonov 1981.

Tubeshoulders are found in all oceans, and five species in four genera are known in the Northeast Pacific Ocean and Bering Sea (Matsui and Rosenblatt 1987). Adults are commonly taken in midwater trawls but have been found at the surface (at night) to below 1000 m (Fitch and Lavenberg 1968). Little is known of their reproduction and early life history except that juveniles of *Sagamichthys abei* migrate upward at night to within 200 m of the surface to feed (Hart 1973).

Table 7
Meristic characters of family Platytroctidae.*

Taxon	Distribution	Vertebrae		Fins				Gill rakers		Branchiostegals
		Precaudal (Total)	Caudal	Dorsal	Anal	Pectoral	Pelvic	Upper	Lower	
<i>Holtbyrnia innesi</i>	Bering Sea	26-30 (46-48)	18-20	18-19	17	16	9	7	15-16	8
<i>Holtbyrnia latifrons</i>	SSC-Brit. Col.	27-28 (46-50)	20	17-20	14-16	16-20	8-9	6-8	17-19	8-9
<i>Maulisia argipalla</i>	SSC-Bering Sea	25-27 (46-47)	19-22	17-20	15-17	18-19	7-8	7-8	16-18	8
<i>Pellisolus eubranchus</i>	Oregon	20-23 (42-44)	20-24	17-19	15-16	18-21	6-8	5-6	17-18	8-9
<i>Sagamichthys abei</i>	S. Calif.-Brit. Col.	30-31 (50-52)	19-21	16-18	14-16	14-18	9-10	7-8	16-18	6-8

*Taxonomy and meristic data from Matsui and Rosenblatt (1987). The northernmost record of *Mirorictus taningi* (reported also as *Normichthys campbelli*) is 35°N but it may occur further north.

Smelts are confined to the Northern Hemisphere, and seven species in six genera are found in the northeastern Pacific. Some species spawn intertidally, others are anadromous. Spawning is protracted and en masse. Spent fish return to deeper water except for *Thaleichthys pacificus*, which experiences high, though not complete, mortality after spawning (Garrison and Miller 1982). Osmerid eggs are generally 0.80-1.1 mm, strong to feebly adhesive, and have numerous oil globules. The adhesive membrane results from the rupturing of an outer "chorion" during spawning which turns out and onto the substrate (Hearne 1983, 1984). In general, larval characteristics include an elongate body shape, gut 75% of standard length, subterminal mouth, conspicuous choroid fissure, stalked pectorals, no dorsal melanophores, a single row of melanophores along the ventral midline of the gut, and a single row of melanophores on the ventral midline of the tail (Hearne 1984). All osmerid larvae possess a single midventral row of melanophores below the gut. *Spirinchus starksi* larvae have a greater number of ventral melanophores than *Spirinchus thaleichthys* or *Thaleichthys pacificus* (Hearne 1983). Myomere counts may be of additional use. Osmerids are the most abundant larvae in the nearshore waters (0-20 km) off Oregon; usually >50% of the larvae collected from January through June are osmerids (B. Mundy, NMFS Southwest Fish. Cent., Honolulu Lab., Honolulu, HI 96822-2396, pers. commun., 1 Oct. 1986.). Presently, it is possible to identify only one of these larvae to genus or species. A complete developmental series of *Mallotus villosus* is known. They are common in our ichthyoplankton collections off Oregon, Washington, and in the Gulf of Alaska.

Table 8
Meristic characters of family Osmeridae.

Taxon	Distribution	Vertebrae		Fins				Gill rakers		Branchiostegals
		Precaudal	Caudal (Total)	Dorsal	Anal	Pectoral	Pelvic	Upper (Total)	Lower	
<i>Allosmerus elongatus</i>	S. Calif.-Brit. Col.	40-44	23-27	9-11	14-17	12-14	8	10-13	23-28	6-7
<i>Hypomesus pretiosus</i>	S. Calif.-Bering Sea	42-44	22-24 (62-70)	8-11	12-17	14-17	8	10-13	21-25	7-8
<i>Mallotus villosus</i>	Wash.-Arctic		(62-73)	10-14	16-23	16-21	9	8-13	24-35	7-8
<i>Osmerus mordax</i>	Brit. Col.-Arctic		(58-68)	8-11	12-16	11-14	8	8-11 (26-37)	18-24	6-8
<i>Spirinchus starksi</i>	S. Calif.-SE Alaska	33-36	25-29	8-11	15-21	10-11	8	8-13	24-31	7-8
<i>Spirinchus thaleichthys</i>	Cent. Calif.-Bering Sea	29-31	24-27 (54-61)	8-10	15-22	10-12	8	10-13	26-34	7-8
<i>Thaleichthys pacificus</i>	Cent. Calif.-Bering Sea		(65-72)	10-13	18-23	10-12	8	4-6	13-18	6-8

MERISTICS

Vertebrae	Total: 62-X-73 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-X-8
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 9-X-9
Dorsal fin	R: 10-X-14
Pectoral fin	R: 16-X-21
Anal fin	R: 16-X-23
Gill rakers	U: 8-X-13 L: 24-X-35

LIFE HISTORY

Range	Washington, 46-48°30'N, to Arctic, not specific
Ecology	Nearshore shelf pelagic, 0-200 m; ^a 750 m ^b
ELH pattern	Oviparous, demersal attached eggs, pelagic larvae
Spawning	Season: Fall (British Columbia); ^c spring (Bering Sea) ^d Area: Gravel beaches ^c Mode: Schools ^c Migration:
Fecundity	Range/function: 3000-6600 (British Columbia) ^c
Age at first maturity	2+ yr ^e
Longevity	>3 yr ^f

^aAndriashev 1954^bAllen and Smith 1988^cHart and McHugh 1944^dBaxter 1975^eTrumble 1973^fHart 1973

Ref: Templeman 1948.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	Adhesive
Hatch size	3-6 mm
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	~75% SL
Length at flexion	After yolk absorption, ~7-8 mm SL
Length at transformation	
Sequence of fin development	Probably caudal, dorsal, anal, pelvics, pectorals

Pigment

- Single row of melanophores along ventral gut midline
- Single row of postanal ventral melanophores developing into double row (ventral view)


Diagnostic characters (see Table 4)

Distinguished from clupeiforms by


- Higher number of myomeres (62-73)
- Pigment: Single rather than double row of mid-ventral melanophores below gut
- Greater gut length (75% SL)
- Presence of adipose fin

Distinguished from other osmerids by

- Combination of greater number of pectoral fin rays (16-21) and anal fin rays (16-23)



Figures A–E, Fahay 1983 (A, after Bigelow and Schroeder 1953; B–E, after Templeman 1948; Atlantic specimens).



Stomiiformes

Fishes of the order Stomiiformes are mostly tropical to temperate, with many being deep-sea. This highly diverse order contains 9 families, 53 genera, and about 248 species (J. Nelson 1984). A total of 6 families, 12 genera, and 22 species are found within the study area. Body forms among stomiiforms range from eel-like (idiacanthids) to deep and extremely compressed (sternoptychids). Most species have luminescent organs, teeth on both premaxilla and maxilla, mouth extending past the eye, an adipose fin, and a dark-brown or black color (primarily silver in *Gonostomatidae* and *Sternoptychidae*). Presently, there is no consensus on the relationships within the order (Weitzman 1974, Fink and Weitzman 1982, Ahlstrom et al. 1984c, Fink 1984). Herein, we treat the gonostomatids and sternoptychids together, and the chauliodontids, melanostomiids, malacosteids, and idiacanthids together. For convenience, we will call the former families the gonostomatoids, and the latter the stomioids, without phylogenetic implications. The identification of gonostomatoid larvae requires a knowledge of developmental data from larvae, juveniles, and adults (including fin rays, teeth, and photophores). Of importance are pigment patterns, position of dorsal and anal fins (caution must be used, since positions change with growth), presence of an adipose fin, and photophores (number, pattern, and sequence of development) (Ahlstrom et al. 1984c). In addition to pigment patterns and meristic and photophore characters, gut structure (trailing or not) is an important feature to consider in the identification of stomioids.

Families in study area: **Gonostomatidae**
Sternoptychidae
Chauliodontidae
Melanostomiidae
Malacosteidae
Idiacanthidae

STOMIIFORMES

Table 9

Definitions of alphabetical symbols used for designating photophores in deep-bodied sternoptychids and other stomiiform fishes (Ahlstrom et al. 1984c, in part).

Code	Definitions
Gonostomatoids	
SO	Symphyseal photophores (organs) located at tip of lower jaw.
Orb	Photophores associated with the eye located anterior and posterior of orbit.
Op	Photophores on opercle series generally three, coded as follows 1/(1+1).
Br (BRP)	Photophores located on the branchiostegal membranes.
Is (I)	Photophores located on the isthmus.
IP	Photophores of the ventral series found from the isthmus to the base of the pectoral fin.
PV	Photophores of the ventral series found from the pectoral fin base to the pelvic (ventral) fin base.
VAV	Photophores of the ventral series found from the pelvic (ventral) fin base to the anal fin base.
AC	Photophores of the ventral series found from the anal fin base to caudal fin base.
IC	Summary of photophores of the ventral series from the isthmus to caudal fin base (IP+PV+VAV+AC).
IV	Summary of photophores of the ventral series from isthmus to pelvic (ventral) fin base (IP+PV).
OV	Photophores of the lateral series from the opercle to pelvic (ventral) fin base.
VA (VALA)	Photophores of the lateral series from the pelvic (ventral) fin base to the anal fin base.
OAA	Summary of photophores of OV plus VA series.
OA (OAB)	Summary of lateral photophores from the opercle to anal fin base (OV+VA).
OAC (OC)	Entire lateral series on body sides just dorsal to ventral series and extending from opercular border, or just medial to it, over anal fin to caudal fin base.
ODM	Photophores (organs) found dorsal to the lateral midline (found only in <i>Gonostoma gracile</i>).
Deep-bodied Sternoptychids	
SO	Subopercle photophore which is equivalent to posteriormost photophore in opercular series of gonostomatoids.
PO	Photophore located anterior to orbit.
PTO	Photophore located posterior to orbit and may be equivalent to upper photophore of opercular series of gonostomatoids.
PRO	Preopercular photophore, used for PO photophore dorsal to ventral limb or preopercle.
Br	Same as gonostomatoid definition.
Is	Same as gonostomatoid definition.
AB	Photophores of ventral series located abdominally between pectoral fin base and pelvic fin base and equivalent to PV in gonostomatoids, plus a few posterior photophores of the IP series.
PAN	Photophores found anterior to anal fin and may be equivalent to VAV or VA in gonostomatoids.
AN	Photophores found above anal fin.
SC	Photophores found on lower (sub) caudal peduncle. Together with AN group may be equivalent to AC in gonostomatoids.
SAB	Photophores located above (supra) to the abdominal series and may be equivalent to VA in gonostomatoids.
SP	Photophores located above (supra) the pectoral fin and may be equivalent to OV in gonostomatoids.
SAN	Photophores located above (supra) to anal photophores and equivalent to part of AC series.


Table 10

Photophore distribution in selected gonostomatoid genera (Ahlstrom et al. 1984c, in part). See Table 9 for photophore definitions.

	No. of rows	SO	ORB	OP	BR	IS	IV	VAV	AC	Photophores in groups of glands?
<i>Argyropelecus</i>	2	No	2	2	6	Yes	18	4	10	Yes
<i>Cyclothone</i>	2	No	1	2	8-11	No	12-14	4-5	12-16	No
<i>Danaphos</i>	2	No	1	2-3	6	Yes	18	5	22-26	Yes
<i>Gonostoma</i>	2	Yes	1	2-3	9	No	11-16	3-10	15-23	No
<i>Sternoptyx</i>	2	No	2	2	3	Yes	15	3	7	Yes

MAJOR PHOTOPHORE GROUPS

A Gonostomatoid



B Sternoptychid




Figure A, Fahay 1983 (after Ozawa 1976); B, Badcock and Baird 1980 (redrawn).

Table 11
Fin position and condition of adipose fin in selected gonostomatoid fishes (Ahlstrom et al. 1984c, in part).

Genera	Adult	Larvae	Adipose fin
<i>Argyropelecus</i>	Anal origin opposite last dorsal fin ray	Anal origin behind dorsal fin	Present or absent
<i>Cyclothone</i>	Anal origin opposite dorsal fin or slightly behind	Same as adult	Absent
<i>Danaphos</i>	Anal origin behind dorsal fin	Same as adult	Absent
<i>Gonostoma</i>	Anal origin opposite or 3-4 rays in advance of dorsal origin	Same as adult	Present or absent
<i>Sternoptyx</i>	Anal origin opposite dorsal origin	Anal origin behind dorsal fin	Present

Table 12
Meristic characters of family Gonostomatidae.

Taxon	Distribution	Vertebrae		Fins					Gill rakers		
		Precaudal (Total)	Caudal	Dorsal	Anal	Pectoral	Pelvic	Caudal	Upper	Lower	Branchiostegals
<i>Cyclothone acclinidens</i>	SSC-Oregon	13-14	17-19	13-15	18-20	9-10	6	6,10+9,6 ^a	6-9	14-17	13-15
<i>Cyclothone atraria</i>	SSC-Bering Sea ^b	12-13	18-20	13-15	18-21	9-10	6		6-8	13-16	12-14
<i>Cyclothone pallida</i>	SSC-Brit. Col.	13-14	17-21	13-15	17-19	9-11	6-7		7-10	14-18	13-15
<i>Cyclothone pseudopallida</i>	SSC-Bering Sea	12-13 ^a	18-20 ^a	13-15	18-21	9-10	6		4-6	12-14	13-15
<i>Cyclothone signata</i>	SSC-Gulf of Alaska	13	17-19	13-14	18-20	9-10	6	6-8,10+9,6-7 ^a	3-5	10-11	12-14
<i>Gonostoma atlanticum</i>	SSC-Oregon	(38-39)		16-18	27-30	9-10	6-7		6-7	11	11-12
<i>Gonostoma gracile</i>	Gulf of Alaska-Bering Sea	18-19	22-24	9-12	22-30	8-10	6-8				11-12

^aE.H. Ahlstrom notes

^bRecords from north of British Columbia are questionable (A.E. Peden, Brit. Col. Prov. Mus., Victoria, B.C., Canada V8V 1X4, pers. commun., 22 Jan. 1987).

This worldwide genus of lightfishes may constitute the most abundant genus of fish in the sea. These small (to 80 mm) bathypelagic fishes are found most commonly at depths of 300 m and greater and have been found as deep as 5300 m. Very little information on life history is available. The eggs are unknown but larvae are common in offshore tropical and temperate areas. *Cyclothone* spp. larvae are discussed at the generic level since only partial series are available for the five species in the study area. Gorbunova (1982a) described the larvae of eight species from the Pacific including four from the study area (all species except *C. atraria*). The descriptions of *C. acclinidens*, *C. pallida*, and *C. pseudopallida* by Gorbunova (1982a) have been questioned by Ozawa and Oda (1986) who described the larvae (>7 mm SL) of five species of *Cyclothone* (all species from the study area except *C. signata*). Since preflexion larvae and transforming larvae linking larval series to adults are generally unavailable, Ozawa and Oda (1986) discussed the need for further investigation.

Preanal length for *Cyclothone* spp. larvae is about 50% SL. Lengths at flexion and transformation are 4-5 mm SL and about 14 mm SL, respectively. Caudal and dorsal fins form first, followed in sequence by the anal, pelvic, and pectoral fins. Photophores are described in Table 10.

Pigment characters for *Cyclothone* spp., based on described species, include

- 2-3 spots along gut with posteriormost spot at anus (sometimes more spots in larger specimens)
- One spot at cleithral symphysis
- Pigment on swimbladder
- Lateral series posterior to pectoral fins, ending prior to swimbladder
- Series over anal fin base (varies from ~5-12 spots and may or may not be evenly spaced; number and spacing of series may be species-specific)
- Some species develop dorsal series similar to anal base series
- Spots at base of anal fin pterygiophores
- Spots usually appear above and below urostyle and in hypural area

Diagnostic characters for *Cyclothone* spp. larvae include

- Distinct dark streak or intense melanophore over and parallel to parhypural on caudal fin base
- Pigmentation over gut and along ventral margin of tail
- Anal fin origin opposite dorsal fin or slightly behind (no ontogenetic movement)
- Conspicuous swimbladder
- No adipose fin
- Larvae generally elongate

Diagnostic characters for species

- *C. acclinidens* (melanophore on dorsal part of tail end)
- *C. pseudopallida* (predorsal melanophores 2-4, see figure)
- *C. atraria* (predorsal melanophores 6 or 7, see figure)
- *C. pallida* (predorsal melanophores 9-12, see figure)
- *C. signata* (no melanophore on dorsal part of tail end, and no predorsal melanophores)


Distinguish from *Gonostoma*

- See Tables 10-12
- Fewer anal fin rays (16-21 vs. 21-31)
- Fewer vertebrae (29-33 vs. 37-40)
- SO photophore absent
- Caudal pigment (present in *Gonostoma atlanticum* but not in *G. gracile*)





Larger postflexion larvae, which differ greatly from adults, undergo a metamorphic stage during which most of the photophores become pigmented simultaneously, the anterior portion of the body shortens, and the anus changes position from near the anal fin to nearer the ventral fin bases.

CYCLOTHONE

C. acclinidens



C. pseudopallida



Figures A-F, Ozawa and Oda 1986; G, Ahlstrom et al. 1984c.

MERISTICS *G. atlanticum* Norman 1930

Vertebrae	Total: 38-X-39 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	11-X-12
Pelvic fin	Abdominal R: 6-7-7
Dorsal fin	R: 16-18-18
Pectoral fin	R: 9-X-10
Anal fin	R: 27-28-30
Gill rakers	U: 6-X-7 L: 11-X-11

MERISTICS *G. gracile* Günther 1878

Vertebrae	Total: 40-X-42 Precaudal: 18-X-19 Caudal: 22-X-24
Branchiostegal rays	11-X-12
Pelvic fin	Abdominal R: 6-X-8
Dorsal fin	R: 9-X-14
Pectoral fin	R: 8-X-11
Anal fin	R: 22-X-30
Gill rakers	U: 7-8-8 L: 12-13-14

LIFE HISTORY

Range	
(<i>G. atlanticum</i>)	South of southern California to Oregon, 42-46°N
(<i>G. gracile</i>)	Gulf of Alaska, 54-60°N to Bering Sea, 54-66°N
Ecology	Meso- and bathypelagic ^a
ELH pattern	Oviparous, pelagic larvae
Spawning^b	Season: Sept-Apr (Japan)
Age at first maturity^b	1 yr (males - then undergo sex reversal at 70-80 mm SL)
Longevity^b	2 yr (females) 1 yr (males)

^a*G. atlanticum* mesopelagic only.^b*G. gracile* only.

Ref: Ahlstrom 1974, Ahlstrom et al. 1984c, Kawaguchi and Marumo 1967, Ozawa 1986b, Sanzo 1931a.

EARLY LIFE HISTORY DESCRIPTION**EGGS (*G. denudatum*)**

Diameter	0.80-0.81 mm
No. of oil globules	One
Oil globule diameter	0.20-0.21 mm
Yolk	
Envelope	
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE (*G. gracile*)**

Preanal length	
Length at flexion	
Length at transformation	
Sequence of fin development	Caudal, dorsal and anal, pectorals and pelvics
Pigment	
• Internal head pigment	
• Swimbladder and peritoneum	


Diagnostic characters

- G. gracile* distinguished from other *Gonostoma* spp. by
- Anal fin origin extremely in advance of dorsal fin origin
- G. gracile* distinguished from *G. atlanticum* by
- Lack of pigment on caudal peduncle and body surface
- G. atlanticum* distinguished from *Cyclothone* spp. by
- Caudal and ventral pigment
 - Swimbladder position

Photophores (see Table 10)

In *G. gracile* and *G. atlanticum*, photophores develop as a group. This differs from other species of *Gonostoma* where photophores develop gradually.

GONOSTOMA
Gonostoma atlanticum



Gonostoma gracile




Figure A, Ahlstrom et al. 1984c (after Ahlstrom 1974); B–E, Kawaguchi and Marumo 1967 (western Pacific specimens).

STERNOPTYCHIDAE

Found in all temperate and tropical seas, hatchetfishes are represented in the northeastern Pacific by seven species in three genera. Adults are found 100-5000 m below the surface (Schultz 1961). Sternoptychid eggs and larvae are planktonic, constituting the third most abundant family in the ichthyoplankton of open ocean waters (after myctophids and gonostomatids) (Ahlstrom 1972, Loeb 1979). *Argyrolepecus hemigymnus* has eggs ranging from 0.92 to 1.04 mm. The yolk is coarsely segmented and contains one oil globule (Ahlstrom 1974). Development of sternoptychids is distinctly divided into two periods (Sanzo 1935): Larval period, and the period of metamorphosis marked by a sharp reduction in body length, change in shape, and photophore and fin-ray development. Distribution and patterns of photophores, especially their sequence of development, are instrumental in larval identification. The following characters may aid in distinguishing between the three sternoptychid genera in the study area (see Tables 13-14).

	<i>Argyrolepecus</i>	<i>Danaphos</i>	<i>Sternoptyx</i>
Anal fin rays	11-13	24-25	14-16
Pectoral fin rays	10-11	13-14	10-11
Vertebrae	34-40	38	28-31
Adipose fin	Present/absent	Absent	Absent
Photophores			
Br	6	6	3
Orb	2	1	2
IV	18	18	15
VAV	4	5	3
AC	10	22-26	7

Table 13
Meristic characters of family Sternoptychidae.

Taxon	Distribution	Vertebrae		Fins					Gill rakers		Branchiostegals
		Precaudal (Total)	Caudal	Dorsal	Anal	Pectoral	Pelvic	Caudal	Upper (Total)	Lower	
<i>Argyropelecus affinis</i>	SSC-Oregon	(39-41)		9	12-13	10-11	6	10-12,10+9,4	(18-22)		10
<i>Argyropelecus hemigymnus</i>	SSC-Wash.	(37-39)		8	11	10-11	6	10,10+9,5	(18-24)		10
<i>Argyropelecus lychnus</i>	SSC-Brit. Col.	(35-37)		9	12	10-11	6	9-11,10+9,6	8-9	12-14	10
<i>Argyropelecus sladeni</i>	SSC-Oregon	(36-38)		9	12	10-11	6	10,10+9,6	(17-21)		10
<i>Danaphos oculatus</i>	SSC-Brit. Col.	(38)		6	24-25	10-11		8,10+9,3	2	11-13	10
<i>Sternoptyx diaphana</i>	SSC-Oregon	(28-30)		9-11	13-16	10-11			6-9		6
<i>Sternoptyx pseudobscura</i>	SSC-Brit. Col.	(27-30)		9-11	13-15	10-11			7-9		6

Table 14
Characters useful in separating larvae of *Argyropelecus* found in the Northeast Pacific (Belyanina 1984, in part).

Character	<i>A. affinis</i>	<i>A. hemigymnus</i>	<i>A. lychnus</i>	<i>A. sladeni</i>
Photophores: SAB, PAN, AN, SC in a continuous row*	Yes	No	No	No
Caudal pigment	At or before transformation (~8 mm SL)	At or before transformation (~8 mm SL)	No	>13 mm SL
Number of postabdominal spines	2	1 (posterior one reduced)	2	2
Number of dorsal fin rays	9	8	9	9
Eye shape (telescopic)	Round	Cylindrical	Round	Round
Presence of well-developed upper preopercular spine	No	No	Yes (pointed downward)	Yes (pointed upward)
Presence of frontal crest	No	No	Yes	Yes

*See Table 9 for definitions of symbols.

MERISTICS

Vertebrae	Total: 39-X-41 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	10-10-10
Caudal fin	10-12, 10+9, 4
Pelvic fin	Abdominal R: 6-6-6
Dorsal fin	R: 9-9-9
Pectoral fin	R: 10-11-11
Anal fin	R: 12-X-13
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Epi- and mesopelagic, 100-610 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**


Preanal length <50% SL increasing with development to ~50% SL

Length at flexion
Length at transformation <8.1 mm SL
Sequence of fin development Caudal and pectorals, dorsal and anal, pelvics
Pigment

- Initially, lightly pigmented on anterior body, increases posteriorly with development
- Characteristic spot occurs on caudal peduncle above subcaudal photophores; in larvae >11 mm SL, spot increases to a patch

Diagnostic characters

- Photophores: *A. affinis* distinguished by the location of supra-abdominal (SAB), preanal (PAN), anal (AN), and subcaudal (SC) photophores in a continuous straight line; AN and SC groups are separated by a gap
- Eyes telescopic
- Spines
 - At 8 mm SL, a spine anterior to spinous dorsal plate
 - At 11 mm SL, long posteriormost spine on spinous dorsal plate
 - At >11 mm SL, enlarged postabdominal spines



Figures A–D, Belyanina 1984.

MERISTICS

Vertebrae	Total: 37-X-39 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	10-10-10
Caudal fin	10, 10+9, 5
Pelvic fin	Abdominal R: 6-6-6
Dorsal fin	R: 8-8-8
Pectoral fin	R: 10-X-11
Anal fin	R: 11-11-11
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Washington, 46-48°30'N
Ecology	Epi- and mesopelagic, 100-731 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	0.92-1.04 mm
No. of oil globules	One
Oil globule diameter	0.26-0.28 mm
Yolk	Segmented
Envelope	
Hatch size	2.5 mm SL; yolk absorbed 4.7 mm SL
Incubation time/temp.	2-2.5 d (temperature unknown)

Pigment

- Embryo unpigmented

Diagnostic characters**LARVAE**

Preanal length	<50% SL increasing with development to ~50% SL
-----------------------	--

Length at flexion

Length at transformation	>6 mm SL; body shrinks 2-3 mm, gut shortens, head deepens, and eyes become telescopic
---------------------------------	---

Sequence of fin development	Caudal and pectorals, dorsal and anal, pelvics
------------------------------------	--

Pigment

- Unpigmented at hatching (except eyes) and through transformation
- After transformation, pigment appears on head in patches and on caudal peduncle

Diagnostic characters

- Elongate compressed body
- Photophores and sequence of development (~6-10 mm SL): Lower OP, BR, posterior IV, anterior IV, posterior AC, anterior AC, OA, ORB, and VAV
- SAB, PAN, AN, and SC photophores not in continuous row
- Postabdominal spines: In larvae >10-12 mm SL, anterior spine becomes thinner and elongated and bends under the posterior spine; posterior spine gradually reduces and becomes spur-shaped
- Number of dorsal fin rays (eight); other sternoptychids usually have nine
- Eye shape more cylindrical than other *Argyrolepecus* spp.




Figure A, Sanzo 1931a; B, Jespersen and Taning 1926; C, E, Belyanina 1984; D, Ahlstrom et al. 1984c (after Sanzo 1931a; A–B, D, Mediterranean specimens).

MERISTICS

Vertebrae	Total: 35-X-37 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	10-10-10
Caudal fin	9-11, 10+9, 6
Pelvic fin	Abdominal R: 6-6-6
Dorsal fin	R: 9-9-9
Pectoral fin	R: 10-X-11
Anal fin	R: 12-12-12
Gill rakers	U: 8-X-9 L: 12-X-14

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epi-, meso-, and bathypelagic, 198-396 m; 4066 m ^a
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**


Preanal length 50-60% SL
Length at flexion
Length at transformation
Sequence of fin Caudal and pectorals, dorsal
development and anal, pelvics
Pigment
• Lightly pigmented, restricted to preanal body

Diagnostic characters

- Upper preopercular spine bent downward whereas same spine in *A. sladeni* is pointed slightly upward
- Pigmentation: Specimens described to 15 mm SL appear to lack pigment on postanal body, specifically in the peduncle area
- Presence of frontal crest

^aHart 1973

Ref: Ahlstrom et al. 1984c, Belyanina 1984.



Figures A–C, Belyanina 1984.

MERISTICS

Vertebrae	Total: 36-X-38
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	10-10-10
Caudal fin	10, 10+9, 6
Pelvic fin	Abdominal
	R: 6-6-6
Dorsal fin	R: 9-9-9
Pectoral fin	R: 10-X-11
Anal fin	R: 12-12-12
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Epi- and mesopelagic, 101-610 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**


Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length 50-60% SL
Length at flexion
Length at transformation
Sequence of fin development Caudal and pectorals, dorsal and anal, pelvics
Pigment
• Initially, appears anteriorly on body in patches and, with development, increases posteriorly
• Spot on peduncle does not appear as early as *A. affinis*, first appearing as a patch in larvae >13 mm SL

Diagnostic characters

- Spines: Upper preopercular spine is developed (pointed slightly upward) and short frontal crests are present
- Distinguished from *A. affinis* and *A. hemigymnus* by
- Combination of nine dorsal spines and SAB, PAN, AN, and SC photophores not in continuous line



Figures A–C, Belyanina 1984.

MERISTICS *S. diaphana* Hermann 1781

Vertebrae	Total: 28-29-30 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	6-6-6
Pelvic fin	Abdominal R: X-X-X
Dorsal fin	R: 9-10-11
Pectoral fin	R: 10-X-11
Anal fin	R: 13-14-16
Gill rakers	U: X-X-X L: X-X-X

MERISTICS *S. pseudobscura* Baird 1971

Vertebrae	Total: 27-29-30 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	6-6-6
Pelvic fin	Abdominal R: X-X-X
Dorsal fin	R: 9-10-11
Pectoral fin	R: 10-X-11
Anal fin	R: 13-14-15
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N ^a
Ecology	Meso- and bathypelagic
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

^a*S. diaphana* only to Oregon, 42-46°N.^bAlthough not well documented, geographic variation among and within species is clearly present.^cA dramatic change in morphology occurs at transformation. Elongate larvae become deep-bodied juveniles. Body length cannot be associated with development.^dBadcock and Baird (1980) cite a personal communication from E.H. Ahlstrom stating that neither of the two forms of *Sternoptyx* spp. from the study area have caudal pigment. A figure of *S. pseudodiaphana* is provided as an example of an early larva with caudal pigment.

Ref: Badcock and Baird 1980, Belyanina 1983.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE** - Genus^b

Preanal length
Length at flexion
Length at transformation 6-14 mm SL^c
Sequence of fin development Pectorals and caudal, anal, dorsal and pelvics
Pigment
• Head and tip of lower jaw
• Peritoneum, becoming embedded with development
• Presence/absence of pigment on caudal peduncle; according to Belyanina (1983) some specimens of *S. diaphana* have pigment whereas specimens of *S. pseudobscura* have none^d






Diagnostic characters

- Prior to metamorphosis, larvae of the two species are indistinguishable; both with spiny ridges on frontal and parietal bones, opercular and post-temporal spines
- Distinguish *S. pseudobscura* juveniles from *S. diaphana* by
 - Rounder body
 - Posterior anal pterygiophores longer
 - SAN higher in position



Photophores (see Table 10)

Order of development in *Sternoptyx* spp. larvae: SO, Br2, AB7, I4, SP3, PTO, AN1 followed by PAN, SC, PRO, and SAN in early juveniles; finally SC.

STERNOPTYX



S. diaphana



S. pseudobscura





Figure A, Ahlstrom et al. 1984c; B, Badcock and Baird 1980 (eastern North Atlantic specimen); C–I, Belyanina 1984.

MERISTICS

Vertebrae	Total: 38-38-38 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	10-10-10
Caudal fin	8, 10+9, 3
Pelvic fin	Abdominal R: X-X-X
Dorsal fin	R: 6-6-6
Pectoral fin	R: 10-X-14
Anal fin	R: 24-X-25
Gill rakers	U: 2-2-2 L: 11-X-13

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Mesopelagic, 183-914 m
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

- Diameter
- No. of oil globules
- Oil globule diameter
- Yolk
- Envelope
- Hatch size
- Incubation time/temp.
- Pigment

Diagnostic characters

LARVAE

- Preanal length** <50% SL (short gut)
- Length at flexion**
- Length at transformation** ~45-50 mm SL
- Sequence of fin development**

Pigment

- Concentrated on preanal body in area below eye and over gut; smaller larvae generally unpigmented except over peritoneum

Diagnostic characters

- Larvae elongate
 - Photophores begin formation ~16.5 mm SL (see Table 15)
 - Photophores in clusters with common bases
 - Metamorphosis gradual with the initial formation of photophores in the BR and PV groups
- Distinguished from *Argyropelecus* spp. and *Sternoptyx* spp. by
- Long anal fin (24-25 fin rays) and short dorsal fin (6 fin rays)

Ref: Ahlstrom 1974, Ahlstrom et al. 1984c.

Table 15
Development of photophores in *Danaphos oculatus* (Ahlstrom et al. 1984c). * See Table 9 for definitions of photophore groups.

Size (mm SL)	Photophore groups								
	ORB	OP	SO	BR	IP	PV	VAV	AC	OA
Adult	1	3	0	(6)	(3)+(4)	(11)	(5)	(3)+16+(4)+1	6
16.5	0	0	0	(2)	0	0	0	0	0
16.5	0	0	0	(3)	0	(3)	0	0	0
19.2	0	0	0	(4)	0	(10)	0	0	0
21.0	1	1	0	(5)	(2)+(4)	(10/11)	0	(2)+0+0+0	0
21.3	1	1	0	(4/5)	(3)+(4)	(10)	0	(3)+0+(2)+0	0
21.8	1	2	0	(5)	(3)+(4)	(11)	(2)	(3)+8+(4)+0	2
24.2	1	2	0	(6)	(3)+(4)	(11)	(2)	(3)+9+(4)+0	2

*Parenthetical numbers indicate photophores found in common glands; nonparenthetical numbers indicate photophores are single.




Figure A, NWAFC original (B. Vinter).

Table 16

Meristic characters of stomioid genera. Most frequent count or range is followed by overall range or infrequent count in parentheses (Kawaguchi and Moser 1984, in part).

Taxon	Vertebrae	Fin rays			
		Dorsal	Anal	Pectoral	Pelvic
Chauliodontidae					
<i>Chauliodus</i>	51-62	6,7 (5-7)	10-12(10-13)	12,13(11-14)	7(6-8)
Melanostomiidae					
<i>Bathophilus</i>	38-45(33-50)	13-16 (9-18)	15-16 (9-18)	1-37	11-16(4-26)
<i>Eustomias</i>	56-69	21-25(20-30)	32-46	0-13	7(6-8)
<i>Opostomias</i>	60	21	24	1+4	8
<i>Tactostoma</i>	80-82	14-16	19-22	0	8-10
Malacosteidae					
<i>Aristostomias</i>	44-56	18-26	24-32	6-10 (3-17)	6
Idiacanthidae					
<i>Idiacanthus</i>	79-85	54-74	34(33-39)	0	6

Table 17

Photophore counts of stomioid genera. Most frequent count or range is followed by overall range or infrequent count in parentheses (Kawaguchi and Moser 1984, in part). Photophore groups as defined by Ahlstrom et al. (1984c) (see Table 9).

Taxon	Photophore groups					
	IP	PV	VAV	AC	OV	VA
Chauliodontidae						
<i>Chauliodus</i>	8-11	17-23	22-30	8-13	17-21	22-29
Melanostomiidae						
<i>Bathophilus</i>	5(4-6)	12-18	11-13(11-17)	5-7 (5-9)	13-14(10-16)	9-11 (8-17)
<i>Eustomias</i>	7-8(9)	27-33(24-36)	13-17(11-21)	17-23(15-25)	26-33(24-37)	13-18(12-22)
<i>Opostomias</i>	4+4	27	17	16	27	17
<i>Tactostoma</i>	8	46	19	12	43	18
Malacosteidae						
<i>Aristostomias</i>	5+3	15-17(14-19)	15-18	9-11(12)	16-19(14-20)	15-17(14-18)
Idiacanthidae						
<i>Idiacanthus</i>	IP+PV=31-36		16-18(15)	13-18	22-25	31-35(30-36)

Table 18

Pigment characters and gut structure (NT = not trailing, T = trailing freely) in larvae and transforming specimens of stomioids (Kawaguchi and Moser 1984, in part).

Taxon	Length of larvae (mm)	Length of transforming specimens (mm)	Dorsal myomere melanophores (n/myomere)	Epaxial myoseptum melanophores (n/myoseptum)	Hypaxial myoseptum melanophores	Gut structure
Chauliodontidae						
<i>Chauliodus macouni</i>	38-49	35-44	0	0	0	NT*
Melanostomiidae						
<i>Bathophilus flemingi</i>	2.9-23.8	—	1 to several	0	0	NT
<i>Eustomias</i> sp.	33	—	7 total	0	0	T
<i>Eustomias</i> sp.	13	—	7 total	0	0	T
<i>Eustomias</i> spp. (4 types)	6-45	—	5-11 total	0	0	T
<i>Opostomias mitsuui</i>	15-21	—	1	0-1 (2-3 posteriorly)	1-2 (3-5 posteriorly)	NT
<i>Tactostoma macropus</i>	5-44	49	0-1	0	1-3	NT
Malacosteidae						
<i>Aristostomias scintillans</i>	43-47	45	14 total	0	0	T
Idiacanthidae						
<i>Idiacanthus antrostomus</i>	4.5-71	67>	0	0	1	T

*Trails slightly.

MERISTICS

Vertebrae	Total: 56-60-62	
	Precaudal: 56-56-56	
	Caudal: 4-4-4	
Branchiostegal rays	16-X-21	
Caudal fin	X, 10+9, X	
Pelvic fin	Abdominal	
	R: 6-7-8	
Dorsal fin	R: 5-6-7	
Pectoral fin	R: 10-X-13	
Anal fin	R: 10-X-13	
Gill rakers	U: 3-3-3	L: 8-8-8

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathypelagic, 76-4231 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	>8 yr ^a

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	2.69-3.17 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Segmented
Envelope	Smooth, clear
Hatch size	6-7 mm SL

Incubation time/temp.**Pigment**

- Pigment in caudal region of yolksac larvae, but soon disappears

Diagnostic characters

- Wide perivitelline space, 0.5 mm
- Yolk segmented
- Large egg size

LARVAE

Preanal length	Long, >75% SL
Length at flexion	>15 mm SL
Length at transformation	Adult ~35-46 mm SL ^b
Sequence of fin development	Dorsal, anal, and pelvics form late in postflexion larvae in adult position

Pigment

- According to Kawaguchi and Moser (1984), unpigmented except for fan-shaped caudal finfold in yolksac larvae. We have not collected any yolksac larvae and therefore have not observed this pigment. A figure of a yolksac *C. sloani* is included for comparison only.

Diagnostic characters

- Morphology: Small head, elliptical eyes, long gut, slender body
- Median finfold shape
- Lack of pigmentation
- Slightly trailing gut

^aFitch and Lavenberg 1968^bMarked shrinkage at transformation.

Ref: Kawaguchi and Moser 1984.




Figure A, Matarese and Sandknop 1984; B, Mito 1961 (western Pacific specimen); C–D, Kawaguchi and Moser 1984.

MERISTICS

Vertebrae	Total: 44-46-48 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	10-10-10
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 15-15-17
Dorsal fin	R: 15-15-16
Pectoral fin	R: 4-5-7, jugular
Anal fin	R: 16-17-17
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Meso- and bathypelagic
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**


Preanal length	>75% SL
Length at flexion	
Length at transformation	≤25 mm SL
Sequence of fin development	Dorsal, anal, and caudal before pectorals and pelvics

Pigment

- One to several melanophores per myomere along dorsum with opposing series along ventral surface
- Head, finfolds, median fins
- Anterodorsal surface of gut and on hindgut
- Pigment more concentrated in larger specimens

Diagnostic characters

- Deep bodied
- Large head, jaws
- Gut large (highly developed trailing s-shaped terminus)
- No midlateral pigment
- Large pigmented median finfolds
- Origin of dorsal and anal fin opposite each other



Figures A–B, Beebe and Crane 1939 (North Atlantic specimens); C, Kawaguchi and Moser 1984.

MERISTICS

Vertebrae	Total: 72-X-78 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	X-X-X
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal? R: 6-X-8
Dorsal fin	R: 20-X-30
Pectoral fin	R: 0-X-13
Anal fin	R: 32-X-46
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Oregon, 42-46°N
Ecology	
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE** - Genus

Preanal length >75% SL (trailing gut)
Length at flexion
Length at transformation ~45 mm SL
Sequence of fin development
Pigment

- 5-11 large melanophores along dorsal midline
- Lower jaw symphysis

Diagnostic characters

- Morphology: Slender body, head elongate and flat
 - Gut: Long, slender, deflected ventrad at anal fin origin, trailing from body
 - Dorsal pigment patches
 - Origin of dorsal fin behind that of anal fin
- Distinguished from *Aristostomias scintillans* (p. 118) by
- Unpigmented gut
 - Lack of ventral body pigment
 - <20 dorsal spots (5-11)

^a Possibly 1-4 species in the area, although only one larval form has been collected.

Ref: Kawaguchi and Moser 1984, Ozawa and Aono 1986.




Figure A, Kawaguchi and Moser 1984 (after Regan 1916).

MERISTICS

Vertebrae	Total: X-X-X	
	Precaudal: X-X-X	
	Caudal: X-X-X	
Branchiostegal rays	10-X-15	
Caudal fin	X, 10+9, X	
Pelvic fin	Abdominal	
	R: 7-X-8	
Dorsal fin	R: 21-X-23	
Pectoral fin	R: 5-5-5	
Anal fin	R: 21-X-24	
Gill rakers	U: 2-2-2	L: 8-8-8

LIFE HISTORY

Range	S. California, 32-34°N, to Brit. Col., 48°30'-55°N
Ecology	Epi- and mesopelagic
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE

Preanal length
Length at flexion
Length at transformation
Sequence of fin development Caudal, dorsal, and anal
before pectorals and pelvics
Pigment

- One expanded melanophore per myomere along dorsum
- 1-2 melanophores on each hypaxial myoseptum
- Area under dorsal fin appears banded
- Dorsally on head
- Gill arch
- Gut terminus

Diagnostic characters

- Morphology
 - Moderately deep body
 - Large mouth
 - Large finfold
 - Elongate sloping snout
 - Eyes small
 - No trailing gut
- Pigment pattern
- Origin of dorsal fin slightly before that of anal fin
- 13-15 myomeres between pelvic and anal fins

Ref: Kawaguchi and Moser 1984, Ozawa and Aono 1986.




Figure A, Kawaguchi and Moser 1984 (western Pacific specimen).

MERISTICS

Vertebrae	Total: 80-X-82 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	13-13-13
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 8-9-10
Dorsal fin	R: 14-X-17
Pectoral fin	Absent in adults
Anal fin	R: 19-19-22
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesopelagic, 31-549 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Summer (Oregon) ^a Area: Mode: Migration:
Fecundity	Range/function: 24,000-66,000 (one spawn/year) ^a
Age at first maturity	6 yr (females) ^a
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.39-1.54 mm
No. of oil globules	One
Oil globule diameter	0.3-0.4 mm
Yolk	Segmented
Envelope	Smooth
Hatch size	5 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

- Oil globule large and ventral
- Egg round

LARVAE

Preanal length	>75% SL
Length at flexion	
Length at transformation	44-49 mm SL
Sequence of fin development	Caudal, dorsal and anal, pectorals and pelvics
Pigment	

- In early larvae, one melanophore per myomere along dorsum and 1-3 melanophores on hypaxial myosepta; postflexion larvae gradually lose the dorsal then ventral melanophores
- Lower jaw, isthmus, pectoral fin base, cleithrum, gut terminus, caudal peduncle

Diagnostic characters

- Morphology: Body elongate and flat, gut slender, finfold moderate
 - Pectoral fin lost at transformation
 - Decrease in pigment with development
 - No trailing gut
 - Finfolds unpigmented
- Distinguished from *Chauliodus macouni* (p. 108) by
- Presence of pigment
 - Posterior position of dorsal fin (origin of dorsal and anal fin opposite each other)

^a Fisher and Pearcy 1983

Ref: Kawaguchi and Moser 1984, Ozawa and Aono 1986.




Figure A, NWAFC original (B. Vinter); B, Kawaguchi and Moser 1984.

MERISTICS

Vertebrae	Total: 54-54-54 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	8-8-8
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 6-X-7
Dorsal fin	R: 21-X-23
Pectoral fin	R: 4-X-8
Anal fin	R: 25-X-29
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi- and mesopelagic
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length
Length at flexion
Length at transformation 45 mm SL
Sequence of fin development Dorsal, anal, and caudal; pectorals, pelvics

Pigment

- Series of paired melanophores along dorsum; initially 14 pairs which increase in number with development
- Paired ventral series develops initially posteriorly and increases in number with development
- Brain, snout, lower jaw, otic region, gular-isthmus region, caudal fin, gut

Diagnostic characters

- Morphology: Body and gut slender, trailing gut, large flat head, finfold moderate
- Distinguished from *Eustomias* spp. (p. 112) by
- Pigment on trailing gut
 - Pigment along ventral midline




Figure A, Kawaguchi and Moser 1984.

MERISTICS

Vertebrae	Total: 81-X-83 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	12-X-18
Caudal fin	X, 10+9, X
Pelvic fin^a	Abdominal R: 6-6-6
Dorsal fin	R: 54-X-66
Pectoral fin	Absent in adults
Anal fin	R: 28-X-43
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to N. California, 38-42°N
Ecology	Meso- and bathypelagic
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function: 14,000 ^a
Age at first maturity	
Longevity	>6 yr (females) ^a <1 yr (males) ^a

EARLY LIFE HISTORY DESCRIPTION

EGGS

- Diameter**
- No. of oil globules**
- Oil globule diameter**
- Yolk**
- Envelope**
- Hatch size**
- Incubation time/temp.**
- Pigment**

Diagnostic characters

LARVAE

- Preanal length**
- Length at flexion**
- Length at transformation** 35-70 mm SL
- Sequence of fin development** Caudal, dorsal and anal, pectorals,^b pelvics^c
- Pigment - Genus**
 - Melanophore on posterior margin of each hypaxial myomere
 - Isthmus
 - Series along trailing gut

Diagnostic characters - Genus

- Morphology
 - Extremely slender body
 - Elongate flat head
 - Elliptical eyes on stalks with cartilaginous support rods (27% BL)
 - Trailing gut
 - Small finfold
- Distinguished from *I. fasciola* by
 - Longer eye stalk and trailing gut

Since specimens are often damaged and pigment patterns are similar, the two species may be difficult to separate. In addition to the above characters, the two species differ in size at various stages of development. *I. antrostomus* larvae are 4.5-71.0 mm SL, transforming at sizes >67 mm SL. *I. fasciola* larvae are 16-28 mm SL, transforming at sizes between 35 and 48 mm SL.

^aFitch and Lavenberg 1968
^bPectoral fins lost at transformation.
^cPelvic fins develop in transforming females.

Ref: Kawaguchi and Moser 1984.




Figure A, Kawaguchi and Moser 1984.

MERISTICS

Vertebrae	Total: X-X-X Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	12-X-18
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 6-6-6
Dorsal fin	R: 56-X-77
Pectoral fin	Absent in adults
Anal fin	R: 38-X-54
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Meso- and bathypelagic
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

- Diameter**
- No. of oil globules**
- Oil globule diameter**
- Yolk**
- Envelope**
- Hatch size**
- Incubation time/temp.**
- Pigment**

Diagnostic characters


LARVAE

- Preanal length**
- Length at flexion**
- Length at transformation** 32-42 mm SL (males)
- Sequence of fin development**
- Pigment**
 - See *I. antrostomus* (p. 120)


Diagnostic characters

- See *I. antrostomus*
- Depending on length of larvae, length of trailing gut may differ

Ref: Kawaguchi and Moser 1984.



Figures A–F, Beebe 1934 (North Atlantic specimens).



Scopelomorpha: Aulopiformes
Myctophiformes

The myctophiforms (Myctophidae and Neoscopelidae) and aulopiforms (J. Nelson 1984) are treated together here. According to J. Nelson (1984), these orders consist of 14 families, and about 75 genera with 429 species. Although several families are benthic (e.g., synodontids), most families consist of deep-sea, pelagic, and benthopelagic forms. Many families (not myctophids) are synchronous hermaphrodites. The diverse early-life-history stages are generally well known and can be distinguished by morphological characters (head, gut, body) and pigment patterns. Additionally, photophore patterns and development are especially helpful in myctophid fishes, while gut pigment (i.e., number and development of peritoneal patches) is an important character distinguishing some of the other families.

Families in study area: **Scopelarchidae**
Notosudidae
Synodontidae
Bathysauridae
Paralepididae
Anopteroideidae
Alepisauridae
Neoscopelidae
Myctophidae

MERISTICS *B. dentata* (Chapman 1939)

Vertebrae	Total: 54-54-55 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	8-8-8
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 9-9-9
Dorsal fin	R: 6-7-8
Pectoral fin	R: 21-24-25
Anal fin	R: 17-19-21

MERISTICS *B. linguoidens*
(Mead and Böhlke 1953)

Vertebrae	Total: 64-64-64 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	X-X-X
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal
Dorsal fin	R: 8-X-9
Pectoral fin	R: 24-X-25
Anal fin	R: 28-X-30

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N ^a
Ecology	Mesopelagic, 200-1000 m ^b
ELH pattern	Oviparous, eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Synchronous hermaphrodites ^c Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

^a *B. linguoidens* only to Oregon, 42-46° N.^b *B. dentata* only, *B. linguoidens* larvae are incompletely known.^c Johnson 1974a^d Larvae of *Benthalbella* spp. are unique among scopolarchids in achieving a large size (50-100 mm SL) while retaining a purely larval form and then exhibiting a very rapid transformation. The largest known larvae of *B. linguoidens* = 85.5 mm SL, and may transform at larger sizes than *B. dentata*. Transformation is complete when peritoneal pigment first appears (uniformly in mesentery dorsal to gut from between pectoral fin bases to behind pelvic fins).

Ref: R.K. Johnson 1984.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE^b**

Preanal length Anus moves posteriorly during transformation

Length at flexion

Length at transformation 50 mm SL^d

Sequence of fin development Caudal; dorsal, anal, and dorsal part of pectoral; pelvics; ventral part of pectoral


Pigment

- No "accessory" pigment
- No "dermal" pigment; develops in transforming larvae and persists in adults


Diagnostic characters

- Gut expansion (in larvae, anus anterior to pelvic fin base but moves posteriorly during transformation)
- Pelvic fin origin ahead of dorsal fin origin
- Adipose fin develops posterior to anal fin
- No peritoneal pigment in larvae

B. linguoides



B. dentata



Figures A–C, Johnson 1974a.

MERISTICS

Vertebrae	Total: 58-X-61 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	10-10-10
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 9-X-10
Dorsal fin	R: 10-X-12
Pectoral fin	R: 10-X-14
Anal fin	R: 16-X-19
Gill rakers	U: 2-2-2 L: 17-X-19

LIFE HISTORY

Range	S. California, 32-34°N, to Aleutian Is., 51-55°N
Ecology	Mesopelagic, 500-800 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Synchronous hermaphrodites ^a Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	Probably small
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**


Preanal length	<50% SL
Length at flexion	
Length at transformation	25-45 mm SL
Sequence of fin development	Caudal, anal, dorsal, pectorals, pelvics (family)
Pigment	<ul style="list-style-type: none"> • Restricted to tail along hypural region

Diagnostic characters—Family

- Hypural pigment: Exact distribution of pigment on the hypural region is diagnostic for species within the family
- Long body, becoming compressed toward tail
- Depressed head
- Posteriorly protruding lobes in brain (corpus cerebelli)
- Conical mass of choroid tissue around narrowed eye; long axis of the narrow eye is horizontal rather than vertical or oblique as it is in most other narrow-eyed larvae
- Anus at midbody; widely separated from anal fin
- Maxillary teeth

^aOkiyama 1984

Ref: Okiyama 1984, Ozawa 1978.



A
(Scopelosaurus smithii) 13.4 mm SL

B
Hypural pigment
27.3 mm

C
175.0 mm

Figure A, Okiyama 1984 (southwestern Pacific specimen); B–C, Bertelsen et al. 1976 (North Atlantic specimens).

MERISTICS

Vertebrae	Total: 50-X-52 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	8-8-8
Caudal fin	7, 10+9, 6
Pelvic fin	Abdominal R: 8-8-8
Dorsal fin	R: 15-X-17
Pectoral fin	R: 16-X-17
Anal fin	R: 11-13-13
Gill rakers	U: 5-X-6 L: 14-X-16

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Bathybenthal
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Synchronous hermaphrodites ^a Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter Ovarian eggs of *B. ferox*
1.2 mm^b

Number of oil globules

Oil globule diameter

Yolk

Envelope

Hatch size

Incubation time/temp.

Pigment

Diagnostic characters

LARVAE^c

Preanal length 70% SL

Length at flexion

Length at transformation >83 mm SL (gradual)^d

Sequence of fin

development

Pigment

- Peritoneal patches, about five; about six short bars laterally on gut
- About 17 vertical bars along body, 8 shorter bars in interspaces between vertical bars
- Several spots on head and fin rays

Diagnostic characters—Family

- Elongate fins
- Anterior placement of dorsal and pelvic fins
- Raised dorsal and anal fin bases
- Long gut terminating in front of anal origin
- Pigment: Peritoneal (5-6 sections) and lateral bars

^aOkiyama 1984

^bWenner 1978

^c*B. mollis* larvae originally described as "*Macristium*" larvae (Johnson 1974b). An illustration of *B. ferox* is provided for comparison and to show fin rays intact.

^dChanges at transformation include shortening of fins, expansion of gape, backward shift of dorsal fin origin, and darkening of body surface, oral cavity, and peritoneum. According to Sulak et al. (1985), the large sizes attained by these larvae suggest a long oceanic existence prior to transformation.





Figure A, Johnson 1974b (Gulf of Mexico specimen); B, Marshall 1960 (Atlantic specimen).

MERISTICS

Vertebrae	Total: 60-62-63 Precaudal: 49-52-54 Caudal: 8-10-11
Branchiostegal rays	18-18-18
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 8-8-8
Dorsal fin	R: 11-X-13
Pectoral fin	R: 13-X-14
Anal fin	R: 12-X-14
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to N. California, 38-42°N ^a
Ecology	Epi- and mesodemersal
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.33-1.44 mm
No. of oil globules	None
Oil globule diameter	
Yolk	
Envelope	Hexagonal surface
Hatch size	<4.7 mm SL
Incubation time/temp.	
Pigment	

- Unpigmented until late stage of development when pigment first appears along dorsal midline
- Few scattered yolk sac spots
- Late-stage embryo resembles newly hatched larva with distinctive pigment along gut

Diagnostic characters

- Hexagonal surface similar to *Pleuronichthys coenosus* (p. 616) except hexagons are wider (0.047 mm vs. 0.035 mm) and irregularly arranged^b

LARVAE^c

Preanal length	>50% SL
Length at flexion	~10.5 mm SL
Length at transformation	
Sequence of fin development	Caudal, anal, dorsal, pectorals, pelvics

Pigment

- Seven evenly spaced pairs of "peritoneal patches" which form gradually
- Spot on postanal ventral midline midway between anus and tail
- Spot in area where hypurals are forming

Diagnostic characters

- Distinctive peritoneal patches (seven)
- Preanal finfold
- Deeper body than other *Synodus* spp.

^aDuring "El Niño" years, adults occur as far north as Puget Sound, Washington. Postflexion larvae were collected off Oregon during the strong El Niño summer of 1982 (B. Mundy, NMFS Southwest Fish. Cent., Honolulu Lab., 2570 Dole St., Honolulu, HI 96822, pers. commun., 1 Oct. 1986).

^bSee Sumida et al. 1979.

^cData on preflexion and postflexion larvae are from E.H. Ahlstrom notes. Illustrations are unavailable.




Figure A, Okiyama 1984.

MERISTICS

Vertebrae	Total: 84-X-87 Precaudal: 41-41-41 Caudal: X-X-X
Branchiostegal rays	8-8-8
Caudal fin	13, 10+9, 13
Pelvic fin	Abdominal R: 8-10-11
Dorsal fin	R: 9-12-12
Pectoral fin	R: 11-11-12
Anal fin	R: 26-26-33
Gill rakers	U: 3-X-9 L: 21-X-31

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epi-, meso-, and bathypelagic
ELH pattern	Oviparous, eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Synchronous hermaphrodites ^a Migration
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length	<50% SL, increasing with development
Length at flexion	
Length at transformation	~45 mm SL
Sequence of fin development	Caudal, anal, dorsal, pectorals, pelvics (family)
Pigment	<ul style="list-style-type: none"> • Tip of jaw, two dorsal and one ventral midline spot • Peritoneal patches increase during development to about seven • With development, head, dorsal and anal fin bases, and caudal peduncle

Diagnostic characters - Family

- Pigment, especially peritoneal patches^b
- Elongation of gut with development
- Morphology^c

Distinguished from *Notolepis rissoi* and *Paralepis atlantica* by


	<i>Lestidiops</i>	<i>Notolepis</i>	<i>Paralepis</i>
No. peritoneal patches (flexion larvae)	3	1	2
(postflexion larvae)	7	12	3
No. myomeres	84-87	72-74	76-83

^aOkiyama 1984


^bPeritoneal pigment patches are sequentially formed with gradual lengthening of the gut. The number of patches (called "sections" by Okiyama 1984) may be species-specific. In general, these patches develop by 5-10 mm SL and persist until 15-45 mm SL. Other characteristic paralepidid pigment may include dorsum of body, caudal peduncle, and caudal and pectoral fins.

^cGeneral morphological characters in paralepidids are a long compressed body, short gut increasing in length with development, head increasing in relative size with development, elongate snout, well-developed preanal finfold, and eyes initially ovoid, becoming round with development.


A



B



C



Figures A, C, Okiyama 1984; B, Moser 1981.

MERISTICS

Vertebrae	Total: 72-X-74 Precaudal: 37-X-39 Caudal: 39-X-41
Branchiostegal rays	8-8-8
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 8-X-12
Dorsal fin	R: 9-X-13
Pectoral fin	R: 10-X-13
Anal fin	R: 29-X-34
Gill rakers	U: 3-X-9 L: 18-X-36

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epi-, meso-, and bathypelagic
ELH pattern	Oviparous, eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Synchronous hermaphrodites ^a Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length	<50% SL, increasing with development
Length at flexion	~13-18 mm SL
Length at transformation	~45 mm SL
Sequence of fin development	Caudal, anal, dorsal, pectorals, pelvics (family)
Pigment	<ul style="list-style-type: none"> • Peritoneal patches, 1 increasing with development to 12 (see <i>Lestidiops ringens</i>) • Above and below notochord on caudal peduncle • Additional pigment along dorsum from dorsal fin origin to tail with development


Diagnostic characters

- See *L. ringens* (p. 134)
- Pigment, peritoneal patches
- Morphology

^aOkiyama 1984

Ref: Okiyama 1984.

A



B




C




D



E



F



Figures A-F, Rofen 1966a (North Atlantic specimens).

MERISTICS

Vertebrae	Total: 60-67-69 Precaudal: 38-40-41 Caudal: X-X-X
Branchiostegal rays	8-8-8
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 9-X-10
Dorsal fin	R: 9-X-12
Pectoral fin	R: 15-X-18
Anal fin	R: 20-X-23
Gill rakers	U: 7-X-9 L: 26-X-32

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59° N
Ecology	Epi-, meso-, and bathypelagic
ELH pattern	Oviparous, eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Synchronous hermaphrodites ^a Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE

Preanal length
Length at flexion
Length at transformation
Sequence of fin development Caudal, anal, dorsal, pectorals, pelvics (family)
Pigment


- Peritoneal patches, one increasing with development to three (see *Lestidiops ringens*, p. 134)
- Few spots above and below notochord in caudal peduncle region which, with development, form a patch
- Patch at base of dorsal and anal fin, on cranium, on upper jaw (late in development)

Diagnostic characters

- See *L. ringens*
- Pigment, peritoneal patches
- Morphology

^aOkiyama 1984

Ref: Okiyama 1984.



Figures A–E, Rofen 1966a (North Atlantic specimens).

MERISTICS

Vertebrae	Total: 76-80-83 Precaudal: 48-52-54 Caudal: 27-29-31
Branchiostegal rays	7-7-7
Caudal fin	14-15, 10+9, 14
Pelvic fin	Abdominal R: 9-X-11
Dorsal fin	Absent
Pectoral fin	R: 12-X-16
Anal fin	R: 14-X-17
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesopelagic
ELH pattern	Oviparous, eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Synchronous hermaphrodites ^a Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE^b**

Preanal length >50% SL
Length at flexion
Length at transformation <50 mm SL
Sequence of fin development Only pectoral anlagen present at 14.2 mm SL
Pigment

- No peritoneal pigment patches, instead uniform peritoneal pigment
- Snout, jaw
- Dorsal midline
- Tail tip

Diagnostic characters

- Morphology: Long slender body, large head, pointed snout
- Fleshy prolongation at jaw tips
- Two large canine teeth
- Gut extending to midbody

^aOkiyama 1984^bData are from one larva.

Ref: Okiyama 1984.

A



B



Figure A, Okiyama 1984; B, Rofen 1966b (North Atlantic specimen).

MERISTICS

Vertebrae	Total: 48-50-52 Precaudal: 19-23-26 Caudal: 24-27-31
Branchiostegal rays	8-8-8
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 8-X-10
Dorsal fin	R: 30-X-45
Pectoral fin	R: 12-X-15
Anal fin	R: 14-X-17
Gill rakers	U: 2-X-6 L: 16-X-24

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathypelagic, 0-1829 m
ELH pattern	Oviparous, eggs unknown, pelagic larvae
Spawning	Season: May (California) ^a Area: Mode: Synchronous hermaphrodites ^b Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE^c**

Preanal length
Length at flexion
Length at transformation^d
Sequence of fin development Pectorals, caudal, dorsal, anal, pelvics
Pigment
• Genus: Preflexion larvae appear to be unpigmented except for pectoral fins
• Postflexion
—Heavy gut pigment
—Pigment patch at anal fin origin
—Pectoral fins
—Saddle in area of adipose fin
—Patch along midline over gut

Diagnostic characters

Distinguished from other species of *Alepisaurus* by

- Four preopercular spines
- Bony ridges on head
- Pigment patch at anal fin base

For family

- Large head and mouth
- Prominent canines on dentary
- Small fins
- Short gut




^aFitch and Lavenberg 1968

^bOkiyama 1984

^cIllustrations of preflexion and postflexion *A. brevirostris* are provided for comparison.

^dTransformation is gradual.

Ref: Okiyama 1984.



Figures A, C, Rofen 1966c (specimens collected off Bermuda); B, Okiyama 1984 (Hawaiian specimen).

MERISTICS

Vertebrae^a	Total: 30-X-31 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	8-8-8
Caudal fin	6-8, 10+9, 5-7
Pelvic fin	Abdominal R: 8-8-8
Dorsal fin	R: 11-X-13
Pectoral fin	R: 18-X-19
Anal fin	R: 10-X-13
Gill rakers^b	U: 3-3-3 L: 8-8-8 (T: 10-X-12)

LIFE HISTORY

Range^c	Brit. Col., 48°30'-55°N
Ecology	Mesopelagic, 300-800 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS** (Ripe ovarian)

Diameter	0.83-0.98 mm
No. of oil globules	One
Oil globule diameter	0.39-0.61 mm
Yolk	
Envelope	
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE** (Postflexion only)

Preanal length	~70% SL
Length at flexion	
Length at transformation	Fin rays formed by 10.0 mm SL
Sequence of fin development	Pectorals, dorsal, anal, caudal, pelvics
Pigment	<ul style="list-style-type: none"> • Small dorsal patches on peritoneum and hindgut

Diagnostic characters

- Pigment patch over hindgut (not solid as in other myctophiforms)
 - Small preopercular spines
- Distinguished from most myctophiforms by
- Lack of peritoneal pigment patches
- For family
- Morphology
 - Deep body
 - Large head
 - Large pectoral fin
 - Round eye without choroid tissue
 - Lightly pigmented

^a Genus vertebral range 29-35.^b Gill raker counts vary geographically.^c Peden and Hughes (1986); previously only off Japan and Hawaii.




Figure A, Okiyama 1984 (southwestern Japan specimen).

MYCTOPHIDAE

Lanternfishes are worldwide in distribution, having 32 genera with 235 species found from the Arctic to the Antarctic. Myctophids generally have large eyes, large terminal mouths, photophores, and a black body color. In the Northeast Pacific, 20 species occur within two subfamilies: Myctophinae and Lampanyctinae. Most of the species (13) are members of the Lampanyctinae.

Myctophids are oviparous and presumably all produce planktonic eggs (Moser et al. 1984a). Eggs of only two species have been described: *Electrona rissoi* by Sanzo (1939) and *Lampanyctodes hectoris* by Robertson (1977). Eggs of both species are small (<1.0 mm), possess a single large oil globule, segmented yolk, smooth chorion, and at least *Lampanyctodes* eggs possess a fragile chorion. We have identified several types of myctophid eggs with these characteristics, in particular a fragile chorion. At least two of these types are probably *Diaphus* and *Stenobranchius* (based on egg characters and the presence of yolksac larvae of those species in the same samples).

The larval photophore complements and the sequence of appearance of photophores are useful characters. Most myctophids develop the Br2 during the larval period (in the Northeast Pacific the only exception is *Taaningichthys*). The Br2 is located posteroventral to the orbit but during transformation assumes a position beneath the orbit on the branchiostegal membrane. Three myctophine genera and eleven lampanyctine genera develop additional photophores during the larval period, with the Br2 always first to appear (Moser et al. 1984a). Of these 14 genera, only 4 are represented in our study area: *Ceratoscopelus*, *Lampadena*, *Diaphus*, and *Notoscopelus*.

The taxonomic section on myctophids here is arranged according to the intrafamilial classification of the group (Table 19), since certain larval characters reflect this classification (e.g., larval eye shape).

Subfamily Myctophinae	(Eyes elliptical)
Tribe Electronini	
	<i>Electrona rissoi</i>
	<i>Protomyctophum crockeri</i>
	<i>Protomyctophum thompsoni</i>
Tribe Myctophini	
	<i>Symbolophorus californiensis</i>
Tribe Gonichthyini	
	<i>Loweina rara</i>
	<i>Tarletonbeania crenularis</i>
	<i>Tarletonbeania taylori</i>
Subfamily Lampanyctinae	(Eyes mostly rounded)
Tribe Lampanyctini	
	<i>Ceratoscopelus townsendi</i>
	<i>Dorsadena</i> (?= <i>Lampadena</i>) <i>yaquinae</i> *
	<i>Lampadena urophaos</i>
	<i>Lampanyctus fernae</i>
	<i>Lampanyctus jordani</i>
	<i>Lampanyctus regalis</i>
	<i>Lampanyctus ritteri</i>
	<i>Parvilux ingens</i>
	<i>Stenobranchius leucopsarus</i>
	<i>Stenobranchius nannochir</i>
	<i>Taaningichthys bathyphilus</i>
Tribe Diaphini	
	<i>Diaphus theta</i>
Tribe Gymnoscopelini	
	<i>Notoscopelus resplendens</i>


*See Moser et al. (1984a); larvae are unknown.

Table 20
Meristic characters of the genera of Myctophidae (Moser et al. 1984a).

Taxon	Vertebrae	Fins								Branchiostegals
		Dorsal	Anal	Pectoral	Pelvic	Caudal*		Gill rakers		
						Upper	Lower	Upper	Lower	
Myctophinae										
<i>Electrona</i>	33-41	12-16	18-22	11-17	8	6-10	6-9	3-10	12-25	7-8
<i>Protomyctophum</i>	35-41	10-14	21-27	14-17	8-9	7-9	6-9	4-7	14-21	8-10
<i>Symbolophorus</i>	36-42	12-16	18-24	12-20	8	8-10	7-9	4-7	12-19	9
<i>Loweina</i>	37-39	10-13	13-17	9-12	7-9	6-7	6-7	2-3	5-10	9
<i>Tarletonbeania</i>	40-42	11-15	16-20	11-16	8	5-8	5-8	4-6	10-12	8
Lampanyctinae										
<i>Ceratoscopelus</i>	35-38	13-15	13-16	12-15	8	6-7	6-7	3-5	9-16	9
<i>Dorsadena</i>		14-15	12-14	15-16	8-9			4-5	12	
<i>Lampadena</i>	35-40	13-16	12-15	13-18	8	8	8-9	3-8	9-18	9
<i>Lampanyctus</i>	30-40	10-19	14-21	0-17	8	6-8	6-8	3-8	9-19	8-11
<i>Parvilux</i>	35-38	14-17	15-18	10-13	8	8	8-9	4-6	11-15	10-11
<i>Stenobranchius</i>	35-38	12-15	14-16	8-10	8	6-8	7-9	5-6	12-14	9-10
<i>Taaningichthys</i>	34-41	11-14	11-14	12-17	8	7-10	6-10	2-5	6-14	8-9
<i>Diaphus</i>	31-37	10-19	11-19	9-14	8	5-8	5-8	4-11	9-21	8-9
<i>Notoscopelus</i>	35-40	21-27	18-21	11-14	8-9	10-14	10-15	4-10	9-22	10

*Principal caudal fin ray count is 10+9. Only secondary fin ray counts are given.

PHOTOPHORE GROUP TERMINOLOGY



Moser and Ahlstrom 1972.

MERISTICS

Vertebrae	Total: 32-X-34
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	8-8-8
Caudal fin	6-8, 10+9, 6-7
Pelvic fin	Abdominal
	R: 8-8-8
Dorsal fin	R: 13-X-15
Pectoral fin	R: 13-X-16
Anal fin	R: 18-19-20
Gill rakers	U: 8-X-9 L: 18-X-21

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Epi- and mesopelagic
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	0.80-0.84 mm
No. of oil globules	One
Oil globule diameter	0.28 mm
Yolk	Segmented
Envelope	Smooth, fragile
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	50% SL, with development 50-75% SL
Length at flexion	6.2-7.0 mm SL
Length at transformation	~10 mm SL
Sequence of fin development	Caudal, pectorals, anal, dorsal, pelvics

Pigment


- Lower jaw symphysis (see 7.9 mm)
- Pectoral fin blade (see 6.3 mm)

Diagnostic characters

- Photophores: Early Br2, usually forms at transformation in other species
- Morphology: Body moderately slender, head large
- Gut saccular, s-shaped
- Space between anus and anal fin origin but not as large as in *Protomyctophum* spp.
- Eyes very narrow
- Transforms earlier than most myctophids (9-10 mm SL)

Distinguished from other myctophids with elliptical eyes by

- Pigment on lower jaw tip and pectoral fin
- Lack of pigment on postanal body



Figures A–D, Moser and Ahlstrom 1970.

MERISTICS

Vertebrae	Total: 36-X-37 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	8-X-12
Caudal fin	7-9, 10+9, 7-8
Pelvic fin	Abdominal R: 8-8-8
Dorsal fin	R: 11-12-13
Pectoral fin	R: 13-16-17
Anal fin	R: 19-22-24
Gill rakers	U: 4-5-6 L: 14-16-18

LIFE HISTORY

Range	South of southern California to Washington, 46-48°30'N ^a
Ecology	Epi- and mesopelagic, 0-500 m
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length	<50% SL
Length at flexion	7.2-8.8 mm SL
Length at transformation	12-15 mm SL, as large as 18 mm SL
Sequence of fin development	Caudal, pectorals, anal, dorsal, pelvics
Pigment	<ul style="list-style-type: none"> • Preflexion larvae unpigmented • Large melanophore over gut • Postanal ventral melanophores in juveniles

Diagnostic characters

- Single spot over gut and lack of pigment elsewhere (flexion and postflexion larvae)

For Genus

Distinguished from other myctophids with elliptical eyes by






- Head small, body slender
- Gut short
- Wide space between anus and anal fin
- Larvae relatively lightly pigmented
- Eyes moderately narrow
- No enlarged fins

Distinguished from *P. thompsoni* by

- Lack of postanal ventral melanophores in preflexion larvae and presence of only one spot over the gut in larger larvae

^aWisner 1974

Ref: Moser and Ahlstrom 1970, Moser et al. 1984a.



Figures A–E, Moser and Ahlstrom 1970.

MERISTICS

Vertebrae	Total: 37-38-39 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	8-X-12
Caudal fin	7-9, 10+9, 6-8
Pelvic fin	Abdominal R: 8-8-8
Dorsal fin	R: 11-12-13
Pectoral fin	R: 14-15-17
Anal fin	R: 21-23-25
Gill rakers	U: 3-4-4 L: 13-14-16

LIFE HISTORY

Range	Cent. California, 34-38°N ^a to Bering Sea, 54-66°N
Ecology	Epi- and mesopelagic
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length <50% SL
Length at flexion 7-10.0 mm SL
Length at transformation 14-17 mm SL
Sequence of fin development Caudal, pectorals, anal, dorsal, pelvics

Pigment


- Postanal ventral melanophores >10 in preflexion larvae decreasing to <10 in flexion larvae; not present in postflexion larvae
- Dorsolateral surface of gut in postflexion larvae

Diagnostic characters

- See *P. crockeri* (p. 150)
- Narrow eye
- Postanal ventral melanophore pattern

^aR.N. Lea, Calif. Dep. Fish Game, 2201 Garden Road, Monterey, CA 93940, pers. commun., 19 Feb. 1987.

Ref: Moser and Ahlstrom 1970.



Figures A–E, Moser and Ahlstrom 1970.

MYCTOPHIDAE *Symbolophorus californiensis* (Eigenmann and Eigenmann 1889)

MERISTICS

Vertebrae	Total: 37-X-40 Precaudal: 16-16-17 Caudal: 22-23-23
Branchiostegal rays	8-X-12
Caudal fin	8-9, 10+9, 8-9
Pelvic fin	Abdominal R: 8-8-8
Dorsal fin	R: 13-14-15
Pectoral fin	R: 15-17-19
Anal fin	R: 19-20-22
Gill rakers	U: 6-6-7 L: 15-16-17

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epi- and mesopelagic, 0-762 m
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Spring-summer ^a Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	7 yr ^a

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE

Preanal length
Length at flexion 8.0-10.0 mm SL
Length at transformation 23.0-24.0 mm SL
Sequence of fin
development Pectorals, pelvics, caudal,
anal, dorsal: pectoral fin
base large and wing-shaped
with supernumerary rays
developing first

Pigment


- Paired fins
- Lateral gut and anus
- Postanal ventral melanophores (preflexion)
- Also on snout, hindbrain, cleithral region, isthmus (lower jaw)

Diagnostic characters

- Distinguished from other myctophids with elliptical eyes by
- Broad head
 - Eyes slightly stalked with conical choroid mass (narrow)
 - Early pectoral fin development, pectorals large with wing-shaped base
 - Early pelvic fin development (unusual for myctophids)
 - Dorsal fin rays develop in finfold
 - Slightly enlarged median finfold
 - Pigment on paired fins

^aFitch and Lavenberg 1968

Ref: Moser and Ahlstrom 1970, 1974; Moser et al. 1984a.



Figures A–E, Moser and Ahlstrom 1970.

MERISTICS

Vertebrae	Total: 37-X-39 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	9-9-9
Caudal fin	6-7, 10+9, 6-7
Pelvic fin	Abdominal R: X-X-X
Dorsal fin	R: X-X-X
Pectoral fin	R: 16-X-18 (larvae) ^a R: 9-X-12 (adults)
Anal fin	R: X-X-X
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range ^b	South of southern California to N. California, 38-42°N, ^c and Oregon, 42-46°N ^d
Ecology	Unknown
ELH pattern	Oviparous, eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

^aLarvae have six more pectoral fin rays than adults.^bRange for larvae; adults widespread.^cMoser and Ahlstrom 1970^dRichardson and Percy 1977

Ref: Moser and Ahlstrom 1970.

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment


Diagnostic characters

LARVAE

Preanal length	55% SL increasing with development to 80% SL
Length at flexion	8.0-11.0 mm SL
Length at transformation	>20.0 mm SL
Sequence of fin development	Pectorals, caudal, anal and dorsal, pelvics
Pigment	<ul style="list-style-type: none"> • Preflexion and flexion larvae <ul style="list-style-type: none"> —Dark band of pigment between olfactory and optic lobes of the brain —Blotch develops medially to each pectoral fin base —Two blotches on dorsal surface of gut merging into one with development —Three postanal body blotches —Median finfold, lower pectoral ray —Dorsal and ventral margins of caudal peduncle

Diagnostic characters

- Among largest myctophid larvae at transformation (>20.0 mm SL)
 - Morphology: Early larvae slender but relative body depth doubles in later stages
 - Only Br2 develops during larval period
- Distinguished from all other myctophids except *Tarleton-beania crenularis* by
- Enlarged median finfolds
- Distinguished from *T. crenularis* (p. 158) by
- Eye moderately narrow, no choroid tissue on ventral surface of eye
 - Pigment: Band on head between eyes



Figures A–E (B, dorsal view), Moser and Ahlstrom 1970.

MERISTICS

Vertebrae	Total: 39-41-42 Precaudal: 17-17-19 Caudal: 22-23-24
Branchiostegal rays	8-9-11
Caudal fin	5-7, 10+9, 5-7
Pelvic fin	Abdominal R: 7-8-9
Dorsal fin	R: 11-12-14
Pectoral fin	R: 11-13-15
Anal fin	R: 17-18-20
Gill rakers	U: 4-5-6 L: 10-11-12

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi- and mesopelagic, 0-832 m
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Winter-spring; ^a Nov-Feb ^b Area: Mode: Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length	50% SL at hatching, 70% SL at 13-14 mm SL, ≥50% SL at transformation
Length at flexion	7.5-10.5 mm SL
Length at transformation	~19-21 mm SL
Sequence of fin development	Caudal, pectorals, anal, dorsal, pelvics
Pigment	<ul style="list-style-type: none"> • Preflexion: Postanal band located at posteriormost 4-5 myomeres, on anus, and over gut • Flexion: Head, finfold, 1-2 postanal melanophores • Postflexion: Dorsal midline anterior to fin

Diagnostic characters

Distinguished from *Loweina rara* (p. 156) by

- See *L. rara*


Distinguished from all other myctophids except *L. rara* by

- Finfold: Dorsal and ventral finfolds greatly enlarged and conspicuously pigmented (more on dorsal)
- Eyes narrow with lunate choroid mass
- Only Br2 develops during larval period
- Enlarged ventral pectoral fin ray (lost during transformation)
- Postanal pigment band

Note: *T. taylori* adults occur in the area but their larvae are inadequately known. Pertseva-Ostroumova (1964) illustrated a 12-mm SL larva as *T. crenularis taylori*, but it has no features that distinguish it from *T. crenularis*.

^aAhlstrom 1965

^bWang 1981



Figures A–E, Moser and Ahlstrom 1970.

MYCTOPHIDAE *Ceratoscopelus townsendi* (Eigenmann and Eigenmann 1889)

MERISTICS

Vertebrae ^a	Total: 35-36-38 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	8-X-12
Caudal fin	6, 10+9, 6-7
Pelvic fin	Abdominal R: 8-X-9
Dorsal fin	R: 13-14-15
Pectoral fin	R: 12-X-14
Anal fin ^a	R: 13-X-14
Gill rakers ^a	U: 4-4-5 L: 10-11-12

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epi- and mesopelagic
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE^b

Preanal length	46% SL, with development 58-61% SL
Length at flexion	6.2-7.0 mm SL
Length at transformation	16.6-21.0 mm SL (may be larger)
Sequence of fin development	Caudal, pectorals, anal, dorsal, pelvics

Pigment

- Generally light pigment
- Pair of melanophores at each side of gut at point of divergence of the free terminal section (retained throughout but becoming embedded with development)
- Series (~11) of postanal ventral melanophores in early larvae, with development usually coalescing to a single spot in postflexion larvae

Diagnostic characters

- Photophores: Br2 (7.0 mm SL); Vn (7.8 mm SL); PLO (8.7 mm SL; develop and remain at pectoral fin base throughout larval period but at transformation they migrate dorsally to just below the lateral line); PO5 (9.0 mm SL); during transformation the remaining photophores develop sequentially
- Distinguished from other myctophids with round eyes by
- Eye elliptical in preflexion larvae (although not perfectly round, the eye is easily differentiated from the narrow eyes of larval Myctophinae)
 - Single spot at anus
 - Preflexion larvae with ~11 postanal ventral melanophores


^aPeden and Hughes (1986) report counts for the following meristic characters:

Precaudal vertebrae	16
Caudal vertebrae	20
Anal fin rays	15
Gill rakers	6-7+1+15

^bH.G. Moser, NMFS Southwest Fish. Cent., P.O. Box 271, La Jolla, CA 92038, pers. commun., 9 Jan. 1984. Illustrations of small larvae are unavailable. A figure of a preflexion larva of *C. warmingi* is included for comparison.

Ref: Moser and Ahlstrom 1974, Moser et al. 1984a.

A



B




Figure A, Miller et al. 1979 (Hawaiian specimen); B, Moser and Ahlstrom 1974.

MERISTICS

Vertebrae^a	Total: 35-36-36 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	9-X-10
Caudal fin	8-9, 10+9, 8-9
Pelvic fin	Abdominal R: 8-8-8
Dorsal fin	R: 14-15-16
Pectoral fin	R: 15-16-17
Anal fin	R: 13-13-14
Gill rakers^a	U: 3-4-5 L: 9-X-11

LIFE HISTORY

Range	South of southern California to Washington, 46-48°30'N
Ecology	Epi- and mesopelagic
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length	50% SL
Length at flexion	By 5-6 mm SL
Length at transformation	19.6 mm SL
Sequence of fin development	Caudal, pectorals, anal, dorsal, pelvics

Pigment - Genus^b
(Pigment for *L. urophaos* more restricted to dorsal and ventral midline)

- Brain, nape, gut, swimbladder
- Double row of large melanophores along dorsal midline with an opposing double row of postanal ventral midline melanophores
- Some with more numerous dorsal and ventral melanophores and internal pigment above the notochord

Diagnostic characters

- Photophores: Br2, PLO, PO5, and Vn+PO1 develop during larval period
- Distinguished from other myctophids with round eyes by
- Dorsal midline pigment anterior to anus


^aPeden and Hughes (1986) report counts of the following meristic characters:

Total vertebrae	38
Precaudal vertebrae	17
Caudal vertebrae	21
Gill rakers	5+1+12

^bDiscussion of pigment is based on genus, since illustrations of small larvae of *L. urophaos* are not available. Figures of a preflexion larva of *L. luminosa* are included for comparison.

Ref: Moser and Ahlstrom 1972; Moser et al. 1984a; H.G. Moser, NMFS Southwest Fish. Cent., P.O. Box 271, La Jolla, CA 92038, unpubl.


A



Dorsal midline pigment
anterior to anus 4.8 mm SL

(Lampadena luminosa)

B




4.8 mm SL

(Lampadena luminosa)

Large spot at dorsal fin origin


C



~4 pairs opposing spots
along dorsal and ventral
midline

13.4 mm SL

D



19.6 mm SL

Figures A–B (A, dorsal view), Miller et al. 1979 (Hawaiian specimens); C–D, Moser and Ahlstrom 1972.

Of the four *Lampanyctus* species found in the study area, we can identify larvae of only *L. regalis* and *L. ritteri*. Generic characters include:

- Pigment above brain
- Myoseptal trunk pigment increasing during postflexion to cover most of anterior trunk at transformation
- Slender body; deepens and becomes robust in some species
- Deep head
- Gut short in early preflexion larvae; with development lengthens to midbody

The following meristic information may aid in identifying the four *Lampanyctus* species.

Species	Distribution	Vertebrae	Fins			
			Dorsal	Anal	Pectoral	Pelvic
<i>Lampanyctus fernalae</i>	N. Calif. - Oregon	36-38	12-14	16-18	12-14	8
<i>Lampanyctus jordani</i>	S. Calif. - Bering Sea	38-40	10-12	17-20	14-17	8
<i>Lampanyctus regalis</i>	SSC - Bering Sea	36-39	14-16	17-19	12-14	8
<i>Lampanyctus ritteri</i>	SSC - Bering Sea	35-38	12-15	16-19	10-13	8-9

MERISTICS

Vertebrae	Total: 36-X-39 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	9-9-9
Caudal fin	6-8, 10+9, 6-8
Pelvic fin	Abdominal R: 8-8-8
Dorsal fin	R: 14-15-16
Pectoral fin	R: 12-13-14
Anal fin	R: 17-17-19
Gill rakers	U: 4-4-4 L: 9-10-10

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi- and mesopelagic
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE^a**

Preanal length	38% SL preflexion, 54% SL at flexion, 67% SL postflexion
Length at flexion	>4 mm SL, <7 mm SL
Length at transformation	>10 mm SL, <28 mm SL
Sequence of fin development	Caudal, pectorals, anal, dorsal, pelvics (pelvics may form early)

Pigment

- Preflexion larvae are unpigmented except for spots on lower jaw tip and snout
- Head: Jaw tips, internal along snout, postorbital, opercle
- Pectoral and pelvic fins
- Large spot at adipose fin

Diagnostic characters

- Photophores: Only Br2 develops during larval period (7.8 mm SL); remaining photophores develop synchronously at end of transformation period
- Distinguished from other myctophids with round eyes by
- Morphology: Initially slender with a small head; with development a deep broad head and body with elongate snout develops; pectorals moderately large
 - Jaws large with teeth
 - Pigment: Large spot at adipose fin

^aH.G. Moser, NMFS Southwest Fish. Cent., P.O. Box 271, La Jolla, CA 92038, pers. commun., 9 Jan. 1984.

Ref: Moser et al. 1984a; Moser, unpubl.





Figure A, NWAFC original (B. Vinter); B, Moser and Ahlstrom 1974.

MERISTICS

Vertebrae	Total: 35-36-38 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	9-9-9
Caudal fin	7-8, 10+9, 7-8
Pelvic fin	Abdominal R: 8-X-9
Dorsal fin	R: 12-13-15
Pectoral fin	R: 10-11-13
Anal fin	R: 16-18-19
Gill rakers	U: 4-4-4 L: 9-10-11

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathypelagic, 20-1098 m
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE^a**

Preanal length	<50% SL in preflexion, with development 50% SL
Length at flexion	>4 mm SL, ~6 mm SL
Length at transformation	>10 mm SL
Sequence of fin development	Caudal, pectorals, anal, dorsal, pelvics
Pigment	<ul style="list-style-type: none"> • Head: Snout, lower jaw, opercle; several spots anterodorsal to eye persist throughout development • Dorsal and ventral midline on tail • Anteroventral to liver

Diagnostic characters

- See *L. regalis* (p. 166)
 - Photophores: Only Br2 develops during larval period; remaining photophores develop synchronously at end of transformation period
- Distinguished from other myctophids with round eyes by
- See *L. regalis*
- Distinguished from *L. regalis* by
- Morphology: Body and head moderately deep; jaws, teeth, and pectorals moderate in size

^aH.G. Moser, NMFS Southwest Fish. Cent., P.O. Box 271, La Jolla, CA 92038, pers. commun., 9 Jan. 1984.

Ref: Ahlstrom 1965; Moser and Ahlstrom 1974; Moser, unpubl.




Figure A, Moser and Ahlstrom 1974.

MERISTICS

Vertebrae	Total: 36-37-38 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	8-X-12
Caudal fin	8, 10+9, 8
Pelvic fin	Abdominal R: 8-8-8
Dorsal fin	R: 14-X-17
Pectoral fin	R: 10-X-13
Anal fin	R: 15-X-18
Gill rakers	U: 4-5-6 L: 12-13-15

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Epi- and mesopelagic
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE

Preanal length <50% SL, with development increasing to 50% SL

Length at flexion

Length at transformation

Sequence of fin development Caudal, pectorals, anal, dorsal, pelvics

Pigment

- Early preflexion larvae probably unpigmented with pigment developing by flexion stage
- Postflexion larvae have pigment above brain, internally above otic region, lateral to cleithrum, and in anteroventral region of liver
- Postanal body with 1-2 dorsal midline melanophores and 1 ventral midline melanophore at caudal peduncle

Diagnostic characters

- Photophores: Only Br2 forms during the larval period
- Distinguished from other myctophids with round eyes by
- Dorsal midline melanophores restricted to 1-2 spots posterior to adipose fin
 - Head and eyes large with the body tapered




Figure A, Moser and Ahlstrom 1974.

MYCTOPHIDAE *Stenobranchius leucopsarus* (Eigenmann and Eigenmann 1890)

MERISTICS

Vertebrae	Total: 36-37-38 Precaudal: 14-15-16 Caudal: 20-21-22
Branchiostegal rays	8-X-12
Caudal fin	6-8, 10+9, 7-8
Pelvic fin	Abdominal R: 8-8-8
Dorsal fin	R: 12-X-14
Pectoral fin	R: 9-X-11
Anal fin	R: 14-15-16
Gill rakers	U: 5-5-6 L: 12-13-14

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathypelagic, 0-2896 m
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Winter-spring (Oregon); ^a Nov-Aug (Calif.); ^b Dec-Feb (Moss Landing, Calif.) ^c Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	3 yr (California) ^b 4 yr (Oregon) ^a
Longevity	3-4 yr ^b

^aSmoker and Percy 1970

^bFast 1960

^cWang 1981

Ref: Ahlstrom 1965, 1972; Moser and Ahlstrom 1974.

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE

Preanal length <50% SL
Length at flexion >5.3 mm SL, ~6-7 mm SL
Length at transformation >12.5 mm SL;
10-19 mm SL^b
Sequence of fin development Caudal, pectorals, anal, dorsal, pelvics
Pigment


- Gut melanophores: Early preflexion, lateral spots at pectoral fin base, midgut, and anus
- Series of postanal ventral melanophores (>15) in preflexion larvae which coalesce to 2 or 3 spots in postflexion larvae
- Postflexion: Above brain, nape, and embedded in trunk myosepta on each side of dorsal midline (not shown)

Diagnostic characters

- Photophores: Only Br2 forms in the larval period
- Distinguished from other myctophids with round eyes by
- Gut pigment: Three spots in preflexion larvae (pectoral fin base, midgut, and anus); with development spots are retained, although their position shifts posteriorly
- Series of postanal ventral melanophores (>15) in preflexion larvae
- Distinguished from *Diaphus theta* (p. 176) by
- Lack of pigment at caudal base
- Number of postanal ventral melanophores (>15): Preflexion larvae usually have >20; postanal ventral midline melanophores appear more embedded in *D. theta* (see figure)

S. nannochir larvae are unknown. The following meristic information may aid in identification.

Total vertebrae	36-38
Dorsal fin rays	13-14
Anal fin rays	14-16
Pectoral fin rays	9-10
Pelvic fin rays	8



Figures A–B, D, NWAFC originals (B. Vinter); C, Moser and Ahlstrom 1974.

MERISTICS

Vertebrae	Total: 34-X-36 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	8-X-12
Caudal fin	X, 10+9, X
Pelvic fin	Abdominal R: 8-8-8
Dorsal fin	R: 11-X-14
Pectoral fin	R: 12-X-14
Anal fin	R: 12-13-14
Gill rakers	U: 3-3-3 L: 6-X-10

LIFE HISTORY

Range	Cent. California, 34-38°N, ^a to Oregon, 42-46°N
Ecology	Meso- and bathypelagic
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length
Length at flexion
Length at transformation
Sequence of fin development Caudal, pectorals, anal, dorsal, pelvics
Pigment - Genus^b

- Above brain, otic region
- One to several opposing melanophores at postanal dorsal and ventral midline
- Late postflexion larvae may develop melanophores along each side of dorsal midline
- Base of caudal rays
- Series of embedded melanophores above spinal column

Diagnostic characters - Genus

- No photophores in larvae
- Distinguished from other myctophids with round eyes by
- Morphology: Body slender, lower jaw projects beyond upper jaw
 - Embedded melanophores above spinal column
 - Opposing dorsal and ventral melanophores on caudal peduncle

^aWisner 1974^bInformation on *T. minimus* larvae is provided for comparison since *T. bathyphilus* larvae are unknown.


Taaningichthys minimus

Figure A–B, Moser and Ahlstrom 1972 (southwestern Pacific specimens).

MERISTICS

Vertebrae	Total: 34-35-36 Precaudal: 15-16-16 Caudal: 18-19-20
Branchiostegal rays	9-9-9
Caudal fin	6-8, 10+9, 6-8
Pelvic fin	Abdominal R: 7-X-8
Dorsal fin	R: 11-13-15
Pectoral fin	R: 10-X-12
Anal fin	R: 12-X-14
Gill rakers	U: 5-X-7 L: 12-15-16

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi- and mesopelagic, 0-792 m
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment


Diagnostic characters**LARVAE**

Preanal length
Length at flexion <6.9 mm SL
Length at transformation
Sequence of fin development Caudal, pectorals, anal,
dorsal, pelvics
Pigment
• Head unpigmented
• Anteroventral surface of liver, midgut, base of
caudal fin rays, swimbladder
• Postanal ventral melanophores in preflexion:
Numerous (<15)

Diagnostic characters

- Number of photophores in larvae^a
- Distinguished from other myctophids with round eyes by
- Body moderately slender
 - Head moderate in size, unpigmented
 - Postanal ventral melanophores (<15)
 - Melanophores at base of caudal fin rays
- Distinguished from *Stenobranchius leucopsarus* (p. 172) by
- See *S. leucopsarus*

^aSequence of formation of photophores: Br2, PO5, PO1 [VO1, PO2, OP2, VO5, PO3, PO4, VLO] (photophores in brackets appear in late larval period).



Figures A–B, D, NWAFC originals (B. Vinter); C, Moser and Ahlstrom 1974.

MERISTICS

Vertebrae	Total: 35-37-38 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	9-9-9
Caudal fin	11-14, 10+9, 10-14
Pelvic fin	Abdominal R: 8-X-9
Dorsal fin	R: 21-22-23
Pectoral fin	R: 11-12-13
Anal fin	R: 18-19-20
Gill rakers	U: 6-6-7 L: 13-14-15

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epi- and mesopelagic
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS** - Genus

Diameter	0.26 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**^{b,c}

Preanal length	<50% SL preflexion, with development 50% SL
-----------------------	--

Length at flexion**Length at transformation**

Sequence of fin development	Caudal, pectorals, anal, dorsal, pelvics
------------------------------------	---

Pigment

- Preflexion: Jaw tips, above brain and swimbladder, lateral cleithral
- Postflexion: Lower jaw, hindbrain, nape, double row of melanophores along dorsal midline beginning at midbody, series along anal fin base, caudal fin base, pelvic and anal fin, internal notochord pigment on trunk

Diagnostic characters


- Photophores: Br2, PO5, Vn, and PLO develop during larval period
- Distinguished from other myctophids with round eyes by
- Morphology: Body moderately deep, head and eyes large
 - Midlateral and dorsal fin base pigment in postflexion larvae
 - Highest dorsal fin ray count among myctophids in study area (21-23)
 - Head pigment: Jaw tips, snout, opercle and brain

^a According to Peden and Hughes (1986) northern records of *N. resplendens* are probably *N. japonicus*.

^b H.G. Moser, NMFS Southwest Fish. Cent., P.O. Box 271, La Jolla, CA 92038, pers. commun., 9 Jan. 1984.

^c Both *N. resplendens* and *N. japonicus* are described and illustrated in Ozawa (1986d). Specimens from the eastern Atlantic are figured here, but, according to Ozawa (1986d), they may differ in pigmentation and in photophore development from specimens found in the Pacific.

Ref: Badcock and Merrett 1976; Moser and Ahlstrom 1972; Ozawa 1986d; Moser, unpubl.



Figures A–C, Badcock and Merrett 1976 (eastern Atlantic specimens); D, Ahlstrom and Moser 1974.



Gadiformes

Gadiform fishes include some of the most commercially important coldwater marine species in the world: the codfishes and their relatives. The order contains between 7 and 13 families, depending on the source, and well over 400 species. Fishes of this order are found worldwide and are generally bathypelagic (except for one freshwater species). Most species are elongate with some having long dorsal and/or anal fins. Pelvic fins are thoracic or jugular, and chin barbels are present in many. Although they are easily recognized and well studied, evolutionary relationships are poorly understood and most recent authors now propose a multiple origin for at least the Gadidae (Markle 1989). The early-life-history stages are well known, especially for the gadids and merlucciids, and have been studied for well over 100 years (Fahay and Markle 1984). Most eggs are pelagic, spherical, and have a single oil globule (lacking in some gadids). Larvae are commonly identified by their distinctive pigment patterns, coiled gut, and general tadpole shape. Morids, melanonids, and macrourids are less well known.

Families in study area: **Moridae**
Melanonidae
Merlucciidae
Gadidae
Macrouridae

MORIDAE

Morids are benthopelagic fishes distributed throughout the world's oceans and are represented by two species in two genera in the study area. Very little is known of the early life history of the codlings. For the few species described, eggs are 0.52-1.16 mm in diameter, have one oil globule, and a smooth chorion (Fahay and Markle 1984). The scarcity of small fish in bottom trawls indicates the very young, and perhaps eggs, are pelagic (Iwamoto 1975). Most morids have precocious pelvic fin ray development. Usually the pelvic fin rays are elongate and probably undergo some ontogenetic reduction. Figures of larvae of *Physiculus nematopus* and *Laemonema* sp. are provided for comparison, since larvae of species in our area are unknown.

Table 21
Meristic characters of family Moridae.

Taxon	Distribution	Vertebrae		Fins					Gill rakers			
		Precaudal (Total)	Caudal	First Dorsal	Second Dorsal	Anal	Pectoral	Pelvic	Caudal (Total)	Upper	Lower	Branchiostegals
<i>Antimora</i> *		24-25 (57-61)	33-35	4-7	48-56	36-49	17-25	5-7				
<i>Antimora microlepis</i>	SSC - Bering Sea	24-25	33	4-5	50-55	37-42	20	6-7	14,5+4,12	5	15	7
<i>Laemonema</i> *		15-17 (50-63)	42-45	5-6	48-75	45-72	15-26	1-3 (usually 2)				
<i>Laemonema longipes</i>	Bering Sea	(51)		6	50-52	45-50	15-17	2-3	(22-25)	5-7	19-20	6

*Data are from Fahay and Markle (1984).



A *Physiculus nematopus***B** *Laemonema* sp.

Figure A, Fahay and Markle 1984; B, Fahay 1983 (North Atlantic specimen; identification is tentative).

MERISTICS

Vertebrae	Total: 58-X-62 Precaudal: 13-13-13 Caudal: 47-47-47
Branchiostegal rays	7-7-7
Caudal fin	23-25, 6+3, 22-25
Pelvic fin	Thoracic R: 5-X-7
Dorsal fin	R: 67-X-80 ^a
Pectoral fin	R: 12-X-16
Anal fin	R: 52-X-61 ^a
Gill rakers	U: 3-X-4 L: 6-X-11

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Mesopelagic, 200-1000 m
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE^b

Preanal length
Length at flexion
Length at transformation
Sequence of fin
development
Pigment

Diagnostic characters

- Large number of secondary caudal fin rays (23-25 upper and 22-25 lower)

^aD. Markle, Dep. Fish. Wildl., Oregon State Univ., Corvallis, OR 97331-3803, pers. commun., 8 May 1986.

^bSpecimens <17 mm SL have not been identified.

Ref: Fahay and Markle 1984.




Figure A, NWAFC original (B. Vinter, Atlantic specimen); B, Fahay and Markle 1984 (South Atlantic specimen).

MERISTICS

Vertebrae	Total: 52-53-55 Precaudal: 23-24-25 Caudal: 29-29-30
Branchiostegal rays	7-7-7
Caudal fin	16-18, 6+2-3, 14-17
Pelvic fin	Thoracic R: 6-X-8
Dorsal fin	1st: 10-11-13 2nd: 37-40-44
Pectoral fin	R: 14-16-16
Anal fin	R: 37-40-44
Gill rakers	U: 4-X-5 L: 13-16-18

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi- and mesopelagic, 0-914 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Jan-June ^a Area: Offshore pelagic; ^b 130-500 m ^c Mode: Schools ^d Migration: South and offshore to spawn (southern subpopulation); within Puget Sound and Straits of Georgia (northern subpopulation) ^b
Fecundity	Range/function: 3419 ^e -496,000 ^f
Age at first maturity	2 yr ^d -4 yr ^g
Longevity	17 yr ^h

^aHart 1973^bThe main offshore population spawns offshore off southern California (Francis and Bailey 1983). Disjunct populations in the Puget Sound-Straits of Georgia area spawn in restricted areas within these regions.^cBailey et al. 1982^dKimura and Milliken 1977^eMacGregor 1971^fMacGregor 1966^gBest 1963^hBeamish 1979

Ref: Ahlstrom and Counts 1955, Fahay and Markle 1984.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.07-1.18 mm (1.12 mm)
No. of oil globules	One
Oil globule diameter	0.27-0.34 mm
Yolk	May appear granular
Envelope	Clear, smooth
Hatch size	2.4 mm NL
Incubation time/temp.	3.1 d/16.6°C
Pigment	<ul style="list-style-type: none"> • On yolk • Characteristic late-stage patches on embryo

Diagnostic characters

- Pigment pattern
- Presence of oil globule
- Eye forms early during embryonic development

LARVAE


Preanal length	<50% SL
Length at flexion	~10 mm SL
Length at transformation	~30-35 mm SL
Sequence of fin development	Caudal, pelvics, 1st dorsal, 2nd dorsal and anal, pectorals

Pigment

- Preflexion larvae: One postanal bar, one large dendritic spot several myomeres posterior to anus; with development, bar becomes diffuse, postanal body pigment increases, especially dorsolaterally
- On head, crown, and snout; increasing with development
- Dorsolaterally on gut

Diagnostic characters

- Gadoid shape but absence of >1 distinct postanal pigment bar
- Two dorsal fins and one anal fin
- Specimens from Puget Sound and Straits of Georgia have pigmented pectoral rays



Figures A–F, Ahlstrom and Counts 1955.

The family Gadidae is represented in the Northeast Pacific Ocean and Bering Sea by five species: Walleye pollock, *Theragra chalcogramma*; Pacific cod, *Gadus macrocephalus*; Pacific tomcod, *Microgadus proximus*; saffron cod, *Eleginus gracilis*; and Arctic cod, *Boreogadus saida*. Identification of gadid larvae prior to 1980 was not possible. Gadid larvae collected during ichthyoplankton studies were routinely reported as "Gadidae." Recent taxonomic studies have continued since 1980 and have allowed the specific identification of all five species in the area (e.g., Matarese et al. 1981 and Dunn and Vinter 1984). Identification is based primarily on differences in pigmentation patterns and meristic characters. The available knowledge of the general early life histories of these five species was recently reviewed by Dunn and Matarese (1987).

Table 22
Characters useful in separating larvae of *Eleginus gracilis*, *Boreogadus saida*, *Gadus macrocephalus*, *Theragra chalcogramma*, and *Microgadus proximus* at specific size ranges (Dunn and Vinter 1984, in part).

Character	Size range (mm SL)	<i>E. gracilis</i>	<i>B. saida</i>	<i>G. macrocephalus</i>	<i>T. chalcogramma</i>	<i>M. proximus</i>
Pigment						
Preanal region						
Ventral gut	4.4-13.5	Double row of small melanophores	Absent in larvae <10mm; spots present anterior to pectoral fins in larvae >10 mm	Large melanophores medially	Relatively few scattered spots, more anterior than posterior	Large melanophores present, more anterior than posterior
Line of lateral pigment	10-18	Begins anterior to anus	Begins under second dorsal fin	Begins anterior to anus	Begins under second dorsal fin	Begins just anterior to anus
Postanal region						
Length of ventral stripes (based on no. of melanophores)	4-6	Both longer than dorsal stripes	Both shorter than dorsal stripes	Anterior stripe longer than dorsal stripe (<5.3 mm)	Posterior stripe longer than dorsal stripe	Anterior stripe longer than dorsal stripe (<5 mm)
No. of myomeres from vertical end of anus to anterior end of first ventral pigment stripe	4-6	4-6	5-7	1-3 (reaches vent by ~5.3 mm)	4-5	1-3 (reaches vent by ~5.0 mm)
Length at which dorsal pigment forms a continuous line	4-15	~10 mm	~7 mm	~5-6 mm	~13 mm	~13 mm
Length at which ventral pigment forms a continuous line	4-10	~7 mm	~10 mm	~5-6 mm	Never merge	~5-6 mm
On ventral margin of body	10-15	In double row on each side of midline	Pigment on midline and scattered on each side	Single row on each side of midline	Pigment on midline and a single row on each side	Single row on each side of midline anteriorly, single row on midline posteriorly
Morphologic						
Position of vent relative to dorsal fins	15-20	Under second dorsal	Under second dorsal	Under second dorsal	Between first and second dorsal	Between first and second dorsal (ultimately under first dorsal)
Meristic						
No. of rays on superior hypural	>13	5	4	4	4	5

MERISTICS

Vertebrae	Total: 53-X-58 Precaudal: 18-X-20 Caudal: 35-X-39
Branchiostegal rays	7-7-7
Caudal fin	21-25, 4+2, 21-25
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	1st: 10-X-17 2nd: 11-X-18 3rd: 16-X-24
Pectoral fin	R: 18-X-19
Anal fin	1st: 13-X-21 2nd: 17-X-23
Gill rakers	U: 8-X-9 L: 29-X-34

LIFE HISTORY

Range	Bering Sea, 54-66°N, to Arctic, not specific
Ecology	Epi- and mesopelagic, 0-731 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Oct-Mar ^a Area: Nearshore ^b Mode: Schools ^b Migration: Nearshore to spawn ^b
Fecundity	Range/function: 9000-21,000 ^a
Age at first maturity	3 yr (females) ^b 2-3 yr (males) ^b
Longevity	7 yr ^b

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.53-1.90 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Smooth
Hatch size	6 mm SL
Incubation time/temp.	
Pigment	


Diagnostic characters**LARVAE**

Preanal length	<50% SL
Length at flexion	11-17 mm SL
Length at transformation	17-30 mm, pelagic until 30-45 mm SL
Sequence of fin development	Caudal, dorsals and anals, pectorals, pelvics
Pigment	<ul style="list-style-type: none"> • Presence of bars • Shorter ventral stripes • Mediolateral pigment

Diagnostic characters (see Table 22)

- Ventral pigment on midline and scattered on each side
 - Rays on superior hypural = 4
- Distinguished from *Theragra chalcogramma* by
- Ventral pigment on midline and scattered on each side. In *T. chalcogramma*, ventral pigment consists of a row along midline and a single row on each side.

^aBain and Sekerak 1978^bCraig et al. 1982



Figures A–D (D, ventral view), Dunn and Vinter 1984; E, NWAFC original (B. Vinter).

MERISTICS

Vertebrae	Total: 57-62-64 Precaudal: 21-X-24 Caudal: 37-X-41
Branchiostegal rays	7-7-7
Caudal fin	22-25, 5+2, 23-26
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	1st: 11-X-16 2nd: 15-X-23 3rd: 18-X-21
Pectoral fin	R: 18-X-21
Anal fin	1st: 20-X-24 2nd: 19-X-22
Gill rakers	U: 2-X-3 L: 17-X-20

LIFE HISTORY

Range	Gulf of Alaska, 54-60°N, to Arctic, not specific
Ecology	Nearshore shelf pelagic, 2-75 m
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Winter ^a Area: Shallow nearshore (2-10 m) ^a Mode: Migration: To shallow water for spawning ^a
Fecundity	Range/function: 28,900-190,700 ^a
Age at first maturity	2 yr ^a
Longevity	9 yr ^a

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.0-1.7 mm (1.3-1.7 mm)
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous, dense
Envelope	Smooth, thick
Hatch size	3.5 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	<50% SL
Length at flexion	11-17 mm SL
Length at transformation	24-27 mm SL
Sequence of fin development	Caudal, dorsals and anals (nearly simultaneous), pelvics, pectorals

Pigment







- Presence of bars
- Ventral stripes longer than dorsal stripes
- Ventral pigment in double row on each side of midline
- Mediolateral pigment begins anterior to anus
- Double row of melanophores along ventral surface of gut

Diagnostic characters (see Table 22)

- Pigment
 - Double row of melanophores along ventral surface of gut
 - Double row of ventral pigment on each side of midline
- Rays on superior hypural = 5

^aWolotira 1985

Ref: Dunn and Matarese 1984, Dunn and Vinter 1984.



Figures A–F (D, ventral view), Dunn and Vinter 1984.

MERISTICS

Vertebrae	Total: 49-54-56 Precaudal: 18-X-21 Caudal: 31-X-35
Branchiostegal rays	7-7-7
Caudal fin	23-24, 4+2, 21-22
Pelvic fin	Thoracic R: 6-X-7
Dorsal fin	1st: 10-12-16 2nd: 11-15-22 3rd: 10-X-21
Pectoral fin	R: 19-20-22
Anal fin	1st: 16-18-27 2nd: 12-17-25
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathybenthal
ELH pattern	Oviparous, demersal eggs, pelagic larvae (small larvae demersal) ^a
Spawning	Season: Jan ^b -July ^c Area: Semi-demersal (73-265 m) ^c Mode: Schools ^b Migration: To deepwater ^d
Fecundity	Range/function: 228,000 ^e -3 million ^f / $F = 12.024 \times L^{2.959}$ ^g
Age at first maturity	2 yr ^d
Longevity	13 yr ^f

^aWalters (1984) reported small larvae are demersal whereas Rugen and Matarese (1988) reported newly hatched larvae quickly rise to above 50 m.

^bMiller et al. 1978

^cHirschberger and Smith 1983

^dKetchen 1961

^eThompson 1962

^fAndriashev 1954

^gKarp 1982

Ref: Dunn and Matarese 1984, 1987; Dunn and Vinter 1984; Matarese et al. 1981; Walters 1984.

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	0.98-1.08 mm (1.02 mm)
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous, dense
Envelope	Smooth, thick
Hatch size	3-4 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	<50% SL
Length at flexion	10-17 mm SL
Length at transformation	25-35 mm SL

Sequence of fin development

Pigment

- Presence of bars
- Ventral gut with large melanophores (in larvae >20 mm SL, small spots occur in two rows along ventral surface of gut)
- Stripe continuity (~5-6 mm SL)
- Mediolateral pigment begins anterior to anus (~5-6 mm SL)

Diagnostic characters (see Table 22)

- Pigment
 - Large melanophores medially along length of gut
 - Single irregular row of pigment on each side of ventral midline
 - Ventral pigment begins at anus after yolksac absorption
 - Rays on superior hypural = 4
- Distinguished from *Theragra chalcogramma* at yolksac stage by
- Less lateral pigment within bars
 - Posterior bar longer, extending closer to tail
 - Presence of about 2-6 spots in the ventral caudal region
 - More pigment in snout area and on mouth
- Distinguished from *T. chalcogramma* at later stages by
- Generally more pigmented, especially on head and gut




Figure A, NWAFC original (B. Vinter); B, E-F (F, ventral view), Dunn and Vinter 1984; C-D, Matarese et al. 1981.

MERISTICS

Vertebrae	Total: 53-56-60 Precaudal: 17-20-21 Caudal: 33-37-40
Branchiostegal rays	6-X-8
Caudal fin	22-26, 5+2, 20-24
Pelvic fin	Thoracic R: 6-X-7
Dorsal fin	1st: 9-X-15 2nd: 16-X-21 3rd: 17-X-24
Pectoral fin	R: 19-19-19
Anal fin	1st: 20-X-29 2nd: 18-X-28
Gill rakers	U: 3-X-5 L: 18-X-23

LIFE HISTORY

Range	Cent. California, 34-38°N, to Bering Sea, 54-66°N ^a
Ecology	Epi- and mesobenthal, 0-275 m
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Winter-spring ^b Area: Mode: Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	
No. of oil globules	Probably none
Oil globule diameter	
Yolk	Homogeneous
Envelope	
Hatch size	3 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	<50% SL
Length at flexion	8-15 mm SL
Length at transformation	22-28 mm, pelagic from 28-45 mm SL
Sequence of fin development	Caudal; 1st anal; 2nd anal; 3rd, 2nd, and 1st dorsal (nearly simultaneously); pelvics; pectorals


Pigment

- Presence of bars
- Single row on each side of ventral midline anteriorly and a single row along ventral midline posteriorly
- Large melanophores scattered on ventral surface of gut

Diagnostic characters (see Table 22)

- Pigment
 - Single row on each side of ventral midline anteriorly and a single row along ventral midline posteriorly
 - Large melanophores scattered on ventral surface of gut
 - Anterior placement of bars
 - Rays on superior hypural = 5
- Distinguished from *Gadus macrocephalus* by
- Dorsal pigment separated in specimens <13 mm SL, bars not continuous
- Distinguished from *Theragra chalcogramma* by
- Anterior bar begins closer to anus

^aThe presence of *M. proximus* larvae in the Bering Sea remains a question.^bRichardson 1977



Figures A–E (C, ventral view), Matarese et al. 1981.

MERISTICS

Vertebrae	Total: 48-51-53 Precaudal: 18-X-20 Caudal: 31-X-34
Branchiostegal rays	6-X-8
Caudal fin	21-24, 4+2, 19-22
Pelvic fin	Thoracic R: 6-X-7
Dorsal fin	1st: 10-12-14 2nd: 12-14-18 3rd: 14-17-21
Pectoral fin	R: 17-20-22
Anal fin	1st: 15-18-22 2nd: 15-18-23
Gill rakers^a	U: 5-X-7 L: 25-X-34

LIFE HISTORY

Range	Cent. California, 34-38°N, to Chukchi Sea, north of 66°N
Ecology	Epi-, meso-, and bathypelagic, 0-975 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Feb-Aug ^a Area: Pelagic (50-460 m) ^a Mode: Schools ^b Migration: Bering Sea, offshore to outer and upper slope; ^c Gulf of Alaska to Shelikof Strait ^d
Fecundity	Range/function: 91,633-1,200,000/ $F=0.1719 \times L^{3.6046}$, L=FL cm; ^e 96,216-1,079,540/ $F=1.2604 \times L^{3.2169}$, L=FL cm ^f
Age at maturity	3-4 yr ^g
Longevity	17 yr ^g

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.35-1.45 mm (1.2-1.8)
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Smooth, clear
Hatch size	3-4 mm SL
Incubation time/temp.	15 d/5°C

Pigment

- Late-stage embryo develops bar pattern

Diagnostic characters

- Late-stage embryo with pigment

LARVAE

Preanal length	<50% SL
Length at flexion	10-17 mm SL
Length at transformation	30-40 mm SL
Sequence of fin development	Caudal, 1st anal, 2nd anal, 3rd dorsal, 2nd dorsal, 1st dorsal, pelvics, pectorals


Pigment

- Presence of bars
- A few melanophores scattered on ventral surface of gut
- Pigment along ventral midline and a single row on each side

Diagnostic characters (see Table 22)

- See *Gadus macrocephalus* (p. 194)
- Pigment
 - A few melanophores scattered on ventral surface of gut
 - Pigment along ventral midline and a single row on each side
- Rays on superior hypural = 4

^aHirschberger and Smith 1983^bTakahura 1954^cSerobaba 1968^dDunn and Matarese 1987^eHinckley 1986 (Bering Sea specimens only)^fMiller et al. 1986 (Shelikof Strait specimens only)^gSalveson and Alton 1976a



Figures A–B, NWAFC originals (B. Vinter); C–E (E, ventral view), Dunn and Vinter 1984; F–G, Gorbunova 1954.

MACROURIDAE

Among the most common of all deep-sea bottom-living fishes, grenadiers are found in all the world's oceans. They have long, tapering bodies and elongate dorsal and anal fins continuous with the tail. A chin barbel is usually present. In the North-east Pacific there are ten species in three genera. Adults have been found in water as shallow as 100 m but are more commonly collected at depths greater than 300 m (Fitch and Lavenberg 1968). Little is known of their early life history. Macrourid eggs are generally 1-2 mm, have a single oil globule, and some species have honeycomb ornamentation on the chorion. Larvae are characterized by an elongate tail, lack of caudal fin, and an elongate pectoral fin peduncle. The transition from larva to juvenile is rapid. The most important morphological change is the loss of the pectoral fin peduncle. Other morphological differences occurring at transformation include small changes in head length, mouth orientation changing from oblique to horizontal, snout becoming more distinct, and the stomach (gut) becoming reduced in prominence (decreasing depth of posterior trunk). The occurrence of juveniles of increasing size at increasing depth suggests an ontogenetic migration of subadults to a benthic existence (Stein 1980a). Some species remain pelagic throughout a prolonged juvenile period (Hubbs and Iwamoto 1977).

Table 23
Meristic characters of family Macrouridae.

Taxon	Distribution	Vertebrae		Fins				Gill rakers		Branchiostegals
		Precaudal (Total)	Caudal	First* Dorsal	Second Dorsal	Anal	Pectoral	Pelvic	Upper (Total)	
<i>Albatrossia pectoralis</i>	SSC - Bering Sea	13-14		9-11	126	131	16-21	6-8	5-7 (5-7)	6
<i>Coryphaenoides acrolepis</i>	SSC - Bering Sea	13-16	70	10-13	138-153	123-135	19-22	7-9	5-7 (5-7)	6
<i>Coryphaenoides armatus</i>	SSC - Bering Sea	13-15		10-12		77	18-22	10-12	1 6-9 (7-9)	6
<i>Coryphaenoides cinereus</i>	Oregon - Bering Sea	13-14		12-16			17-23	8-10	1 9-11 (9-12)	6
<i>Coryphaenoides filifer</i>	S. Calif. - Bering Sea	(84)		13-16			18-23	9-10	(8-10)	6
<i>Coryphaenoides leptolepis</i>	SSC - SE Alaska	12		10-12			18-22	9-11	1 8-10 (8-11)	6
<i>Coryphaenoides liocephalus</i>	Brit. Col.	(84)		11	114	113	20	10		6
<i>Coryphaenoides longifilis</i>	Aleutian Is. - Bering Sea	14-15		14-16			15-19	9-10	2-3 12-13	6
<i>Coryphaenoides yaquinae</i>	Cent. Calif. - Oregon			9-12			16-22	8-11	(11-12)	
<i>Nezumia stelgidolepis</i>	SSC - Brit. Col.	13	73-77	10-13			20-26	8-11	(8-12)	7

*First dorsal fin count includes two spinous rays.

Table 24
Morphological characters useful in identifying macrourid larvae likely to occur off the coast of Oregon (Stein 1980a).

	<i>Albatrossia pectoralis</i>	<i>Coryphaenoides</i>				
		<i>C. acrolepis</i>	<i>C. armatus</i>	<i>C. cinereus</i>	<i>C. filifer</i>	<i>C. leptolepis</i>
Larval body pigment	Unknown	Melanophores on dorsum and venter, absent on last 20% of tail and on midline	Unknown	Unknown	Melanophores widely scattered on dorsum around dorsal fin only	Melanophores on trunk and head, closely spaced, not posterior to anal fin ~10th ray
First dorsal fin rays	7-9	9-11	8-10	10-12	11-15	8-10
Pelvic fin rays	6-8	8-9	10-11	8-10	9-10	9-10
Pyloric caeca	12-16	12-14	10-13	5-7	8-12	11
Rostral scutes ^a	Absent	Strong	Absent	Strong	Strong	Absent
Size at which pectoral fin peduncle is reduced	—	9.4-9.8 mm HL ^b	18.2 mm HL	—	14.1-14.6 mm HL	6.2-15.0 mm HL
Precaudal vertebrae	13-14	14-15	13-15	13-14	—	12
Gas glands ^{a,c}	2	4	5-6	4	4	6
Retia ^{a,c}	2	4	5-6	4	4	6

^a Adult/juvenile characters.
^b Since macrourid larvae often have damaged tails, body lengths are generally described by head lengths (HL).
^c The swimbladder in postlarvae and juveniles can be examined by making an incision in the side of the abdominal cavity.

MERISTICS

Vertebrae	Total: 86-86-86 Precaudal: 14-X-15 Caudal: 70-70-70
Branchiostegal rays	6-6-6
Caudal fin	Absent
Pelvic fin	Thoracic R: 8-X-9
Dorsal fin^a	1st: 9-X-11 2nd: 138-X-153
Pectoral fin	R: 19-20-22
Anal fin	R: 123-X-135
Gill rakers	U: X-X-X L: 5-6-7

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Meso- and bathybenthal
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	Ripe eggs 2 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**


Preanal length
Length at flexion
Length at transformation
Sequence of fin development
Pigment
• Melanophores on dorsal and ventral body surface
• Lateral pigment absent on posterior 20% of body

Diagnostic characters (see Table 24)

- Number of first dorsal fin rays (9-11)
- Pelvic fin rays (8-9)
- Number of gas glands (four)
- Presence of rostral scutes in juveniles >16 mm HL

^aRange of counts for the first dorsal does not include "spines."

Ref: Stein 1980a.



Figures A–D, Stein 1980.

MERISTICS

Vertebrae	Total: 84-84-84 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	Absent
Pelvic fin	Thoracic R: 9-9-10
Dorsal fin^a	1st: 11-X-15 2nd: X-X-X
Pectoral fin	R: 18-22-23
Anal fin	R: X-X-X
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Bathybenthal
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE


Preanal length
Length at flexion
Length at transformation
Sequence of fin development
Pigment

- Melanophores widely scattered on upper dorsal body around dorsal fin

Diagnostic characters (see Table 24)

- Number of first dorsal fin rays (11-15)
- Pelvic fin rays (9-10)
- Number of gas glands (four)
- Pigment pattern: Fine pigment spots under first dorsal fin

^aRange of counts for the first dorsal does not include "spines."



Figures A–C, Stein 1980. Figure C is a composite drawn from two specimens. The head has been reconstructed.

MERISTICS

Vertebrae	Total: X-X-X Precaudal: 12-12-12 Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	Absent
Pelvic fin	Thoracic R: 9-X-10
Dorsal fin^a	1st: 8-X-10 2nd: X-X-X
Pectoral fin	R: 18-19-22
Anal fin	R: X-X-X
Gill rakers	U: 1-1-1 L: 8-X-10

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59°N
Ecology	Bathybenthal
ELH pattern	Oviparous, eggs probably pelagic, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length
Length at flexion
Length at transformation
**Sequence of fin
development**
Pigment


- Melanophores on trunk, gut, and head dense, but not present posterior to anal fin (~10th ray)^b

Diagnostic characters (see Table 24)

- Pigment pattern: Pigment to 10th anal fin ray
- Number of gas glands (six)
- Number of first dorsal fin rays (8-10)

^aRange of counts for the first dorsal does not include "spines."

^bAccording to Stein (1980a), this pattern may be similar in *C. armatus*.



Figures A–B, Stein 1980.

MERISTICS

Vertebrae	Total: 86-X-90 Precaudal: 13-13-13 Caudal: 73-X-77
Branchiostegal rays	7-7-7
Caudal fin	Absent
Pelvic fin	Thoracic R: 8-X-11
Dorsal fin	S: 10-X-13 R: X-X-X
Pectoral fin	R: 20-X-26
Anal fin	R: X-X-X
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epi- and mesobenthic
ELH pattern	Oviparous, eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE** - Genus

Preanal length <30% SL
Length at flexion
Length at transformation
Sequence of fin development
Pigment

- Frontal patch
- First dorsal fin patch
- Heavy over entire gut
- Ventral midline series and double lateral spots approximately first half of postanal body

Diagnostic characters

- Lateral pigment on postanal body appears to be distinctive perhaps at generic level, but whether it appears in other macrourine larvae cannot be determined at this time




Figure A, Fahay and Markle 1984. North Atlantic specimen tentatively identified as *Nezumia*.



Ophidiiformes

The ophidiiforms occupy mostly benthic habitats ranging from the shallow tropics to abyssal depths and subarctic locations. Adults have long tapering bodies, with or without a caudal fin, and long dorsal and anal fins. The order comprises 4 families, 86 genera, and 294 species (J. Nelson 1984). The two suborders are defined according to mode of reproduction: Ophidioidei (Ophidiidae) are oviparous and Bythitoidei (Aphyonidae and Bythitidae) are viviparous (Cohen and Nielsen 1978). Ophidiiform eggs of oviparous forms are pelagic, may have one oil globule, and are spherical or ellipsoidal (Gordon et al. 1984). Larvae are not well known; of the eight species found in the study area only two larval series have been identified and are presented here. A third series of an unidentified ophiidid has been tentatively identified as *Spectrunculus grandis*.

Families in study area: **Ophidiidae**
Aphyonidae
Bythitidae

OPHIDIIDAE

Fishes of this group are found primarily in tropical and temperate waters, although a few occur in subarctic locations. In the northeastern Pacific there are six species within five genera in the family. Larvae are known only for *Chilara taylori*, but a second series has been tentatively identified as *Spectrunculus grandis*. These larvae occur in samples collected off northern California, Oregon, and Washington. Since meristic characters, especially vertebral counts, are unavailable for several species in the area, only a tentative identification can be made. Available meristics, however, match those of *Spectrunculus grandis*. These larvae resemble those of the pleuronectid, *Embassichthys bathybius*, and have been routinely mixed with them in samples. They can be distinguished from *E. bathybius* (p. 580) larvae by number of myomeres (>75 vs. 65), less finfold pigment, and a less pronounced loop in the gut.

Table 25
Meristic characters of family Ophidiidae.


Taxon	Distribution	Vertebrae		Fins				Gill rakers		Branchiostegals
		Precaudal	Caudal	Dorsal	Anal	Pectoral	Pelvic	Upper	Lower	
<i>Bassozetus</i> sp.	Oregon									
<i>Chilara taylori</i>	SSC - Oregon	18-19	68-72	198-216	156-170		I,1	1-4	5-9	7
<i>Dicrolene filamentosa</i>	SSC - Oregon			100-104	84-90	23-29	I,2	5	17	8
<i>Holcomycteronus profundissimus</i>	Oregon	18-21		107-118	80-95	15-17	I,2			8
<i>Spectrunculus grandis</i> *	Cent. Calif. - Gulf of Alaska	23	56	103-140	73-106	26-30	I,2	3-4	8-9	8
<i>Spectrunculus radcliffei</i>	Gulf of Alaska									

*Aboussouan and Rasonarivo (1986) described a larva as *S. grandis* from the Indian Ocean. The counts are the following:

precaudal vertebrae	18-19
caudal vertebrae	53
dorsal fin	118
anal fin	96
pectoral fin	23/24
pelvic fin	I,2
caudal fin	4+4/5


Ophidiidae (tentatively *Spectrunculus grandis*)

A



15.6 mm SL

B



29.8 mm SL

Figures A–B, NWAFC originals (B. Vinter).

MERISTICS

Vertebrae	Total: 86-88-91 Precaudal: 18-18-19 Caudal: 68-70-72
Branchiostegal rays	7-7-7
Caudal fin	X, 4+5, X
Pelvic fin	Jugular
	S: 1-1-1 R: 1-1-1
Dorsal fin	R: 198-X-216
Pectoral fin	R: 22-X-25
Anal fin	R: 156-X-170
Gill rakers	U: 1-X-4 L: 5-X-9

LIFE HISTORY

Range ^a	South of southern California to Oregon, 42-46°N
Ecology	Epi- and mesobenthal, 1-244 m
ELH pattern	Oviparous, eggs unknown, pelagic larvae
Spawning	Season: Larvae collected in fall (Baja California) ^b Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE

Preanal length <50% SL
Length at flexion
Length at transformation 22-30 mm SL
Sequence of fin development
Pigment

- Caudal fin pigment
- Ventral (double series) with development, extends along base of brain and onto snout
- Ventral gut


Diagnostic characters

- Gut loops form at 14 mm SL
- Distinctive pigment pattern
- High number of myomeres (86-91)

^aCenter of distribution is Pt. Eugenia, Baja California.

^bAmbrose et al. 1983

Ref: Ambrose et al. 1983.



Figures A–E (C, ventral view), Ambrose et al. 1983.

Fishes of this group are found in tropical and temperate waters, freshwater, and at abyssal depths. The families Aphyonidae and Bythitidae, which form the suborder, are represented in the Northeast Pacific by four species within four genera. Only larvae of *Brosomphycis marginata* are known. All bythitoid males possess an intromittent organ, some with pseudoclaspers.

Table 26
Meristic characters of suborder Bythitoidei.

Taxon	Distribution	Vertebrae		Fins				Gill rakers		Branchiostegals
		Precaudal (Total)	Caudal	Dorsal	Anal	Pectoral	Pelvic	Upper (Total)	Lower	
Aphyonidae										
<i>Barathronus pacificus</i>	N. Calif. - Oregon	37-38	46-51	71-75	62-69	25-26	I,1	5-7	26-28	
<i>Sciadonus pedicellaris</i>	SSC - Oregon	43-47	36-39	91-93	42-47	11-14	I,1	(14-15)		9-10
Bythitidae										
<i>Brosomphycis marginata</i>	SSC - SE Alaska	16-17	47-49	99-110	72-81	20-26	I,2			7
<i>Cataetyx rubrirostris</i>	S. Calif. - Oregon	(60-62)		102-109	76-82		I,1	3		8

MERISTICS

Vertebrae	Total: 63-64-65 Precaudal: 16-16-17 Caudal: 47-48-49
Branchiostegal rays	7-7-7
Caudal fin	
Pelvic fin	Jugular S: 1-1-1 R: 2-2-2
Dorsal fin	R: 99-101-110
Pectoral fin	R: 20-X-26
Anal fin	R: 72-75-81
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59°N
Ecology	Epi- and mesobenthic, 3-256 m
ELH pattern	Probably ovoviviparous, pelagic larvae
Spawning	Season: Spring ^a Area: Mode: Migration:
Fecundity	Range/function: 12,000-30,000 ^a
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	9.0 mm NL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE^b**

Preanal length	<50% SL
Length at flexion	>13.5-18.0 mm SL
Length at transformation	
Sequence of fin development	
Pigment	<ul style="list-style-type: none"> Newly hatched larvae have pigment on lower jaw Alternating dorsal/ventral spots Above/below anterior portion of tail tip (more pronounced in specimens <12.5 mm SL) Swimbladder and on gut

Diagnostic characters

- Pigment pattern: Dorsal/ventral spots

^aHart 1973^bA figure of a postflexion bythitid from the Atlantic Ocean is included for comparison.





Figure A, NWAFC original (B. Vinter); B–C, Gordon et al. 1984 (C, North Atlantic specimen).



Lophiiformes

The order Lophiiformes includes some of the most unusual fishes known. Adults exhibit several forms: Dorsoventrally flattened and ventrally compressed forms found in shallow-water benthic habitat, and globose, flabby-bodied forms found at meso- and bathypelagic depths. Worldwide there are 18 families, 63 genera, and 262 species (Pietsch 1984). All families are characterized by having a luring apparatus (illicium) bearing a terminal bait (esca) originating from three modified anterior dorsal spines and a narrow tube-like gill opening located near the pectoral fin. Extreme sexual dimorphism is common among the fishes within the suborder Ceratioidei. Males are one-third the size of females or smaller and sometimes are permanently attached (parasitic) to them. Eggs of most lophiiforms are spawned in gelatinous “veils” which may be up to 1.5 m wide and 12 m long (Pietsch 1984). Several species have eggs attached to the adult (Pietsch and Grobecker 1987). Larvae are pelagic and many have large heads and an envelope of highly inflated skin. Neither eggs nor larvae have been collected in the study area.

Families in study area: **Ceratiidae**
Oneirodidae

CERATIOIDEI

The suborder Ceratioidei is represented within the study area by members of the families Oneirodidae and Ceratiidae. Ceratioids differ from other members of the Lophiiformes in being meso- and bathypelagic, lacking pelvic fins (except larval Caulophrynidae), and in having extreme sexual dimorphism.

Oneirodidae

Found worldwide, oneirodids (dreamers) are represented by five species within three genera (*Oneirodes*, *Chaenophryne*, and *Bertella*) in the Northeast Pacific. Reproduction is accomplished by facultative or non-parasitic attachment of the male onto the body of the female (Pietsch 1976). Eggs and larvae are pelagic. Eggs are probably released in gelatinous veils which might break up in plankton nets. Eggs are presumably small, generally 0.5-0.8 mm, and hatching occurs between 2.5 and 3.5 mm SL (Bertelsen 1984). Larvae have a transparent envelope of gelatin, under colorless skin, which may serve as an aid to flotation (Idyll 1964, Pietsch 1984).

According to Bertelsen (1984), larval characters for oneirodids include the following:

- Presence in larval males of a rudiment of illical bone
- Moderately elongate body shape (body depth up to 80-90% in most ceratioids but less pronounced in oneirodids)
- Larvae surrounded by inflated transparent skin (similar to other ceratioids)
- Head length usually about 45% SL
- Pigment: Ceratioids possess four main pigment areas (peritoneal, dorsal, caudal, and opercular). In oneirodids, the opercular pigment is dense and occurs in different patterns

At metamorphosis (usually between 8 and 10 mm SL), larvae descend into deeper waters.

Ceratiidae

Recently, two ceratiids (seadevils) have been identified from the Bering Sea, *Ceratias holboelli* and *Ceratias* sp. (Pietsch 1986). Larval ceratiids are distinguished from other ceratioid families in having the following combination of characters (Pietsch 1986):

- Body “hump-backed,” mouth subvertical
- Female with caruncles on dorsal surface of trunk
- 4-5 dorsal fin rays, 4 anal fin rays
- Pectoral fins not reaching beyond dorsal and anal fins
- Pelvic fins absent

Table 27
Meristic characters of Ceratioidei.

Taxon	Distribution	Vertebrae		Fins			Branchiostegals
		Precaudal (Total)	Caudal	Dorsal	Anal	Pectoral	
Ceratiidae							
<i>Ceratias holboelli</i>	Bering Sea			4-5	4		
<i>Ceratias</i> sp.	Bering Sea			4-5	4		
Oneirodidae							
<i>Bertella idiomorpha</i>	SSC - Gulf of Alaska	5	15	5-6	4-5	17-21	6
<i>Chaenophryne longiceps</i>	SSC - Oregon		(21)	6-7	5-6	17-22	6
<i>Chaenophryne melanorhabdus</i>	SSC - Brit. Col.	9	10	6-8	5-6	16-17	6
<i>Oneirodes bulbosus</i>	Oregon - Bering Sea	4	16	6-7	4	15-18	6
<i>Oneirodes thompsoni</i>	N. Calif. - Bering Sea	4	16	5-6	4	14-17	6

MERISTICS

Vertebrae	Total: 21-21-21 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Absent
Dorsal fin	R: 6-7-7
Pectoral fin	R: 17-19-22
Anal fin	R: 5-5-6
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Mesopelagic, 200-1000 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Males attach to body of female ^a Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE^b**

Preanal length
Length at flexion
Length at transformation
Sequence of fin development
Pigment


- Pigment appears in all four pigment areas
 - Peritoneal
 - Anterior body, begins as a small dorsal patch
 - Opercular
 - Caudal, initially a few spots

Diagnostic characters

- Generally more lightly pigmented than *Oneiroides* spp.

^aPietsch 1976

^bBecause of the possibility of involvement of other species in the description, and geographic variation, specimens in our area could differ.



Figures A–E, Bertelsen 1951 (North Atlantic specimens).

MERISTICS

Vertebrae	Total: 20-20-20 Precaudal: 4-4-4 Caudal: 16-16-16
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Absent
Dorsal fin	R: 6-X-7
Pectoral fin	R: 15-16-18
Anal fin	R: 4-4-4
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Oregon, 42-46°N, to Bering Sea, 54-66°N
Ecology	Mesopelagic, 200-1000 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Males attach to body of female ^a Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment


Diagnostic characters**LARVAE^b**

Preanal length
Length at flexion
Length at transformation
Sequence of fin
development
Pigment - Species group


- Pigment in three out of the four main pigment areas
 - Peritoneal
 - Anterior body
 - Opercular
- Caudal pigment is light and appears late in development

Diagnostic characters^aPietsch 1976^bFigures of *Oneirodes eschrichti* group (Bertelsen 1951) are presented for comparison only. Because of the possibility of involvement of other species in the description, and geographic variation, specimens of *Oneirodes* in our area could differ.

Ref: Bertelsen 1984.



Figures A–E, Bertelsen 1951 (North Atlantic specimens from the *O. eschrichti* group).



Gobiesociformes

The gobiesociforms (clingfishes) are mostly small, inshore bottom-dwelling species occurring in tropical and temperate seas. Noted primarily for having the pelvic fins modified into a sucking disc, the order has 2 families, 36 genera, and 114 species (J. Nelson 1984). All but one genus and four species are members of Gobiesocidae. Eggs are demersal, attached to substrate or kelp, ovate to ellipsoidal, and 0.7-1.9 mm. Oil globules (1-100) coalesce to one during development (Allen 1984). Larvae are well developed at hatching and may have a fully formed pelvic disc at this time (Marliave 1975a). Most are heavily pigmented and have long guts (50-70% SL) (Allen 1984).

Family in study area: **Gobiesocidae**

MERISTICS

Vertebrae	Total: 32-33-34 Precaudal: 12-14-14 Caudal: 19-19-20
Branchiostegal rays	6-6-6
Caudal fin	Total rays=11-13
Pelvic fin	Disc S: 1-1-1 R: 4-4-4
Dorsal fin	R: 13-16-16
Pectoral fin	R: 21-22-23
Anal fin	R: 13-14-15
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59°N
Ecology	Intertidal, nearshore, 0-8 m
ELH pattern	Oviparous; demersal, attached, guarded eggs; pelagic larvae
Spawning	Season: Winter-spring (British Columbia) ^a Area: Demersal, on underside of rocks, usually intertidal ^a Mode: Polygamous males guard eggs (laid in monolayer) ^a Migration:
Fecundity	Range/function: 194-382 per female ^b
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.68-1.92 mm (1.78)
No. of oil globules	
Oil globule diameter	
Yolk	Bilobed in embryo and yolksac larva
Envelope	
Hatch size	5.7 mm SL, yolksac present until 6.2 mm SL
Incubation time/temp.	30 d/9-11°C
Pigment	

Diagnostic characters

- Adhesive, hemispherical, flat at point of attachment

LARVAE

Head length	21-35% SL; gut length (preanal minus head length) 25% SL
Length at flexion	7 mm SL
Length at transformation	10-13 mm SL
Sequence of fin development	Pectoral fin buds form at 7.0 mm SL; pelvic disc is formed by 7.3 mm SL and fully functioning by 9.1 mm SL

Pigment

- Dorsally on gut
- With development, on snout and dorsal region of head






Diagnostic characters

- Large melanophores on dorsal surface of gut, lack of pigment on lateral surface
- Forms pelvic disc
- Number of myomeres (32-34)

^aMarliave 1975a

^bJohnson 1970

Ref: Allen 1984, Allen and Ilg 1983.



Figures A, C–E, Allen and Ilg 1983; B, NWAFC original (B. Vinter).

MERISTICS

Vertebrae	Total: 35-X-36 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	5-X-7
Caudal fin	Total rays=8
Pelvic fin	Disc S: 1-1-1 R: 4-4-4
Dorsal fin	R: 6-7-8
Pectoral fin	R: 14-15-17
Anal fin	R: 6-7-8
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Intertidal, nearshore
ELH pattern	Oviparous; demersal, attached eggs; pelagic larvae
Spawning	Season: Area: Kelp beds, on blades ^a Mode: Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.3 mm
No. of oil globules	One
Oil globule diameter	
Yolk	
Envelope	
Hatch size	<4 mm SL; yolk sac present at 4 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	~50% SL
Length at flexion	
Length at transformation	
Sequence of fin development	
Pigment (based on figure only)	<ul style="list-style-type: none"> • Lateral melanophores (40-50) • No postanal ventral melanophores • No hypural pigment

Diagnostic characters

- Pelvic disc at hatching
- Number of myomeres (35-36)

Note: Marliave (1975a) reared *Rimicola muscarum* in the laboratory. Eggs (1.3 mm in diameter) had a single oil globule, and late-stage embryos developed a pelvic disc. Hatching occurred when embryos reached 4.0 mm SL. He described them as identical to *Gobiesox maeandricus* larvae except that they were markedly smaller, without nasal pigment, and with the pelvic disc at hatching. According to Marliave (Vancouver Public Aquarium, P.O. Box 3232, Vancouver, B.C., Canada V6B 3X8, pers. commun., 16 Oct. 1986), *R. muscarum* probably lacks a pelagic stage and the specimen illustrated here from Allen (1979) is misidentified.

^a Allen 1984

Ref: Allen 1979, Marliave 1975a.

A





Figure A, Allen 1979.



Atherinomorpha: Beloniformes
Atheriniformes

The Beloniformes and Atheriniformes together comprise a group of fishes found in freshwater and marine habitats within tropical and temperate areas. The Beloniformes (not always accorded ordinal status [J. Nelson 1984]), consists of 5 families, 37 genera, and 180 species (Collette et al. 1984a). Marine forms of this order are mostly epipelagic, the best known being exocoetids, or flying fish. There are 6 families (White et al. 1984), 49 genera, and 235 species of atheriniforms (J. Nelson 1984). The two orders belong to the superorder Atherinomorpha and share the development of large demersal eggs with filaments and oil globules that coalesce at the vegetal pole (Collette 1984). Filaments may be short or long, grouped or evenly scattered, and adhesive or non-adhesive. Beloniform larvae are well formed at hatching and many have a preanal finfold and beak-like jaws (Collette et al. 1984a). Atheriniform larvae have direct development (i.e., no specialized larval or juvenile stages), a preanal finfold, and a single row of melanophores on the dorsal midline (White et al. 1984).

Families in study area: **Scomberesocidae**
Atherinidae

MERISTICS

Vertebrae	Total: 62-65-69 Precaudal: 37-38-40 Caudal: 24-27-29 ^a
Branchiostegal rays	14-X-15
Caudal fin	X, 7+8, X
Pelvic fin	Abdominal R: 6-6-6
Dorsal fin	R: 14-X-18 ^a
Pectoral fin	R: 12-13-15
Anal fin	R: 18-X-21 ^a
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epipelagic, 0-200 m
ELH pattern	Oviparous, eggs attach to flotsam (kelp) with adhesive filaments, pelagic larvae
Spawning	Season: Winter-fall; ^b peaks Feb-July ^c Area: Mode: Schools ^b Migration:
Fecundity	Range/function: ~1800 during each of 6-7 spawnings ^b
Age at first maturity	2 yr ^b
Longevity	5 yr ^b

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.58 (1.68 mm) × 2.13 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Adhesive filaments, 12-15 in polar cluster and 1 lateral
Hatch size	6.0-8.5 mm SL (family)
Incubation time/temp	
Pigment	

Diagnostic characters

- Adhesive filaments in polar cluster
- Single lateral filament
- Slightly ovoid

LARVAE

Preanal length	65% SL
Length at flexion	
Length at transformation	
Sequence of fin development	Caudal, dorsal and anal, pectorals, pelvics (caudal forms at hatching)

Pigment

- Dense body pigment

Diagnostic characters (see Table 3)

- Persistent preanal finfold
- Unpigmented area on ventrolateral region of caudal peduncle
- At 20-40 mm SL, upper and lower jaw are slightly elongate but do not form prominent beak; with growth a slight beak develops

^aCollette et al. 1984a^bFitch and Lavenberg 1971^cWang 1981




Figure A, Matarese and Sandknop 1984; B, Collette et al. 1984a; C–D, NWAFC originals (B. Vinter); E, Moser 1981; F, Uchida et al. 1958.

MERISTICS^a

Vertebrae	Total: 44-48-52 Precaudal: 32-34-37 Caudal: 11-13-15
Branchiostegal rays	5-X-6
Caudal fin	
Pelvic fin	Abdominal S: 1-1-1 R: 5-5-5
Dorsal fin	S: 6-X-10 R: 8-X-14
Pectoral fin	R: 13-13-13
Anal fin	S: 1-1-1 R: 19-X-25
Gill rakers	U: 4-X-8 L: 21-X-34

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Nearshore shelf pelagic
ELH pattern	Oviparous, eggs with adhesive filaments, pelagic larvae
Spawning	Season: May-July ^b Area: Mudflats, eelgrass ^b Mode: Schools ^b Migration: To mudflats in sloughs ^a
Fecundity	Range/function:
Age at first maturity	2-3 yr ^c
Longevity	6-7 yr ^b

^a Meristic information from White et al. (1984) is very different from the information in our database. Due to confusion in the literature over the definition of spines and rays, only total elements are reported by White et al. (1984) and they are as follows:

Vertebrae	43-49	Pectoral fin	12-15
First dorsal fin	3-7	Anal fin	9-14
Second dorsal fin	10-14	Gill rakers	14-27

^b Hart 1973

^c Fitch and Lavenberg 1975

^d Additional unpubl. data provided by W. Watson, H.J. Walker, and R. Davis (W. Watson, Marine Ecological Consultants, 531 Encinitas Blvd., Suite 110, Encinitas, CA 92024, pers. commun., 3 Nov. 1986).

^e Wang 1981

^f According to White et al. (1984), two egg types are reported for the species, one with 40-78 filaments attached at both ends and the other with about 6 filaments attached only at one end.

Ref: Wang 1981, White et al. 1984.

EARLY LIFE HISTORY DESCRIPTION^d

EGGS

Diameter	1.5-1.7 mm ^e
No. of oil globules	Initially may have one to many, but they usually coalesce to one

Oil globule diameter

Yolk	Amber, granular
Envelope	6-7 filaments ^f
Hatch size	4.3-4.9 mm SL; yolk absorbed by 7 mm SL
Incubation time/temp.	9 d/15-18°C

Pigment

- Dorsal and lateral spots on head, few anterior spots on yolk sac

Diagnostic characters

- Narrow perivitelline space

LARVAE

Preanal length	30-40% SL
Length at flexion	7.7-10.5 mm SL
Length at transformation	15 mm SL
Sequence of fin development	Caudal, pectorals, anal rays and 2nd dorsal, pelvics, 1st dorsal and anal spines

Pigment

- Dorsal midline from snout to caudal peduncle
- Mediolateral beginning above gut
- Dorsal and ventral surface of gut
- Ventrally on tail

Diagnostic characters

Distinguished from *Atherinopsis californiensis* by

- Presence of melanophores along ventral body midline and ventral gut




Figure A, White et al. 1984; B–E (D, ventral view), Watson and McGowen, unpubl.

MERISTICS^a

Vertebrae	Total: 50-51-52
	Precaudal: 9-12-14
	Caudal: 37-39-41
Branchiostegal rays	5-X-6
Caudal fin	
Pelvic fin	Abdominal
	S: 1-1-1 R: 6-6-6
Dorsal fin	S: 6-X-10 R: 11-X-14
Pectoral fin	R: 15-15-15
Anal fin	S: 1-1-1 R: 21-X-26
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Nearshore shelf pelagic
ELH pattern	Oviparous, eggs with filaments in a mass or attached to substrate, pelagic larvae
Spawning	Season: Fall-spring ^b Area: On algae, eelgrass ^b Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION^c**EGGS**

Diameter	1.9-2.5 mm ^b
No. of oil globules	Many (23-44) which coalesce to one or few
Oil globule diameter	0.4-0.6 mm (consolidated)
Yolk	Yellow-orange, granular
Envelope	13-19 filaments scattered on surface
Hatch size	7.5-8.6 mm SL; yolk absorbed by 10 mm SL
Incubation time/temp.	17-19 d/12-16°C
Pigment	<ul style="list-style-type: none"> • Dorsal spots along nape

Diagnostic characters

- Narrow perivitelline space
- 1-2 cm filaments

LARVAE

Preanal length	<50% SL
Length at flexion	8.8 to ≤11.5 mm SL
Length at transformation	18.1-19.5 mm SL
Sequence of fin development	Caudal, pectorals, anal rays and 2nd dorsal, pelvics, 1st dorsal and anal spines

Pigment

- Dorsal midline from snout to caudal peduncle, sometimes with a break in nape area
- Mediolateral usually begins posterior to gut during preflexion stage

Diagnostic characters

Distinguished from *Atherinopsis affinis* by

- First 5-6 postanal myomeres always lack ventral pigment through the larval and early juvenile stages
- Lack of ventral midline melanophores until midflexion
- Mediolateral pigment originates further posterior

^aMeristics from White et al. (1984) are as follows:

Vertebrae	46-53
First dorsal fin	4-9
Second dorsal fin	10-15
Pectoral fin	14-17
Anal fin	20-29
Gill rakers	18-44

^bWang 1981

^cAdditional unpubl. data provided by W. Watson, H.J. Walker, and R. Davis (W. Watson, Marine Ecological Consultants, 531 Encinitas Blvd., Suite 110, Encinitas, CA 92024, pers. commun., 3 Nov. 1986).

Ref: Wang 1981, White et al. 1984.





Figure A, White et al. 1984; B–E (C, ventral view), Watson and McGowen, unpubl.



Lampriformes

Fishes of the order Lampriformes are extremely diverse. Occupying meso- and epipelagic habitats, some are deep-bodied while others are long and ribbon-like. Extreme specialization and unique body shapes have caused conflicting proposals for evolutionary relationships within the order (Olney 1984). According to Olney (1984), there are 7 families, 12 genera, and 21 families worldwide (not found in polar seas). Lampriforms share the unique arrangement of protrusible premaxilla and maxilla, or specializations thereof. Early-life-history information is available for only four genera. Eggs are large (1.7-4.0 mm) with thick resilient chorions, pelagic, and may be shaded with amber, pink, or red hues. Advanced stages of eggs of some forms are easily recognized due to precocious development of anterior dorsal and pelvic rays and distinctive pigment patterns. Newly hatched lampriforms are identified by their well-developed protrusible jaws and elongate anterior dorsal and pelvic fin elements which are often ornamented with highly pigmented serial or terminal swellings (Olney 1984).

Families in study area: **Lampridae**
Trachipteridae

MERISTICS^a

Vertebrae	Total: 43-X-46 Precaudal: 19-X-21 Caudal: 23-24-25
Branchiostegal rays	6-X-7
Caudal fin	Total rays = 30-32
Pelvic fin	Abdominal R: 13-X-17
Dorsal fin	R: 48-X-52
Pectoral fin	R: 21-X-24
Anal fin	R: 33-X-42
Gill rakers	U: 2-X-3 L: 13-X-14

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesopelagic
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Spring ^b Area: Mode: Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS** - Family

Diameter	Large, 1.7-4.0 mm
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous in known forms
Envelope	Thick, resilient
Hatch size	
Incubation time/temp.	~3 weeks
Pigment	

Diagnostic characters

- Precocious development of fins
- Ovarian eggs have thick chorion with amber tint

LARVAE

Preanal length	<50% SL, increasing with development to ≥50% SL
-----------------------	---

Length at flexion**Length at transformation**

Sequence of fin development	Anterior dorsal and pelvics precocious; dorsal, anal, and caudal; pectorals
------------------------------------	---

Pigment

- Initially on crown and dorsal surface of gut; with development, increases along entire body except for tail region and above anal fin

Diagnostic characters

Distinguished from *Trachipterus altivelis* (p. 246) by


- Lack of ornamentation on dorsal and pelvic fins (when fins are intact)
- Rapid change in body form from slender at hatching to deep-bodied by 10.6 mm SL

Lampriform characters

- Well-developed protrusible jaws
- Differentiated guts with open lumen and little yolk
- Elongate anterior dorsal elements (may be ornamented) and well-developed pelvic elements (may be ornamented with lengths to 40-60% NL)

^aData from Olney (1984) in part, only total elements reported.

^bFitch and Lavenberg 1971



Figures A–C, Olney 1984.

MERISTICS

Vertebrae	Total: 90-92-94 Precaudal: 35-35-39 Caudal: 53-59-59
Branchiostegal rays	6-X-7
Caudal fin	
Pelvic fin	Thoracic R: 6-X-7
Dorsal fin	R: 160-X-191
Pectoral fin	R: 10-X-11
Anal fin	Absent
Gill rakers	U: 3-X-5 L: 9-X-11

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesopelagic, 0-900 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	2.6-3.7 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Clear, smooth, thick
Hatch size	
Incubation time/temp.	
Pigment	<ul style="list-style-type: none"> • Head and gut • Finfold • Ornamentation on elongate rays

Diagnostic characters

- Precocious development of elongate rays in anterior dorsal fin and pelvic fins

LARVAE

Preanal length	<50% SL, with development increasing to >50% SL
Length at flexion	
Length at transformation	
Sequence of fin development	Anterior dorsal and pelvics precocious, posterior dorsal, caudal, pectorals
Pigment	<ul style="list-style-type: none"> • Initially only on head and anterior/dorsal gut; with development pigment appears on lateral surface of gut, along body over gut, and in a series above notochord along 3/4 BL • Several spots develop above and below notochord in caudal region

Diagnostic characters

- See *Lampris guttatus* (p. 244)





Figure A, Matarese and Sandknop 1984; B–C, NWAFC originals (B. Vinter).



Beryciformes

Fishes of the order Beryciformes are cosmopolitan in distribution. The composition of the order is subject to much variation, but according to Keene and Tighe (1984), it consists of 16 families, 42 genera, and 155 species. Although several characters define the beryciforms (e.g., a high number of pelvic and caudal fin rays and the presence of dorsal, anal, and pelvic fin spines), none are unique to the order. No information is available on eggs, and other early-life-history stages are known for only six families. In the Northeast Pacific, larvae are known for only two families, Anoplogastridae and Melamphaidae. Anoplogastrids have large heads with short deep bodies, and well-developed preopercular, rostral, and cranial spines. Larvae of the best known family, Melamphaidae, are longer-bodied and more slender with spination generally restricted to the preopercle.

A summary of meristic characters is provided for members of the suborder Cetomimoidei (Barbourisiidae, Cetomimidae, Rondeletiidae), as larvae are unknown.

Families in study area: **Anoplogastridae**
Melamphaidae
Barbourisiidae
Cetomimidae
Rondeletiidae

MERISTICS

Vertebrae	Total: 25-X-28 Precaudal: 12-12-12 Caudal: 16-16-16
Branchiostegal rays	8-X-9
Caudal fin	X, 17, X
Pelvic fin	Thoracic S: 1-1-1 R: 6-6-6
Dorsal fin	R: 17-18-19
Pectoral fin	R: 14-15-16
Anal fin	R: 8-X-9
Gill rakers	U: 7-X-11 L: 7-X-11

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Epi- and mesopelagic, below 610 m
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Summer (California) ^a Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE

Preanal length
Length at flexion 4.5-6.0 mm SL
Length at transformation
Sequence of fin development Pelvic fins last to develop

Pigment

- At 6 mm SL, lateral surface of body to caudal peduncle, gut, and pectoral fin base pigmented
- Pigment increases with development

Diagnostic characters

- Spination
 - Serrate frontal ridge terminates in a short stout supraocular spine
 - Long serrate parietal spine
 - Long serrate preopercular spine directed posteroventrad

Juveniles

- Increase in dark pigment
- Reduction of spines

^aFitch and Lavenberg 1968

Ref: Keene and Tighe 1984.





Figure A, Keene and Tighe 1984.

MELAMPHAIDAE

Melamphoids, or bigscales, are small (15 cm), darkly colored, bathypelagic fish found in most oceans. They are distinguished by exceptionally thin skull bones, sometimes with ridges or crests. They are represented in the study area by five species within four genera: *Melamphaes*, *Poromitra*, *Scopeloberyx*, and *Scopelogadus*. A complete descriptive series is known only for *Melamphaes lugubris*. Generally, small melamphaid larvae (2-10 mm) are relatively elongate and slender as compared with later stages, and possess early and rapidly developing pelvic fins which are long and darkly pigmented. *Melamphaes*, *Scopeloberyx*, and *Scopelogadus* larvae usually have two spots near the posterior end of the dorsal and anal fin margin; with development these spots spread both anteriorly and posteriorly along the dorsal and ventral midlines (Keene and Tighe 1984). Other pigment occurs on the cranium, peritoneum, and along the caudal peduncle. Although usually damaged, the second or third dorsal fin ray is elongate in larvae of these three genera. Larger larvae (5-10 to 20 mm) can generally be distinguished by body shape, presence of preopercular spines (e.g., *Poromitra*), and meristic characters. The following generic meristic characters are based on Ebeling (1962) and E.H. Ahlstrom notes, in part.


Genus	Vertebrae	Fins			
		Dorsal	Anal	Pectoral	Pelvic
<i>Melamphaes</i>	27-31	III, 13-16	I,7-9	14-17	I,7
<i>Poromitra</i>	24-29	III, 9-14	I,8-11	13-15	I,7-8
<i>Scopeloberyx</i>	23-27	II-III, 10-13	I,7-9	12-14	I,7-8
<i>Scopelogadus</i>	24-26	II, 10-12	I,7-9	13-15	I,7-8

A *Melamphaes lugubris* MELAMPHAES




6.2 mm SL

B *Melamphaes lugubris*




8.3 mm SL

C *Poromitra crassiceps* POROMITRA




7.9 mm SL

D *Poromitra* sp.




13.5 mm SL

E *Scopeloberyx robustus* SCOPELOBERYX




13.0 mm SL

F *Scopeloberyx* sp.



6.5 mm SL

G *Scopelogadus bispinosus* SCOPELOGADUS



8.0 mm SL

Figures A–G, Keene and Tighe 1984 (B–G, North Atlantic specimens).

MERISTICS

Vertebrae	Total: 28-X-31 Precaudal: 11-12-12 Caudal: 16-X-19
Branchiostegal rays	8-8-8
Caudal fin	X, 10+9, X
Pelvic fin	Thoracic S: 1-1-1 R: 7-7-7
Dorsal fin	S: 3-3-3 R: 14-15-16
Pectoral fin	R: 15-16-17
Anal fin	S: 1-1-1 R: 7-8-9
Gill rakers	U: 5-X-6 L: 15-X-18

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Mesopelagic, 200-1000 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS** - Family

Diameter	0.81-0.96 mm SL
No. of oil globules	One to many
Oil globule diameter	
Yolk	Segmented
Envelope	Clear, spherical
Hatch size	Probably <2 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	45-55% SL
Length at flexion	~6 mm SL
Length at transformation	
Sequence of fin development	Pelvics, dorsal and caudal, anal, pectorals
Pigment	<ul style="list-style-type: none"> • Dorsally on top of head • Peritoneum • Postanal: Two spots near posterior of dorsal and anal anlagen which spread anteriorly and posteriorly along dorsal and ventral midline • Spot on caudal peduncle

Diagnostic characters

- Pigment (see above): Patches, bands
 - 2nd or 3rd dorsal ray elongate (until 5-10 mm SL)
 - Pelvics: (2-10 mm SL) develop rapidly and are long, fragile, and pigmented
- Distinguished from *Sebastolobus* spp. (p. 336) at sizes <5 mm SL by
- Postanal pigment band more anterior, beginning at myomere 15
 - Precocious development of pelvic fin

M. parvus larvae are unknown. The following meristic information may aid in identification.

Total vertebrae	27-29
Precaudal vertebrae	11-12
Caudal vertebrae	16-17
Dorsal fin rays	III, 13-15
Anal fin rays	I, 7-9
Pectoral fin rays	14-15
Pelvic fin rays	I, 7




Figure A, NWAFC original (B. Vinter); B-E, Keene and Tighe 1984.


CETOMIMOIDEI

Fishes of the suborder Cetomimoidei are bathypelagic with whale-shaped bodies, large mouths, highly distensible stomachs, and luminous tissue on the body. Eyes may be reduced or rudimentary. Whalefishes are divided into three families: Cetomimidae, Rondeletiidae, and Barbourisiidae. Worldwide in distribution, the suborder consists of 8 genera and 18 species. Of these, five species in five genera have been collected in the northeastern Pacific. Adults are taken by deep tows generally in excess of 400 m, and then only rarely. No information on life history, eggs, or larvae is available (Keene and Tighe 1984).

Table 28
Meristic characters of suborder Cetomimoidei.

Taxon	Distribution	Vertebrae		Fins				Gill rakers		Branchiostegals
		Precaudal (Total)	Caudal	Dorsal	Anal	Pectoral	Pelvic	Upper	Lower	
Barbourisiidae			(42-44)							
<i>Barbourisia rufa</i>	Cent. Calif. - Arctic	17-18	25	20-22	16-18	13-14	6	4-6	14-16	6-7
			(42-44)							
Cetomimidae			(51-52)							
<i>Cetomimus</i> sp.	Bering Sea									
<i>Cetostoma regani</i>	Cent. Calif. - Oregon			30	27	14	Absent			8
<i>Gyrinomimus</i> sp.*	N. Calif. - Brit. Col.									
Rondeletiidae			(24-27)							
<i>Rondeletia loricata</i>	S. Calif. - Oregon	10	16	13-16	13-14	9-11	5	4-6	14-16	8
			(24-26)							

*Peden et al. 1985.



Zeiformes

The zeiforms occur in tropic and temperate areas of all oceans, in benthic and pelagic habitats from shallow to deep-water areas; little else is known of the life history of most fish in this group. Adults are generally deep-bodied with large eyes, greatly distensible jaws, and dorsal, anal and/or pelvic fin spines. The order is loosely organized according to a number of characters, none of which are unique. Presently, there are 6 families, 21 genera, and about 36 species throughout their distribution (J. Nelson 1984). The order is represented in the study area by one taxon, *Alloctytus* sp., a member of the Oreosomatidae. Eggs of only three species are known. Sizes range from 1.0 to 2.25 mm in diameter; all have a smooth chorion, homogeneous yolk, and a single oil globule (Tighe and Keene 1984). Larvae are generally deep-bodied and may be heavily pigmented. Forms exhibit a variety of armaments including serrated cranial ridges and spines, preopercular spines, and hardened cones or scaly knobs.

Family in study area: **Oreosomatidae**

MERISTICS

Vertebrae	Total: 39-39-40 Precaudal: 13-X-14 Caudal: 26-26-26
Branchiostegal rays	7-7-7
Caudal fin	Total rays = 12-18
Pelvic fin	Thoracic S: 1-1-1 R: 6-6-6
Dorsal fin	S: 6-X-7 R: 30-X-36
Pectoral fin	R: 19-X-21
Anal fin	S: 2-X-3 R: 28-X-33
Gill rakers	U: 6-6-6 L: 18-X-21

LIFE HISTORY

Range	Cent. California, 34-38°N, to Bering Sea, 54-66°N
Ecology	Meso- and bathybenthal, 366-732 m
ELH pattern	Eggs and larvae unknown
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE^b

Preanal length
Length at flexion
Length at transformation
Sequence of fin
development
Pigment

Diagnostic characters - Family

- Limited data on prejuveniles only; pelagic prejuveniles are oval in outline and possess a leathery skin with distinct hardened cones or scaly knobs laterally and ventrally

^a *Allocyttus folletti* and *A. verrucosus* may occur in the study area, but identification to species is not possible. Kobayashi et al. (1968) suggested *A. folletti* should be a synonym of *A. verrucosus*.

^b Early-life-history stages of *Allocyttus* spp. are unknown. An illustration of a prejuvenile of *Oreosoma atlanticum* is presented for comparison only. A photograph of an *Allocyttus verrucosus* juvenile (95 mm TL) is presented in Kobayashi et al. (1968).






Figure A, Abe and Kaji 1972 (Tasman Sea specimen).





Gasterosteiformes

The order Gasterosteiformes is made up of small, cryptically colored fish found in freshwater or marine habitats in tropical and temperate areas. Most species have small mouths at the end of tubular snouts and dermal plates covering the body. Historically divided into two or three orders, the group, as supported by Pietsch (1978) and Fritzsche (1984), contains 10 families with well over 200 species. Eggs are broadcast or deposited in nests or on algae, or for syngnathids, in male brood pouches. Eggs of families other than Syngnathidae are 1.5-2.1 mm in diameter, may have one or more oil globules, and have thick chorions. Most gasterosteiform larvae hatching from externally incubated eggs (except gasterosteids) have the same distinctive characters (i.e., elongate snout and small mouth, dermal plates) as the adults. Although adult gasterosteids are sometimes collected in plankton samples, early-life-history stages are unlikely to occur and are not treated here.

Families in study area: **Aulorhynchidae**
Gasterosteidae
Syngnathidae

MERISTICS

Vertebrae	Total: 54-54-56
	Precaudal: 24-25-26
	Caudal: 29-29-31
Branchiostegal rays	4-4-4
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 4-4-4
Dorsal fin	S: 23-X-27 R: 9-X-11
Pectoral fin	R: X-X-X
Anal fin	S: 1-1-1 R: 9-X-10
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59°N
Ecology	Nearshore shelf pelagic
ELH pattern	Oviparous; demersal, adhesive, guarded eggs; larvae briefly pelagic
Spawning	Season: Late spring-summer (Brit. Col.); ^a Feb-July (Calif.); ^b year-round (Calif.) ^c Area: Demersal, on algae or marine plants ^a Mode: Nest building; males territorial, polygamous ^c Migration:
Fecundity	Range/function: 150-600/mass ^a
Age at first maturity	1 yr ^c
Longevity	9 yr ^b

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	2.0-2.1 mm
No. of oil globules	One
Oil globule diameter	
Yolk	Yellow to yellow-green
Envelope	Thick
Hatch size	8 mm TL
Incubation time/temp.	2-6 wk/10°C
Pigment	

Diagnostic characters

- Protracted hatch always occurs, embryos develop asynchronously

LARVAE

Preanal length	60-65% SL
Length at flexion	>16 mm TL
Length at transformation	
Sequence of fin development	Pectorals; dorsal rays, anal, and caudal; dorsal spines; pelvics


Pigment

- Gut: Along dorsal surface and ventral midline
- Postanal body: Dorsal and ventral midline to tail; along notochord initially over anal fin but increasing anteriorly
- Head: Dorsal and posterior to eye

Diagnostic characters

- Pigment generally heavier than on syngnathids
- Bony plates develop by notochord flexion
- Elongated snout

^aMarliave 1975a^bFitch and Lavenberg 1975^cLimbaugh 1962



Figures A–D, Marliave 1976.

MERISTICS

Vertebrae	Total: 56-61-64 Precaudal: 19-20-21 Caudal: 37-41-43
Branchiostegal rays	2-X-3
Caudal fin	
Pelvic fin	Absent
Dorsal fin	R: 28-38-43
Pectoral fin	R: 11-X-13
Anal fin	R: 2-X-3
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59°N
Ecology	Intertidal, nearshore
ELH pattern	Oviparous, eggs and larvae develop in brood pouch
Spawning	Season: Spring-summer ^a Area: Inshore protected areas ^b Mode: Males retain eggs and developing larvae until the young have reached a juvenile stage of development ^b Migration:
Fecundity	Range/function: 225 ^a
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.0-1.5 mm
No. of oil globules	One or more
Oil globule diameter	0.4-0.5 mm
Yolk	Bright yellow
Envelope	
Hatch size	4.5-5.0 mm TL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE^c**

Preanal length	35-40% TL
Length at flexion	
Length at transformation	
Sequence of fin development	
Pigment	• Sparse at hatching, developing to form vertical blotches covering the head and body

Diagnostic characters^a Wang 1981^b Hart 1973^c Larval illustrations of *Syngnathus leptorhynchus* are unavailable. Illustrations of early-life-history stages of other members of the genus are presented for comparison and may differ from early-life-history stages of *S. leptorhynchus*.



A *Syngnathus fuscus***B** *Syngnathus schlegeli*

Figure A, Ryder 1887 (reversed, Atlantic specimen from brooding pouch); B, Chung 1977 (western Pacific specimen).



Scorpaeniformes

Most scorpaeniforms are bottom-oriented and have large heads, large rounded pectoral fins, rounded caudal fins, and many have head spines or bony plates (or combinations thereof). One character unites the group: a bony suborbital stay extending from the third suborbital bone to the opercle. Within the Scorpaeniformes, there are 20 families, 269 genera, and about 1160 species (J. Nelson 1984). The order is usually divided into four suborders: Scorpaenoidei, Anoplopomatoidei, Hexagrammoidei, and Cottoidei. Although members of the order are distributed worldwide, more than 250 species occur in the North Pacific Ocean and Bering Sea of which 190 are endemic.

Reproduction is oviparous except in *Sebastes* where eggs are internally fertilized and live feeding young are extruded. Larvae are extremely diverse. Head spines are common in larval scorpaenoids and cottoids.

Families in study area: **Scorpaenidae**
Anoplopomatidae
Hexagrammidae
Cottidae
Agonidae
Cyclopteridae

SCORPAENIDAE

Scorpaenidae

In the study area this speciose family consists of *Sebastes*, with about 40 species, and *Sebastolobus* with 3 species. The larvae can be identified to genus, but identification is presently possible for only a few of the species of *Sebastes*. Information is included here which will aid in identifying species for which early-life-history series are known, and also characters are presented which may prove useful in identifying species whose larvae are not yet known. Full text pages are provided for those species for which an illustration is available of a postflexion larva and at least one other stage. Species with illustrations of only an extrusion larva or pelagic juvenile are treated in the comparative sections. For *Sebastes* species accounts, a section on head spines has been substituted for the egg section, and counts of lateral line pores and total gill rakers have been added to the meristic section.

Sebastinae

Among the 70 or so species of *Sebastes* in the Northeast Pacific, 40 species are known to occur in the study area (Table 29). Meristic information on the following pages has been updated from recent sources (Chen 1986; Laroche, in prep.). *Sebastes* are ovoviviparous (or viviparous¹) and give birth to live young which are similar in stage of development to first-feeding larvae of oviparous scorpaenids. Eggs, generally between 0.75 and 1.9 mm diameter, are retained in the lumen of the ovary after ovulation and internal fertilization. Eggs possess a homogeneous yolk, narrow perivitelline space, fragile smooth chorion, and one to many oil globules. Hatching precedes extrusion with newborn larvae, 3.8-7.5 mm SL, having already utilized most of their yolk.

Newly extruded larvae have functional eyes, jaws, and pectoral fins. Notochord flexion occurs at about 6-12 mm SL and transformation (acquisition of adult complement of fin rays) occurs at about 15-25 mm SL. Many species have a distinct pelagic juvenile stage that may last until about 60 mm SL. Preflexion larvae have relatively slender bodies and compact guts (13-23% SL). Snout-to-anus length may increase from 40-50% SL to >60% SL in some species during the larval period. Preflexion larvae usually have pigment over the gut and a series of postanal ventral melanophores. Some species may have pigment along the dorsal midline which develops gradually. Other pigment may occur on the brain, jaws, opercle, fins (especially pectorals), and caudal peduncle.

A prominent feature in *Sebastes* larvae is head spination. Pterotic, parietal, and preopercular spines form before larvae undergo flexion, and other spines appear gradually thereafter. The full complement of head spines generally develops during the larval period, and some spines that develop in the larval period (e.g., pterotic, anterior preopercles, and lower posttemporal, and others in some) are overgrown during the juvenile stage (Washington et al. 1984b).

Larval series of 16 species of *Sebastes* occurring in our study area are known. Extrusion or yolk-exhaustion larvae have been illustrated for 32 species, and pelagic juveniles have been illustrated for 33 species. Although *Sebastes* larvae are often abundant in plankton samples from the Northeast Pacific in winter and spring, the specific identity of most cannot presently be determined.

Sebastolobinae

Sebastolobus is represented by three species in the study area. Eggs are extruded in bilobed gelatinous masses which float at the surface. The eggs are slightly elliptical and have homogeneous yolk, a narrow perivitelline space, and a smooth chorion. A single oil globule (0.18-0.20 mm diameter) is present (Pearcy 1962, Moser 1974).

Besides differences in meristic characters, *Sebastolobus* larvae can be distinguished from those of *Sebastes* in that preflexion larvae have a postanal band of pigment instead of the ventral and possibly dorsal midline melanophores in *Sebastes*. A parietal ridge develops in *Sebastolobus* and terminates in a double spine, the parietal and nuchal. In *Sebastes* the parietal spine develops singly, usually with a smaller nuchal spine immediately posterior to it.

¹Boehlert and Yoklavich (1984) demonstrated that in *Sebastes melanops* the embryos receive some nutrition from the ovarian fluid during gestation.

Table 29

Distribution, parturition season, and meristic characters of genus *Sebastes*. Range of counts followed by mode in parentheses. All have modal counts of 13 dorsal spines (except *S. polyspinis* with 14^a) and 3 anal spines (Chen 1986).

Taxon	Distribution	Parturition	Fin rays			Total gill rakers on first arch	No. of lateral line pores
			Dorsal	Anal	Pectoral		
<i>S. aleutianus</i>	S. Calif.-Bering Sea	Apr ^b	13-15 (14)	7-8 (7)	17-19 (18)	28-33 (31)	30-33
<i>S. alutus</i>	S. Calif.-Bering Sea	Jan-May ^c	13-16 (15)	7-9 (8)	17-19 (18)	34-39 (35/38)	46-55
<i>S. auriculatus</i>	SSC-SE Alaska	Mar-June ^d	12-14 (13)	6-7 (7)	16-19 (18)	25-29 (27)	42-50
<i>S. aurora</i>	S. Calif.-Brit. Col.	Mar-May ^e	12-13 (13)	5-7 (6)	17-19 (17)	24-28 (27)	28-31
<i>S. babcocki</i>	S. Calif.-Bering Sea	Apr-May ^b	13-15 (14)	6-8 (7)	17-20 (19)	29-33 (31)	42-51
<i>S. borealis</i>	Cent. Calif.-Bering Sea	Apr ^c	12-15 (13)	6-8 (7)	17-20 (19)	27-31 (30)	28-32
<i>S. brevispinis</i>	SSC-Bering Sea	June ^b , June-July ^c	13-15 (14)	7-7 (7)	17-18 (18)	33-36 (33)	46-51
<i>S. caurinus</i>	SSC-Gulf of Alaska	Apr ^b	11-14 (13)	5-7 (6)	16-18 (17)	26-31 (28)	39-47
<i>S. chlorostictus</i>	SSC-Wash.	July ^f	11-13 (12)	5-7 (6)	16-18 (17)	31-36 (33)	35-43
<i>S. ciliatus</i>	Brit. Col.-Bering Sea		14-16 (15)	6-9 (8)	17-19 (18)	32-35 (35)	42-53
<i>S. crameri</i>	S. Calif.-Bering Sea	Feb ^c	13-15 (13)	7-7 (7)	18-20 (19)	30-34 (32)	40-51
<i>S. diploproa</i>	SSC-Gulf of Alaska	May-June ^b , July ^c	11-14 (12)	5-7 (7)	17-18 (18)	32-37 (34)	32-42
<i>S. elongatus</i>	SSC-Gulf of Alaska	May-June ^b	12-14 (13)	6-7 (6)	16-18 (17)	29-33 (31)	37-47
<i>S. emphaeus</i>	N. Calif.-Gulf of Alaska	Aug-Sept ^g	13-15 (14)	6-7 (7)	16-18 (17)	37-41 (40)	40-46
<i>S. entomelas</i>	SSC-Gulf of Alaska	Apr ^c , Nov-Mar ^h	14-16 (15)	7-9 (8)	18-19 (18)	34-38 (36)	54-60
<i>S. flavidus</i>	S. Calif.-Gulf of Alaska	Mar ^c , Nov-Feb ^h	14-15 (14)	7-8 (8)	17-18 (18)	34-37 (36)	49-55
<i>S. glaucus</i>	Bering Sea		14-17 (15)	7-9 (8)	18-20 (19)	35-41 (38)	
<i>S. goodei</i>	SSC-Brit. Col.	Dec-Mar ^h	13-14 (14)	8-9 (8)	16-18 (17)	34-39 (36)	50-57
<i>S. helvomaculatus</i>	SSC-Gulf of Alaska	June ^c	12-14 (13)	6-7 (6)	15-17 (16)	28-33 (30)	34-45
<i>S. jordani</i>	SSC-Gulf of Alaska	Mar ^c , Nov-Mar ^h	13-16 (14)	9-10 (9)	19-22 (20)	40-47 (44)	53-64
<i>S. maliger</i>	Cent. Calif.-Gulf of Alaska	Apr ^c	11-13 (13)	6-7 (7)	16-18 (17)	29-33 (31)	34-45
<i>S. melanops</i>	Cent. Calif.-Aleutian Is.	Apr ^b	14-15 (15)	7-9 (8)	18-19 (19)	34-38 (37)	47-55
<i>S. melanostictus</i>	Bering Sea		13-14 (14)	7-7 (7)	18-18 (18)	32-35 (33)	
<i>S. melanostomus</i>	SSC-Wash.		12-14 (13)	6-8 (7)	18-20 (19)	30-34 (31)	28-33
<i>S. miniatus</i>	SSC-Brit. Col.	Nov-Mar ^h	13-15 (14)	6-7 (7)	17-18 (18)	36-42 (41)	41-48
<i>S. mystinus</i>	SSC-Aleutian Is.	Nov-Jan ⁱ	15-17 (16)	8-10 (9)	17-19 (18)	32-38 (35)	47-53
<i>S. nebulosus</i>	S. Calif.-SE Alaska		12-14 (13)	6-7 (7)	17-18 (18)	27-31 (28)	38-43
<i>S. nigrocinctus</i>	Cent. Calif.-SE Alaska	May ^c	12-15 (14)	6-7 (7)	18-19 (19)	28-32 (28)	36-46 ^k
<i>S. paucispinis</i>	SSC-Gulf of Alaska	Jan-Feb ^c , Nov-Apr ^j	13-15 (14)	8-10 (9)	14-16 (15)	27-32 (28)	51-62
<i>S. pinniger</i>	SSC-Gulf of Alaska	Jan-Mar ^c , Nov-Mar ^f	13-15 (14)	7-7 (7)	16-18 (17)	41-44 (43)	40-47
<i>S. polyspinis</i>	Brit. Col.-Bering Sea		13-16 (15)	7-9 (8)	17-19 (18)	35-39 (36)	43-53
<i>S. proriger</i>	S. Calif.-Bering Sea	July ^c	14-15 (15)	7-7 (7)	16-17 (17)	36-41 (38)	48-55
<i>S. rastrelliger</i>	SSC-Oregon		12-13 (13)	6-6 (6)	18-20 (19)	17-25 (23)	40-49
<i>S. reedi</i>	N. Calif.-SE Alaska	Apr ^c	13-15 (14)	7-8 (7)	18-20 (19)	30-36 (34)	47-55
<i>S. ruberrimus</i>	SSC-Gulf of Alaska	June ^b , July ^c	14-16 (15)	7-7 (7)	18-19 (19)	26-30 (29/30)	39-45
<i>S. rufus</i>	SSC-Oregon	Dec-May ^e	13-16 (15)	8-9	17-19 (18)	32-37 (35)	49-56
<i>S. saxicola</i>	SSC-Gulf of Alaska	Feb ^c , Nov-Mar ^h	11-13 (12)	6-7 (7)	16-18 (16)	30-35 (32)	35-43
<i>S. variegatus</i>	Brit. Col.-Gulf of Alaska	June ^c	14-15 (14)	6-7 (7)	17-19 (18)	37-41 (38)	43-52
<i>S. wilsoni</i>	S. Calif.-Gulf of Alaska	June ^c	13-15 (14)	5-6 (6)	16-18 (17)	37-42 (39)	37-46
<i>S. zacentrus</i>	S. Calif.-Gulf of Alaska	July ^c	13-15 (14)	7-8 (7)	16-18 (17)	33-37 (35)	39-47

^aIncorrectly reported as 13 instead of 14 in Chen (1986:Table 2). L.-C. Chen, Dep. Zool., San Diego St. Univ., San Diego, CA 92192, pers. commun., 21 July 1988.

^bOff Canada (Hart 1973).

^cOff Canada (Westrheim 1975).

^dPuget Sound (Washington et al. 1978).

^eNorth-central California (Wyllie Echeverria 1987).

^fOff California (Moser 1967).

^gPuget Sound (Moulton 1975).

^hOff north-central California (Phillips 1964).

ⁱOff California (Wales 1952).

^jOff California (Frey 1971).

^kOne rare count of 50.

Head spines


There are differences in terminology for *Sebastes* head spines between the two major sources of early-life-history descriptions (i.e., Richardson and Laroche 1979, Moser and Ahlstrom 1978). Most of the differences are semantic; however, the numbering of anterior preopercular spines is a substantive difference (Table 30). Among the species studied to date, posterior preopercular spine 1 has not had an anterior spine associated with it. Therefore, Richardson and Laroche (1979) gave the number 1 to the anterior preopercular spine associated with the second posterior preopercular spine. Moser and Ahlstrom (1978) gave the same numbers to the anterior and posterior preopercular spines. Thus the same anterior preopercular spine is numbered one less by Richardson and Laroche (1979) than it is by Moser and Ahlstrom (1978). Succeeding descriptions by these and other authors perpetuate these differences (e.g., Washington et al. 1984b). Here we use the system of Moser and Ahlstrom (1978).

There are also differences between the *Sebastes* head spine terminology used for adults and that used for larvae (Table 30). This situation resulted from the difficulty of determining spine/bone associations by external examination of adults and is complicated by the overgrowth of some spines during development (Richardson and Laroche 1979). The major points of confusion are that the supracleithral of larvae is called the upper posttemporal spine of adults, and the cleithral spine of larvae is called the supracleithral spine of adults. Head spine patterns which may help identify adult *Sebastes* are listed in Table 31.

Table 30

Head spine terminology in *Sebastes*, listed by sequence in which the spines develop in *S. melanostomus* (Moser and Ahlstrom 1978). Names used by Richardson and Laroche (1979) are listed only when they differ from those used by Moser and Ahlstrom (1978).

Abbreviation	Moser and Ahlstrom (1978)	Richardson and Laroche (1979)	Bone of origin	Adult spine
PA	Parietal		Parietal	Parietal
NU	Nuchal		Parietal	Nuchal
PSO	Postocular		Frontal	Postocular
PT	Pterotic		Pterotic	(Overgrown)
LPST	Lower posttemporal	Inferior posttemporal	Posttemporal	(Overgrown)
UOP	Upper opercular	Superior opercular	Opercle	Opercular
UIO-1	1st upper infraorbital	Superior infraorbital series, 1st	Infraorbital 1	(Overgrown)
LIO-1	1st lower infraorbital	Inferior infraorbital series, 1st	Infraorbital 1	Lachrymal projection (suborbital spine)
APO-2	2nd anterior preopercular	Anterior preopercular series, 1st	Preopercle	(Overgrown)
PPO-2	2nd posterior preopercular	Posterior preopercular series, 2nd	Preopercle	Preopercular
APO-3	3rd anterior preopercular	Anterior preopercular series, 2nd	Preopercle	(Overgrown)
PPO-3	3rd posterior preopercular	Posterior preopercular series, 3rd	Preopercle	Preopercular
APO-4	4th anterior preopercular	Anterior preopercular series, 3rd	Preopercle	(Overgrown)
PPO-4	4th posterior preopercular	Posterior preopercular series, 4th	Preopercle	Preopercular
PPO-1	1st posterior preopercular	Posterior preopercular series, 1st	Preopercle	Preopercular
PPO-5	5th posterior preopercular	Posterior preopercular series, 5th	Preopercle	Preopercular
UIO-4	4th upper infraorbital	Superior infraorbital series, 4th	Infraorbital 3	(Overgrown)
SC	Supracleithral		Supracleithrum	Cleithral
LIO-2	2nd lower infraorbital	Inferior infraorbital series, 2nd	Infraorbital 1	Lachrymal projection (suborbital spine)
LIO-3		Inferior infraorbital series, 3rd	Infraorbital 1	(Overgrown)
LOP	Lower opercular	Inferior opercular	Opercle	Opercular
UIO-2	2nd upper infraorbital	Superior infraorbital series, 2nd	Infraorbital 1	(Overgrown)
UPST	Upper posttemporal	Superior posttemporal	Posttemporal	Supracleithral
SPO	Supraocular		Frontal	Supraocular
UIO-3	3rd upper infraorbital	Superior infraorbital series, 3rd	Infraorbital 2	(Overgrown)
NA	Nasal		Nasal	Nasal
PRO	Preocular		Lateral ethmoid (prefrontal)	Preocular
CL	Cleithral		Cleithrum	(Overgrown)
IOP	Interopercular		Interopercular (interopercle)	Gill cover spine
TM	Tympanic		Frontal	Tympanic
SOP	Subopercular		Subopercular (subopercle)	Gill cover spine
CO	Coronal		Frontal	Coronal



Positions and abbreviations of larval head spines in larval *Sebastes*. Based on a 16.0-mm stained larva of *Sebastes melanostomus* augmented to show the position of the coronal (CO) and 3rd lower infraorbital (LIO-3) spines. From Moser and Ahlstrom (1978) and Laroche (in prep.).

Table 31
Presence and absence of head spines which are diagnostic for adults of members of the genus *Sebastes*
(+ indicates presence, 0 indicates absence, blank indicates unknown character state).

	Pre-ocular	Supra-ocular	Post-ocular	Tympanic	Coronal		Pre-ocular	Supra-ocular	Post-ocular	Tympanic	Coronal
<i>S. aleutianus</i>	+	+	+	+	+	<i>S. maliger</i>	+	0	+	+	0
<i>S. alutus</i>	+	+	+	+	0	<i>S. melanops</i>	0	0	+	+	0
<i>S. auriculatus</i>	+	0	+	+	+	<i>S. melanostictus</i>			+	+	
<i>S. aurora</i>	+	+	+	+	0	<i>S. melanostomus</i>	+	+	+	+	0
<i>S. babcocki</i>	+	0	+	+	0	<i>S. miniatus</i>	+	+	+	+	0
<i>S. borealis</i>	+	+	+	+	+	<i>S. mystinus</i>	+	+	+	+	0
<i>S. brevispinis</i>	0	0	+	+	0	<i>S. nebulosus</i>	+	0	+	+	0
<i>S. caurinus</i>	+	0	+	+	0	<i>S. nigrocinctus</i>	+	0	+	+	+
<i>S. chlorostictus</i>	+	+	+	+	0	<i>S. paucispinis</i>	+	0	0	+	0
<i>S. ciliatus</i>	0	+	+	+	0	<i>S. pinniger</i>	+	+	+	+	0
<i>S. crameri</i>	+	+	+	+	0	<i>S. polyspinis</i>	0	0	+	+	0
<i>S. diploproa</i>	+	0	+	+	0	<i>S. proriger</i>	+	0	+	+	0
<i>S. elongatus</i>	+	0	+	+	0	<i>S. rastrelliger</i>	+	0	+	+	0
<i>S. emphaeus</i>	+	0	+	+	0	<i>S. reedi</i>	+	+	+	+	0
<i>S. entomelas</i>	+	+	+	+	0	<i>S. ruberrimus</i>	+	+	+	+	0
<i>S. flavidus</i>	0	0	+	+	0	<i>S. rufus</i>			+	+	0
<i>S. glaucus</i>			+	+	0	<i>S. saxicola</i>	+	0	+	+	0
<i>S. goodei</i>	0	0	+	0	0	<i>S. variegatus</i>	+	0	+	+	0
<i>S. helvomaculatus</i>	+	+	+	+	0	<i>S. wilsoni</i>	+	0	+	+	0
<i>S. jordani</i>	+	0	+	+	0	<i>S. zacentrus</i>	+	0	+	+	0


Extrusion or Yolk-Exhaustion Larvae

Various authors have illustrated and described larvae which have been extruded from pregnant female *Sebastes*. In some cases, the larvae have been maintained in seawater for several days after extrusion, and in a few cases (e.g., Stahl-Johnson 1985, Moser and Butler 1987) the larvae were fed and reared. With handling, pregnant *Sebastes* will release larvae that are not full-term, and such larvae may be represented in some illustrations. Furthermore, some illustrations were drawn by adding observed pigmentation for a particular species to a general outline of an extrusion *Sebastes* larva (c.f. Westrheim et al. 1968a). The generalized outline used by Westrheim et al. (e.g., 1968a) and DeLacy et al. (1964) lacked pectoral fins, apparently the result of an oversight. Several illustrations of extrusion larvae have been published for some species, and there are notable differences among some of these illustrations. One of the illustrations of an extrusion larva purported to be *Sebastes zacentrus* by Harling et al. (1971) was obviously misidentified, as discussed by Laroche and Richardson (1981). With these problems in mind, pages 273-280 present all of the published, and some previously unpublished, illustrations of extrusion larvae of *Sebastes* spp. which occur in the study area, not so much for use in species identifications but as reference for further research.


EXTRUSION OR YOLK-EXHAUSTION LARVAE

S. aleutianus

A




B




S. alutus

C



D




E




S. auriculatus

F




G





Figures A, E, Westrheim et al. 1968a; B-C, Efremenko and Lisovenko 1970; D, F, DeLacy et al. 1964; G, Moser et al. 1977.

EXTRUSION OR YOLK-EXHAUSTION LARVAE

A *S. aurora*



S. babcocki



S. brevispinis









Figure A, Moser et al. 1985; B, Westrheim et al. 1968a; C, F, DeLacy et al. 1964; D, E, Efremenko and Lisovenko 1970; G, NWAFC original (B. Vinter).

EXTRUSION OR YOLK-EXHAUSTION LARVAE


S. caurinus




D *S. chlorostictus*



E *S. ciliatus*



F *S. crameri*



Figures A, D, Moser et al. 1977; B, DeLacy et al. 1964; C, NWAFC original (B. Vinter); E, Harling et al. 1971; F, Westrheim et al. 1968a.

EXTRUSION OR YOLK-EXHAUSTION LARVAE


S. diploproa

A



\bar{x} = 4.7 mm TL


B



~ 5.0 mm TL


S. elongatus

C



\bar{x} = 4.9 mm TL


D



4.9 mm


S. entomelas

E




\bar{x} = 4.8 mm TL

F



4.8 mm

G *S. flavidus*




~ 4.5 mm TL


Figures A, C, Westrheim et al. 1968b; B, G, DeLacy et al. 1964; D, Moser et al. 1977; E, Harling et al. 1971; F, Moser and Butler 1987.

EXTRUSION OR YOLK-EXHAUSTION LARVAE


A *S. goodei*





B *S. helvomaculatus*




C *S. jordani*




S. maliger



F *S. melanops*




G *S. melanostomus*



Figures A, G, Moser et al. 1977; B, D, Westrheim et al. 1968b; C, F, NWAFC originals (B. Vinter); E, DeLacy et al. 1964.

EXTRUSION OR YOLK-EXHAUSTION LARVAE


A *S. miniatus*




B *S. mystinus*




C *S. paucispinis*




D *S. pinniger*



E *S. polyspinis*




F *S. proriger*



Figures A, C–D, Moser et al. 1977; B, Wold, unpubl.; E, NWAFC original (B. Vinter); F, Westrheim et al. 1968b.

EXTRUSION OR YOLK-EXHAUSTION LARVAE


A *S. reedi*



\bar{x} = 6.0 mm TL


S. ruberrimus

B




\bar{x} = 4.4 mm TL

C




\bar{x} = 4.2 mm TL

D



~ 5.0 mm TL


E *S. rufus*



5.3 mm


S. saxicola

F



4.7 mm

G




\bar{x} = 5.2 mm TL

Figures A, G, Westrheim et al. 1968a; B, Harling et al. 1971; C, Westrheim et al. 1968b; D, DeLacy et al. 1964; E-F, Moser et al. 1977.

EXTRUSION OR YOLK-EXHAUSTION LARVAE


S. variegatus

A



\bar{x} = 4.8 mm TL


B



4.6 mm SL

S. zacentrus


C



\bar{x} = 5.0 mm TL


(Misidentified, see Laroche and Richardson 1981)

D




\bar{x} = 4.2 mm TL

E




\bar{x} = 4.5 mm TL

F



\bar{x} = 5.0 mm TL

G



4.7 mm SL

Figures A, C, F, Harling et al. 1971; B, G, NWAFC originals (B. Vinter); D, Efremenko and Lisovenko 1970; E, Westrheim et al. 1968b.

Sebastes Pelagic Juveniles

The following six pages present illustrations of pelagic juveniles of 33 of the 40 species of *Sebastes* occurring in the study area. Pelagic juveniles of the remaining species are unknown or have not yet been illustrated. Illustrations are grouped to form a pictorial key based primarily on pigment characters of the juveniles as derived from Laroche (in prep.). Species are grouped on the pages according to predominant pigment characters; within these groups, species are separated according to other pigment characters. Several species possess identical states for these characters. Identification of these will require use of additional characters such as meristics, body shape, head spination, time and location of collection, and other pigment characters. These characters should also be checked on fish separated based on the pictorial key to verify their identity. Variation in pigment can be expected due to size and other differences in specimens of a species. Also, pigment characters of species for which pelagic juveniles are unknown may closely resemble those species illustrated here, which would lead to misidentification unless other characters were examined.

PELAGIC JUVENILES
Black pigment blotch at posterior of spinous dorsal fin











Diffuse body pigment	Pectoral fin base unpigmented	Anal fin unpigmented	Lateral midline pigmented	<p>A <i>S. melanops</i></p>  <p>45.3 mm SL</p>
			Lateral midline unpigmented	<p>B <i>S. flavidus</i></p>  <p>42.0 mm SL</p>
		Ventral caudal peduncle pigmented	Ventral caudal peduncle unpigmented	<p>C <i>S. entomelas</i></p>  <p>48.0 mm SL</p>
		Anal fin pigmented	<p>D <i>S. mystinus</i></p>  <p>43.0 mm SL</p>	
	Pectoral fin base pigmented	<p>E <i>S. miniatus</i></p>  <p>28.0 mm SL</p>		
		<p>F <i>S. pinniger</i></p>  <p>40.0 mm SL</p>		
Banded body pigment				

Figure A, Laroche and Richardson 1980; B–F, Laroche, in prep.

PELAGIC JUVENILES
 Fringe of pigment on spinous dorsal fin
 Body pigment diffuse

Anal fin without distinct stripe of pigment	No fringe of pigment on pectoral and pelvic fins	<p>A <i>S. alutus</i></p>  <p>57.0 mm SL</p>
	Fringe of pigment on pectoral and pelvic fins	<p>B <i>S. paucispinis</i></p>  <p>46.0 mm SL</p>
Both soft dorsal and anal fins have distinct stripe of pigment	Caudal fin with distinct band of pigment	<p>C <i>S. aleutianus</i></p>  <p>57.0 mm SL</p>
	Caudal fin without distinct band of pigment	<p>D <i>S. diploproa</i></p>  <p>39.0 mm SL</p>


Figures A–D, Laroche, in prep.

PELAGIC JUVENILES
Body with three or more distinct bands of pigment

Saddles or bands do not extend onto dorsal or anal fins		Saddles or bands extend onto spinous and/or soft dorsal fins				
Bands slant anteroventrally		Bands extend onto soft dorsal fin only	Bands extend onto both spinous and soft dorsal fins			
		Fringe of pigment on pectoral	Pectoral fin totally pigmented	Pectoral fin pigmented proximally		
			Two or more bands of pectoral pigment	Pectoral uniformly pigmented	Distinct stripe on anal fin	Anal fin pigment uniform
	F <i>S. saxicola</i> 36.0 mm SL	E <i>S. reedi</i> 49.0 mm SL	D <i>S. nigrocinctus</i> 53.0 mm SL	C <i>S. camerti</i> 39.0 mm SL	B <i>S. melanostomus</i> 46.2 mm SL	A <i>S. caurinus</i> 36.0 mm SL
G <i>S. goodeti</i> 37.0 mm SL						

PELAGIC JUVENILES
Diffuse body and dorsal fin pigment
Ventral caudal peduncle pigmented


A *S. auriculatus*



31.0 mm SL


Body pigment aligned
along myosepta

B *S. proriger*



38.0 mm SL


C *S. zacentrus*



30.0 mm SL

No distinct pigment along proximal margin of caudal fin


D *S. rastrelliger*



27.0 mm SL

Distinct stripe of pigment
along proximal margin of
soft dorsal and anal fins

E *S. brevispinis*




41.0 mm SL

Distinct pigment along proximal margin of caudal fin







No distinct stripe of
pigment along proximal
margin of soft dorsal
and anal fins

PELAGIC JUVENILES
Diffuse body and dorsal fin pigment
Ventral caudal peduncle unpigmented



Figures A–D, Laroche, in prep.; E, NWAFC original (B. Vinter).

PELAGIC JUVENILES
 Body with fewer than three pigment bands, with stripes of pigment,
 or with a patch of pigment on the caudal peduncle

Patch of pigment on caudal peduncle*	Uniform heavy body and head pigment	<p>A <i>S. helvomaculatus</i></p>  <p>41.6 mm SL</p>
	Unpigmented area on trunk	<p>B <i>S. ruberrimus</i></p>  <p>28.0 mm SL</p>
	Longitudinal stripes on body of larger specimens	<p>C <i>S. elongatus</i></p>  <p>32.0 mm SL</p>
No distinct patch of pigment on caudal peduncle*	Body with two bands of pigment	<p>D <i>S. babcocki</i></p>  <p>16.7 mm SL</p>
	<p>E <i>S. chlorostictus</i>*</p> <p>*Smaller specimens of <i>S. chlorostictus</i> have patch of pigment on caudal peduncle, becomes band in larger specimens.</p>	 <p>21.0 mm SL</p>
		<p>F <i>S. aurora</i></p>  <p>13.5 mm SL</p>

Figures A–E, Laroche, in prep.; F, Moser et al. 1985.

MERISTICS

Vertebrae	Total: 27-27-27
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-14 R: 13-14-15
Pectoral fin	R: 17-18-19
Anal fin	S: 3-3-3 R: 7-7-8
Gill rakers	U: 8-9-11 L: 20-22-24
	(T: 28-31-33)
Lateral line pores	30-33

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Meso- and bathybenthal, 183-732 m
ELH pattern	Ovoviparous, pelagic larvae
Spawning	Season: Apr ^a Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	+
Supraocular	+
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	4.1-5.3 mm TL
Preanal length	
Length at flexion	
Length at transformation	
Sequence of fin development	
Pigment	<ul style="list-style-type: none"> • Extrusion larvae: Series along ventral body • Postflexion larvae >19.3 mm SL: Pigment covering entire body except on lower cheek, pectoral fin base, and tail tip

Diagnostic characters**PELAGIC JUVENILES**

- Body pigment light, diffuse
- Banded pigment on median fins

^aHart 1973

Ref: Laroche, in prep.






Figure A, Westrheim et al. 1968a; B-C, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-27 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	9-11, 8+7, 9-11
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 12-13-14
Pectoral fin	R: 16-18-19
Anal fin	S: 3-3-3 R: 6-7-7
Gill rakers	U: 8-8-10 L: 18-20-21 (T: 25-27-29)
Lateral line pores	42-50

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59°N
Ecology	Nearshore shelf demersal
ELH pattern	Ovoviviparous, pelagic larvae
Spawning	Season: Mar-June ^a Area: Mode: Migration:
Fecundity	Range/function: 52,000-339,000 ^b
Age at first maturity	5 yr (females) ^c
Longevity	

^aWashington et al. 1978^bHart 1973^cWyllie Echeverria 1987^dDescription of pigment and illustrations based on laboratory-reared specimens which may be more melanistic than wild-caught specimens.Ref: Laroche, in prep.; Stahl-Johnson (1985), who noted a mistake in Moser et al. (1977); yolk-depleted *S. auriculatus* figure reversed with *S. caurinus*.**HEAD SPINES**

Preocular	+
Postocular	+
Coronal	+
Supraocular	0
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	4.7-6.7 mm SL
Preanal length	<50% SL
Length at flexion	Between 7.4 and 9.0 mm SL

Length at transformation

Sequence of fin development	Caudal and pectorals, pelvics, dorsal and anal
-----------------------------	--

Pigment^d


- Mostly on dorsal head, with development covering entire head
- Nape
- Gut: Dorsal and ventrolateral surface, increasing laterally with development
- Dorsal and ventral midline melanophores begin posterior to anus; with development, dorsal melanophores extend to nape
- Increase in lateral pigment on posterior half of body and to hypural region
- Pectoral fins

Diagnostic characters




- Distinguished from *S. caurinus* (p. 296) by
- Dorsal midline melanophores postanally at birth, gradually spread forward to nape
 - More pigment on opercular than in *S. caurinus*

PELAGIC JUVENILES

- Diffuse body pigment
- Opercular pigment blotch present
- Lateral midline pigmented



Unpigmented area dorsally,
myomeres 1-8



Figures A-C, Stahl-Johnson 1985 (reared); D, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: 10-10-10 Caudal: 16-16-16
Branchiostegal rays	7-7-7
Caudal fin	10, 8+7, 9-10
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 12-13-13
Pectoral fin	R: 17-17-19
Anal fin	S: 3-3-3 R: 5-6-7
Gill rakers	U: 7-8-8 L: 18-19-20 (T: 24-27-28)
Lateral line pores	28-31

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epi-, meso-, and bathybenthal, 125-768 m
ELH pattern	Ovoviviparous, pelagic larvae
Spawning	Season: Mar-May (north-central California) ^a Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

HEAD SPINES

Preocular ^b	+
Postocular	+
Coronal	0
Supraocular	+
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	4.4 mm SL
Preanal length	~44% increasing with development to 64% SL
Length at flexion	6.5-8.6 mm SL
Length at transformation	13 mm SL
Sequence of fin development	Caudal; pectorals; pelvics, dorsal, and anal

Pigment

- Head: Dorsal spots spreading with development to opercle then to entire head
- Snout and jaws pigmented
- Gut: Posterior pigment increasing dorsolaterally with development
- Postanal body: Dorsal and ventral midline over body (between postanal myomeres 4 and 15); lateral pigment increases with development to form a band
- Pectoral fin first, other fins with development


Diagnostic characters

- Morphology: Snout-to-anus length greater than in other species studied
- Pigment: Snout, postanal band


PELAGIC JUVENILES^c

- Low total gill raker count (24-28) and lateral line pore count (28-31) are diagnostic for the species
- Body pigment faintly banded with unpigmented caudal peduncle
- Pectoral rays heavily pigmented medially




^aWyllie Echeverria 1987^bAccording to Moser et al. (1985), the preocular spine develops by 10.5 mm SL but is not visible on their 13.5-mm SL specimen shown here.^cLargest pelagic juvenile collected is 34.4 mm SL.



4.4 mm



4.8 mm



Figures A–E, Moser et al. 1985.

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 13-14-15
Pectoral fin	R: 17-19-20
Anal fin	S: 3-3-3 R: 6-7-8
Gill rakers	U: X-X-X L: X-X-X (T: 29-31-33)
Lateral line pores	42-51

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthic, 91-475 mm
ELH pattern	Ovoviparous, pelagic larvae
Spawning	Season: Apr-May ^a Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	4 yr (females) ^b
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	0
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	4.2-5.3 mm
Preanal length	
Length at flexion	
Length at transformation	
Sequence of fin development	
Pigment	<ul style="list-style-type: none"> • Extrusion larvae <ul style="list-style-type: none"> —Series along ventral body —Shorter series along posterior dorsal midline • Postflexion larvae >14.6 mm SL <ul style="list-style-type: none"> —Wide postanal band —Light pigment dorsally on head and laterally on gut

Diagnostic characters**PELAGIC JUVENILES**

- Body pigment in two bands
- Little pigment on median fins

^aHart 1973^bWyllie Echeverria 1987

Ref: Laroche, in prep.






Figure A, Westheim et al. 1968a; B-C, Laroche, in prep.

MERISTICS

Vertebrae	Total: 25-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	9-11, 8+7, 9-11
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 11-13-14
Pectoral fin	R: 16-17-18
Anal fin	S: 3-3-3 R: 5-6-7
Gill rakers	U: 8-X-10 L: 19-X-21 (T: 26-28-31)
Lateral line pores	39-47

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Nearshore shelf demersal, 0-183 m
ELH pattern	Ovoviparous, pelagic larvae
Spawning	Season: Apr ^a Area: Mode: Migration:
Fecundity	Range/function: 20,000-640,000 ^{a/} $F = 0.000000027404 \times L^{4.9567}$, L = TL mm; ^b $F = 0.000000034554 \times L^{5.30011}$, L = TL mm ^c
Age at first maturity	3-4 yr ^d 6 yr (females) ^e
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	0
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	5.3 mm SL
Preanal length	41% SL increasing with development to 49% SL
Length at flexion	Between 7.5 and 9.5 mm SL

Length at transformation

Sequence of fin development	Caudal and pectorals, pelvics, dorsal and anal
-----------------------------	--

Pigment

- Head: Dorsal and nape spreading, with development, to entire head
- Gut: Dorsal and ventrolateral surface, increasing laterally with development
- Dorsal midline melanophores extend from head to just anterior to tail with no break over gut
- Ventral midline melanophores, with development increasing ventrolaterally and in hypural region

Diagnostic characters

- See notes on *S. auriculatus* (p. 290) about reared specimens
- Distinguished from *S. auriculatus* by
- Until midflexion, more dorsal midline pigment anterior to anus
- Less pigment in the opercular region (flexion and postflexion)

PELAGIC JUVENILES

- Four pigment bands on body
- Fins pigmented

^aHart 1973^bWashington et al. 1978^cDeLacy et al. 1964^dPatten 1973^eWyllie Echeverria 1987







Figure A, NWAFC original (B. Vinter); B–C, Stahl-Johnson 1985 (reared); D, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 13-13-15
Pectoral fin	R: 17-18-19
Anal fin	S: 3-3-3 R: 7-7-7
Gill rakers	U: 8-9-10 L: 21-23-25 (T: 30-32-34)
Lateral line pores	40-51

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthal, 29-549 m
ELH pattern	Ovoviviparous, pelagic larvae
Spawning	Season: Feb (Brit. Col. ^a -Oregon ^b); Nov-Mar in southern part of range ^c Area: Mode: Migration:
Fecundity	Range/function: 50,000-609,800 ^c
Age at first maturity	5-6 yr ^c 4 yr (females) ^d
Longevity	30 yr ^c

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	+
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	5.7 mm SL
Preanal length	54-65% SL
Length at flexion	8.0-9.3 mm SL
Length at transformation	16-21 mm SL ^e
Sequence of fin development	Caudal and pectorals, pelvics, dorsal and anal

Pigment

- Extrusion larvae: Series along ventral body
- Pigmented pectorals and pelvics
- Heavy nape and top of head
- No dorsal midline
- ~11 ventral midline melanophores reducing to ~4 on caudal peduncle
- Spinous dorsal fin
- Tip of lower jaw

Diagnostic characters


- Heavy nape and paired fin pigment
- Pigment on spinous dorsal fin

PELAGIC JUVENILES


- Five pigment bands on body, four extending onto dorsal fin
- Paired fins heavily pigmented

^aHart 1973^bHitz 1962^cPhillips 1964^dWyllie Echeverria 1987^eJuveniles become benthic at 40-60 mm SL.


A



B



C



D




Figure A, Westrheim et al. 1968a; B–C, Richardson and Laroche 1979; D, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 11-12-14
Pectoral fin	R: 17-18-18
Anal fin	S: 3-3-3 R: 5-7-7
Gill rakers	U: 9-10-12 L: 23-24-27 (T: 32-34-37)
Lateral line pores	32-42

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesobenthal, 91-579 m
ELH pattern	Ovoviviparous, pelagic larvae
Spawning	Season: May-June; ^a July; ^b May-Aug (Oregon) ^c Area: Mode: Migration:
Fecundity	Range/function: 14,000-304,000 ^d
Age at first maturity	4-5 yr ^d 7 yr (females) ^e
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	0
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size 5.2 mm

Preanal length

Length at flexion

Length at transformation

Sequence of fin

development

Pigment

- Extrusion larvae: Series along ventral body
- Postflexion larvae >8.7 mm SL
 - Pigment at base of parietal spine, increasing with development to head and nape
 - Internal spots along anal fin pterygiophores
 - Along hypural margin

Diagnostic characters

PELAGIC JUVENILES

- Body and fins (except distal portions) pigmented
- Spinous dorsal fin with fringe of pigment
- Distinct band of pigment on soft dorsal and anal fins

^aHart 1973

^bWestrheim 1975

^cHitz 1962

^dPhillips 1964

^eWyllie Echeverria 1987




Figure A, Westrheim et al. 1968b; B-D, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-26
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 12-13-14
Pectoral fin	R: 16-17-18
Anal fin	S: 3-3-3 R: 6-6-7
Gill rakers	U: 8-9-10 L: 20-22-23
	(T: 29-31-33)
Lateral line pores	37-47

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesobenthic, 61-402 m
ELH pattern	Ovoviviparous, pelagic larvae
Spawning	Season: May-June ^a
	Area:
	Mode:
	Migration:
Fecundity	Range/function:
Age at first maturity	7 yr (females) ^b
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	0
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size ~5 mm

Preanal length

Length at flexion

Length at transformation

Sequence of fin

development

Pigment

- Extrusion larvae: Series along ventral body
- Postflexion larvae >17.4 mm SL
 - Along dorsal body margin
 - Internal row of spots along lateral midline
 - Above and below notochord along caudal peduncle
 - A few spots along ventral body margin

Diagnostic characters

PELAGIC JUVENILES

- Body pigment blotchy, longitudinal stripe on larger specimens
- Fins lightly pigmented, or unpigmented

^aHart 1973

^bWyllie Echeverria 1987

Ref: Laroche, in prep.






Figure A, Moser et al. 1977; B-C, Laroche, in prep.

MERISTICS

Vertebrae	Total: 27-27-28
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 13-14-15
Pectoral fin	R: 16-17-18
Anal fin	S: 3-3-3 R: 6-7-7
Gill rakers	U: 11-11-13 L: 28-30-31 (T: 37-40-41)
Lateral line pores	40-46

LIFE HISTORY

Range	N. California, 38-42°N, to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesobenthic, 10-366 m
ELH pattern	Ovoviparous, pelagic larvae
Spawning	Season: Aug-Sept ^a Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	2-3 yr ^a
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	0
Tympanic	0
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**


Hatch size
Preanal length
Length at flexion
Length at transformation
Sequence of fin development
Pigment
• Postflexion larvae >16.4 mm SL
—Along base of soft dorsal fin rays
—Along base of parietal spine
—Along lateral line in caudal peduncle area

Diagnostic characters**PELAGIC JUVENILES**

- Body pigment mainly dorsal to midline
- Opercular blotch present
- Lateral midline pigmented
- Little fin pigment

^aMoulton 1975

Ref: Laroche, in prep.



Figures A–B, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-27 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 14-15-16
Pectoral fin	R: 18-18-19
Anal fin	S: 3-3-3 R: 7-8-9
Gill rakers	U: 9-10-11 L: 24-26-28 (T: 34-36-38)
Lateral line pores	54-60

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesobenthic, 0-375 m
ELH pattern	Ovoviparous, pelagic larvae
Spawning	Season: Nov-Mar; ^a Apr (British Columbia) ^b Area: Mode: Migration:
Fecundity	Range/function: 55,600-915,200 ^a
Age at first maturity	3-4 yr (California) ^a 5 yr (females) ^c
Longevity	28 yr ^d 56 yr (females) ^e 57 yr (males) ^e

^aPhillips 1964^bWestrheim 1975^cWyllie Echeverria 1987^dToole 1982^eR. Mandapat, Wash. Dep. Fish., 7600 Sand Point Way N.E., Seattle, WA 98115-0070, pers. commun., 1 June 1987.^fJuveniles are pelagic at 55-75 mm SL.

Ref: Laroche, in prep.; Laroche and Richardson 1981; Moser and Butler 1987; Washington et al. 1984b.

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	+
Tympanic	+
Inferior infraorbital	3

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	4.5-5.0 mm SL (pre-extrusion)
Preanal length	56-61% SL
Length at flexion	9.9-12.9 mm SL, complete at 14 mm SL
Length at transformation	21.7-30.6 mm SL ^f
Sequence of fin development	Caudal and pectorals, pelvics, dorsal and anal
Pigment	<ul style="list-style-type: none"> • Extrusion larvae: Short series midway along ventral body • Larvae >10 mm SL <ul style="list-style-type: none"> —Beneath dorsal fin —Above and below notochord at tail —Dorsal and ventral margin of caudal peduncle —Moderate on paired fins

Diagnostic characters

- Pigment along dorsal fin base
- Relatively slender body
- No pigment at anal ray bases
- Pigment at tip of notochord

PELAGIC JUVENILES

- Pigment blotch on last few dorsal fin spines
- Body heavily pigmented
- Pectoral fin base, anal fin, and ventral caudal peduncle unpigmented







Figure A, Moser and Butler 1987; B-C, Laroche and Richardson 1981; D, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	12, 8+7, 12-13
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 12-13-13 R: 14-14-15
Pectoral fin	R: 17-18-18
Anal fin	S: 3-3-3 R: 7-8-8
Gill rakers	U: 8-11-12 L: 24-26-27 (T: 34-36-37)
Lateral line pores	49-55

LIFE HISTORY

Range	S. California, 32-34°N, to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesobenthal, 0-549 m
ELH pattern	Ovoviparous, pelagic larvae
Spawning	Season: Nov-Feb; ^a Mar ^b Area: Mode: Migration:
Fecundity	Range/function: 48,000-632,800/ $F = 82721.8 \times L - 323516^c$
Age at first maturity	3-5 yr ^a 7 yr (females) ^d
Longevity	

HEAD SPINES

Preocular	0
Postocular	+
Coronal	0
Supraocular	0
Tympanic	+
Inferior infraorbital	3

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	4.5 mm SL (pre-extrusion)
Preanal length	50-75% SL
Length at flexion	Complete at 10.1 mm SL
Length at transformation	23-27 mm SL ^e
Sequence of fin development	Caudal and pectorals, pelvics, dorsal and anal
Pigment	

- Extrusion larvae: Series along ventral body
- See also *S. melanops* (p. 314)
- Moderate on paired fins
- Over tip of notochord
- Along base of second dorsal fin, developing all along dorsal midline
- Dorsal and ventral margin of caudal peduncle
- At bases of some dorsal and anal fin rays

Diagnostic characters


- See also *S. melanops*
- Number of pectoral fin rays (18)
- Lateral line pores 49-55, usually >50
- Slender caudal peduncle
- Heavy pigment develops along entire dorsal midline
- Pigment at bases of some dorsal and anal fin rays

PELAGIC JUVENILES


- Pigment blotch on last few dorsal fin spines
- Body heavily pigmented, diffuse
- Ventral caudal peduncle pigmented

^aPhillips 1964^bWestrheim 1975^cGunderson et al. 1980^dWyllie Echeverria 1987^eJuveniles are pelagic at 40-50 mm SL.


A



B



C



D




Figure A, DeLacy et al. 1964; B-C, Laroche and Richardson 1980; D, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 3-5-5
Dorsal fin	S: 12-13-14 R: 12-13-14
Pectoral fin	R: 15-16-17
Anal fin	S: 3-3-3 R: 6-6-7
Gill rakers	U: 8-9-9 L: 19-21-22 (T: 28-30-33)
Lateral line pores	34-45

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesobenthic, 25-549 m
ELH pattern	Ovoviparous, pelagic larvae
Spawning	Season: June ^a Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	8 yr (females) ^b
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	+
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	4.1 mm SL
Preanal length	56-63% SL
Length at flexion	7.7-8.8 mm SL
Length at transformation	12.0-18.6 mm SL ^c
Sequence of fin development	Caudal and pectorals, pelvics, dorsal and anal

Pigment

- Extrusion larvae: Series along ventral body
- Lack of body pigment, with development caudal peduncle patch
- Pigmented area at base of long parietal spine
- Pigmented fringes of pectoral and pelvic fins

Diagnostic characters

- Pigmented fringes of pectoral and pelvic fins
- Lack of body pigment
- Long serrate parietal and middle posterior preopercular spines


PELAGIC JUVENILES

- Body pigment diffuse, heavier over gut and on caudal peduncle

^aWestrheim 1975^bWyllie Echeverria 1987^cJuveniles are pelagic to 60 mm SL.

Ref: Laroche, in prep.; Richardson and Laroche 1979; Washington et al. 1984b; Westrheim 1975.


A



\bar{x} = 4.1 mm TL

Long parietal spine


B



Pigmented fringes on pectoral and pelvic fins

Wal 8.0 mm SL


C



Caudal peduncle pigment patch

Wal 13.4 mm SL

D



Wal 41.6 mm SL

Figure A, Westrheim et al. 1968a; B–C, Richardson and Laroche 1979; D, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 13-14-16
Pectoral fin	R: 19-20-22
Anal fin	S: 3-3-3 R: 9-9-10
Gill rakers	U: 12-12-12 L: 31-31-31 (T: 40-44-47)
Lateral line pores	53-64

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesobenthic, 0-350 m
ELH pattern	Ovoviparous, pelagic larvae
Spawning	Season: Nov-Mar; ^a Mar ^b Area: Mode: Migration:
Fecundity	Range/function: 7000-50,000 ^c
Age at first maturity	3 yr (females) ^d
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	0
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION

LARVAE

Hatch size	5.4 mm SL
Preanal length	36%, with development 54% SL
Length at flexion	8-10 mm SL
Length at transformation	~27-30 mm SL ^e
Sequence of fin development	Caudal and pectorals, pelvics, dorsal and anal

Pigment

- Brain (dorsal head on crown)
- Pigment over dorsolateral surface of gut
- Dorsal and ventral postanal midline (see illustrations)

Diagnostic characters

- Most slender *Sebastes* sp. larvae described to date (body depth <25% SL)
- Large larvae (5.4 mm SL at hatching, 8-10 mm SL at flexion)
- Opposing dorsal and ventral postanal midline pigment
- Anterior placement of anus in juveniles/adults
- Large gap between anus and origin of anal fin

PELAGIC JUVENILES

- Elongate shape, large gap between anus and origin of anal fin
- Body pigment diffuse, but myosepta pattern evident

^aPhillips 1964

^bWestrheim 1975


^cHart 1973

^dWyllie Echeverria 1987


^eJuveniles are pelagic at 30-63 mm SL.

Ref: Laroche, in prep.; Moser et al. 1977; Washington et al. 1984b.

A




B



Ventral midline longer than dorsal midline


C



Gap between anus and anal fin origin

Both dorsal and ventral midline equal in length, extending from midbody to just anterior to caudal peduncle

D



E




Figure A, NWAFC original (B. Vinter); B-D, Moser et al. 1977; E, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	12, 8+7, 12-13
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-14 R: 14-15-15
Pectoral fin	R: 18-19-19
Anal fin	S: 3-3-3 R: 7-8-9
Gill rakers	U: 9-11-13 L: 23-26-28 (T: 34-37-38)
Lateral line pores	47-55

LIFE HISTORY

Range	Cent. California, 34-38°N, to Aleutian Is., 51-55°N
Ecology	Epi- and mesobenthic, 0-366 m
ELH pattern	Ovoviparous, pelagic larvae
Spawning	Season: Jan; ^a Feb-Apr ^b Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	7 yr (females) ^c
Longevity	

HEAD SPINES

Preocular	0
Postocular	+
Coronal	0
Supraocular	0
Tympanic	+
Inferior infraorbital	3

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	
Preanal length	50-75% SL
Length at flexion	Complete at 10.6 mm SL
Length at transformation	24-33 mm SL ^d
Sequence of fin development	Caudal and pectorals, pelvics, dorsal and anal
Pigment	

- Extrusion larvae: Series along ventral body
- Larvae >10.6 mm SL
 - Head, on crown and opercle
 - Pigment over dorsolateral surface of gut
 - Dorsal midline heavier than ventral midline
 - Hypural margin
 - Along posterior lateral line, extends anteriorly along notochord with development
 - Paired fin blades moderately pigmented
 - Discrete melanophores at the articulation of several dorsal and anal fin rays

Diagnostic characters

- Distinguished from *S. flavidus* (p. 308) by
- High pectoral fin ray count (usually 19)
 - Fewer lateral line pores (usually <50)
 - Deeper, shorter caudal peduncle

PELAGIC JUVENILES

- Body pigment heavy, diffuse
- Pigment blotch on last few dorsal fin spines
- Lateral midline pigmented

^a Westrheim 1975^b Hart 1973^c Wyllie Echeverria 1987^d Juveniles are pelagic at 40-50 mm SL.







Figure A, NWAFC original (B. Vinter); B–D, Laroche and Richardson 1980.

SCORPAENIDAE *Sebastes melanostomus* (Eigenmann and Eigenmann 1890)

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	9-10, 8+7, 9-11
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 12-13-14
Pectoral fin	R: 18-19-20
Anal fin	S: 3-3-3 R: 6-7-8
Gill rakers	U: 8-X-9 L: 22-22-22 (T: 30-31-34)
Lateral line pores	28-33

LIFE HISTORY

Range	South of southern California to Washington, 46-48°30'N
Ecology	Epi-, meso-, and bathybenthal, 125-768 m
ELH pattern	Ovoviviparous, pelagic larvae
Spawning	Season: Abundance of newborn larvae peaks in February off S. California ^a Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	8 yr (females) ^b
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	+
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION

LARVAE

Hatch size	4.5 mm SL
Preanal length	37% SL increasing with development to 62% SL
Length at flexion	6.2-7.2 mm SL
Length at transformation	~16 mm SL ^c
Sequence of fin development	Caudal and pectorals, pelvics, dorsal and anal

Pigment

- Preflexion larvae
 - Single row postanal ventral midline (~8 spots)
 - Brain
 - Lower jaw, streak along upper jaw
 - Dorsolaterally on gut
 - Pectoral fin
- Flexion larvae: Opercle blotch and upper jaw streak
- Postflexion and pelagic juveniles: Anterior bar; with development, two additional bars form, under second dorsal and on caudal peduncle

Diagnostic characters

- Pigment
 - Jaw, opercle blotch, postanal bars
 - Blades of paired fins heavily pigmented
 - Band of pigment on body centered at about dorsal spine 5-7
- Head spination (see p. 271)
 - Large serrate parietal spine
 - Large weakly serrate preopercle angle spine
- Deep bodied (>35% SL)

PELAGIC JUVENILES

- Banded body pigment extends onto dorsal and anal fins
- Pectoral fin, but not base, pigmented

^aMoser and Ahlstrom 1978

^bWyllie Echeverria 1987

^cLargest pelagic juvenile collected is 46.2 mm SL; smallest demersal juvenile collected is 36.0 mm SL.







Figure A, Moser et al. 1977; B-C, Moser and Ahlstrom 1978; D, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 13-14-15
Pectoral fin	R: 17-18-18
Anal fin	S: 3-3-3 R: 6-7-7
Gill rakers	U: X-X-X L: X-X-X (T: 36-41-42)
Lateral line pores	41-48

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epi- and mesobenthal, 0-274 m
ELH pattern	Ovoviviparous, pelagic larvae
Spawning	Season: Nov-Mar ^a Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	5 yr (females) ^b
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	+
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size 4.3 mm

Preanal length

Length at flexion

Length at transformation

Sequence of fin

development

Pigment

- Extrusion larvae: Short series midway along ventral body
- Postflexion larvae >12.3 mm SL
 - Dorsolateral patch under spinous dorsal
 - Along dorsal and anal fin ray bases
 - Series along ventral midline toward caudal, becoming internal
 - Along hypural margin

Diagnostic characters**PELAGIC JUVENILES**

- Body and head pigment diffuse, heavy
- Pigment blotch on last few dorsal fin spines
- Paired fins pigmented

^aPhillips 1964^bWyllie Echeverria 1987

Ref: Laroche, in prep.






Figure A, Moser et al. 1977; B-C, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-27 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 15-16-17
Pectoral fin	R: 17-18-19
Anal fin	S: 3-3-3 R: 8-9-10
Gill rakers	U: 9-10-12 L: 23-25-29 (T: 32-35-38)
Lateral line pores	47-53

LIFE HISTORY

Range	South of southern California to Aleutian Is., 51-55°N
Ecology	Epi- and mesopelagic, 0-549 m
ELH pattern	Ovoviparous, pelagic larvae
Spawning	Season: Nov-Jan ^a Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	6 yr (females) ^b
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	+
Tympanic	+
Inferior infraorbital	3

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size 5.2 mm; 3.8 mm SL^c

Preal length

Length at flexion

Length at transformation

Sequence of fin

development

Pigment

- Extrusion larvae: Short series along ventral body^c
- Preflexion larvae
 - Spots develop at tip of lower jaw and on pectoral fin blade^c
 - Spots form at junction of cleithra
- Postflexion larvae >12.8 mm SL
 - Along body margins beneath soft dorsal fin and posterior to anal fin
 - Hypural margin
 - Internal and external spots along lateral line

Diagnostic characters**PELAGIC JUVENILES**

- Body and head pigment diffuse, heavy
- Pigment blotch on last few dorsal fin spines

^aWales 1952

^bWyllie Echeverria 1987

^cL. Wold and G. Moreno, Moss Landing Mar. Lab., Moss Landing, CA 95039-0450, pers. commun., 26 July 1988. Drawing (Fig. A) by L. McMasters. Based on results from California Sea Grant Project R/F 115.


Ref: Efremenko and Lisovenko 1970; Laroche, in prep.

A




3.8 mm NL

B




12.8 mm SL

C



21.2 mm SL

D



43.0 mm SL

Figure A, Wold, unpubl.; B–D, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-15 R: 13-14-15
Pectoral fin	R: 14-15-16
Anal fin	S: 3-3-3 R: 8-9-10
Gill rakers	U: 8-8-9 L: 20-21-21 (T: 27-28-32)
Lateral line pores	51-62

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesobenthal, 0-475 m
ELH pattern	Ovoviviparous, pelagic larvae
Spawning	Season: Jan-Feb; ^a Nov-Apr ^b Area: Semi-demersal (25-305 m) ^c Mode: Migration:
Fecundity	Range/function: 20,000 ^d - 2,440,000 ^e (may mature more than one brood per year) ^c
Age at first maturity	3-4 yr ^d 4 yr (females) ^f
Longevity	30 yr ^g 60 yr ^h

^a Westrheim 1975^b Frey 1971^c Moser 1967^d Phillips 1964^e Gunderson et al. 1980^f Wyllie Echeverria 1987^g Hart 1973^h Beamish 1979

Ref: Laroche, in prep.; Moser 1967; Moser et al. 1977.

HEAD SPINES

Preocular	+
Postocular	0
Coronal	0
Supraocular	0
Tympanic	+
Inferior infraorbital	3-4

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	4-5 mm SL
Preanal length	<50%, with development 50-75% SL
Length at flexion	7.2-9.7 mm SL
Length at transformation	~15 mm SL
Sequence of fin development	Caudal and pectorals, pelvics, dorsal and anal

Pigment


- Fringed margins of pectoral and pelvic fins
- Few (6-14) ventral midline melanophores migrate with development to form large patch on caudal peduncle

Diagnostic characters

- Early development of elongate pigmented paired fins
- Midlateral pigment blotch on caudal peduncle
- Gap between anus and anal fin origin

PELAGIC JUVENILES

- Body pigment diffuse, light
- Paired and first dorsal fins heavily pigmented distally



Figures A–D, Moser et al. 1977; E, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 13-14-15
Pectoral fin	R: 16-17-18
Anal fin	S: 3-3-3 R: 7-7-7
Gill rakers	U: 12-14-15 L: 26-28-31 (T: 41-43-44)
Lateral line pores	40-47

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesobenthic, 0-425 m
ELH pattern	Ovoviviparous, pelagic larvae
Spawning	Season: Jan-Mar (Brit. Col.); ^a Nov-Mar (California) ^b Area: Mode: Migration:
Fecundity	Range/function: 260,000-1,897,600 ^c / $F=64221.3 \times L - 2330029$ ^d
Age at first maturity	3-5 yr ^c 9 yr (females) ^e
Longevity	

^a Westrheim 1975^b Moser 1967^c Phillips 1964^d Gunderson et al. 1980^e Wyllie Echeverria 1987^f Larvae are pelagic to 50 mm SL.

Ref: Laroche, in prep.; Moser et al. 1977; Richardson and Laroche 1979; Waldron 1968; Washington et al. 1984b.

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	+
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	3.6-4.0 mm SL
Preanal length	59-64% SL
Length at flexion	<7.8-8.8 mm SL
Length at transformation	12.8-18.4 mm SL ^f
Sequence of fin development	Caudal and pectorals, pelvics, dorsal and anal

Pigment

- Initially: Lower jaw, ventral surface of gut and posterior gut, short dorsal and ventral midline series at ~3/4 BL between myomeres 18 and 24
- Short dorsal and ventral midline series on caudal peduncle
- Lightly pigmented paired fins
- Head: Dorsal surface; with development, patch also on upper part of opercle
- Paired fins have large melanophores on blades

Diagnostic characters

- Pigment: Opercular spot, spots on nape
- Morphology: Deep bodied (~40% SL)
- Spines: Large serrate parietal and third posterior preopercular spine

PELAGIC JUVENILES

- Body pigment banded, mainly above lateral midline
- Pigment blotch on last few dorsal fin spines, little pigment on rest of fins




Figure A, Moser et al. 1977; B–C, Richardson and Laroche 1979; D, Laroche, in prep.

MERISTICS

Vertebrae	Total: 27-27-27 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 14-15-15
Pectoral fin	R: 16-17-17
Anal fin	S: 3-3-3 R: 7-7-7
Gill rakers	U: 10-11-13 L: 26-27-30 (T: 36-38-41)
Lateral line pores	48-55

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthic, 12-274 m
ELH pattern	Ovoviparous, pelagic larvae
Spawning	Season: July ^a Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	0
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size

Preanal length

Length at flexion

Length at transformation

Sequence of fin

development

Pigment

- Extrusion larvae: Series along ventral body
- Postflexion larvae >16.8 mm SL
 - On head dorsally
 - Along dorsal body margin and anal fin ray base
 - Internal and external spots along lateral midline
 - Hypural margin

Diagnostic characters

PELAGIC JUVENILES

- Body elongate, pigment diffuse
- Little fin pigment

^aWestheim 1975

Ref: Laroche, in prep.






Figure A, Westheim et al. 1968b; B-C, Laroche, in prep.

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: X-X-X R: 13-14-15
Pectoral fin	R: 18-19-20
Anal fin	S: 3-3-3 R: 7-7-8
Gill rakers	U: 9-10-11 L: 23-24-27 (T: 30-34-36)
Lateral line pores	47-55

LIFE HISTORY

Range	N. California, 38-42°N, to SE Alaska, 55-59°N
Ecology	Epi- and mesobenthal, 141-366 m
ELH pattern	Ovoviparous, pelagic larvae
Spawning	Season: Apr ^a Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	+
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size
Preanal length
Length at flexion
Length at transformation
Sequence of fin
development

Pigment

- Extrusion larvae: Short series along ventral body
- Postflexion larvae >18.5 mm SL
 - Dorsally on head
 - Along dorsal body margin
 - Hypural margin

Diagnostic characters**PELAGIC JUVENILES**

- Body pigment banded, heavy
- Fins heavily pigmented

^aWestrheim 1975

Ref: Laroche, in prep.






Figure A, Westrheim et al. 1968a; B-C, Laroche, in prep.

MERISTICS

Vertebrae	Total: 27-27-28 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	13-14, 8+7, 12-14
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 13-15-16
Pectoral fin	R: 17-18-19
Anal fin	S: 3-3-3 R: 8-X-9
Gill rakers	U: X-X-X L: X-X-X (T: 32-35-37)
Lateral line pores	49-56

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N ^a
Ecology	Epi- and mesobenthic
ELH pattern	Ovoviparous, pelagic larvae
Spawning	Season: Dec-May (north-central California) ^b Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	3 yr (females, California) ^b
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	+
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	4.3-4.8 mm BL
Preanal length	38-43% BL
Length at flexion	7.2-7.6 mm BL

Length at transformation**Sequence of fin****development****Pigment**

- Increases from extrusion through flexion
- Tip of lower jaw
- Line develops on maxillaries
- Above brain and on nape
- Develops on opercular region
- Short ventral midline series
- Dorsal midline series develops
- Some internal pigment posteriorly above notochord
- Base and blade of pectoral fin


Diagnostic characters**PELAGIC JUVENILES**

- Relatively light, uniform pigment
- Midlateral line pigmented
- Myosepta outlined with pigment

^aNWAFRC meristic database^bWyllie Echeverria 1987

Ref: Moser and Butler 1987.


A



Short line of ventral
midline melanophores

4.8 mm BL

B




Short line of dorsal
midline melanophores

Lower jaw
pigment

Pectoral fin
pigment

6.1 mm BL

C




Nape pigment

Internal notochord pigment

Maxillary pigment

7.6 mm BL

D



36.0 mm SL

Figures A–C, Moser and Butler 1987; D, NWAFC original (B. Vinter).

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 11-12-13
Pectoral fin	R: 16-16-18
Anal fin	S: 3-3-3 R: 6-7-7
Gill rakers	U: 8-10-10 L: 22-23-26 (T: 30-32-35)
Lateral line pores	35-43

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesobenthic, 46-421 m
ELH pattern	Ovoviviparous, pelagic larvae
Spawning	Season: Dec-Mar; ^a Feb ^b Area: Mode: Migration:
Fecundity	Range/function: 13,500-230,000 ^a
Age at first maturity	2-4 yr ^a 2 yr (females) ^c
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	0
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE****Hatch size****Prenatal length****Length at flexion****Length at transformation****Sequence of fin
development****Pigment**

- Extrusion larvae: Series along ventral body, shorter series along posterior dorsal midline
- Postflexion larvae >17 mm SL
—Pigment aligned along epaxial musculature
—Internal and external spots along lateral midline

Diagnostic characters**PELAGIC JUVENILES**

- Body pigment banded
- Median fins pigmented

^aPhillips 1964^bWestrheim 1975^cWyllie Echeverria 1987

Ref: Laroche, in prep.






Figure A, Moser et al. 1977; B-C, Laroche, in prep.

MERISTICS

Vertebrae	Total: 27-27-27 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-13-13 R: 13-14-15
Pectoral fin	R: 16-17-18
Anal fin	S: 3-3-3 R: 7-7-8
Gill rakers	U: 9-10-11 L: 23-25-27 (T: 33-35-37)
Lateral line pores	39-47

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesobenthal, 25-475 m
ELH pattern	Ovoviviparous, pelagic larvae
Spawning	Season: July (British Columbia) ^a Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

HEAD SPINES

Preocular	+
Postocular	+
Coronal	0
Supraocular	0
Tympanic	+
Inferior infraorbital	2

EARLY LIFE HISTORY DESCRIPTION**LARVAE**

Hatch size	4.3 mm SL
Preanal length	50-75% SL
Length at flexion	7.4-9.0 mm SL
Length at transformation	~13.5-20.0 mm SL; 13.7-19.6 mm ^{b,c}

Sequence of fin development	Caudal and pectorals, pelvics, dorsal and anal
-----------------------------	--

Pigment

- Extrusion larvae: Posterior gut, along ventral midline, and on head
- Larvae >7.4 mm SL: A few melanophores along ventral midline, head, and dorsal gut
- Postflexion larvae >12.7 mm SL: Along dorsal midline and fin base, on caudal peduncle

Diagnostic characters

- Moderately pigmented pectorals and pelvics
- Lack of body pigment
- Long, deeply serrate parietal spine and posterior preopercular spine
- Little ventral postanal midline pigment
- Development of dorsal midline pigment (>10 mm SL)
- Large head (>40% SL), deep body (>30% SL)

PELAGIC JUVENILES

- Body pigment diffuse, lateral midline pigmented
- Little fin pigment

^aWestrheim 1975^bWashington et al. 1984b^cJuveniles are pelagic to 65 mm SL, although they may become demersal at 35 mm SL.




Figure A, NWAFC original (B. Vinter); B-C, Laroche and Richardson 1981; D, Laroche, in prep.

MERISTICS *S. alascanus* Bean 1890

Vertebrae	Total: 29-30-31	
	Precaudal: 10-10-11	
	Caudal: 18-20-21	
Branchiostegal rays	7-7-7	
Caudal fin	X, 8+8, X	
Pelvic fin	Thoracic	
	S: 1-1-1	R: 5-5-5
Dorsal fin	S: 14-16-17	R: 8-9-10
Pectoral fin	R: 20-21-23	
Anal fin	S: 3-3-3	R: 3-5-5
Gill rakers	U: 5-6-8	L: 12-14-17

MERISTICS *S. altivelis* Gilbert 1896

Vertebrae	Total: 28-X-29	
	Precaudal: X-X-X	
	Caudal: X-X-X	
Branchiostegal rays	7-7-7	
Caudal fin	X, 8+8, X	
Pelvic fin	Thoracic	
	S: 1-1-1	R: 5-5-5
Dorsal fin	S: 15-X-16	R: 8-X-10
Pectoral fin	R: 22-X-24	
Anal fin	S: 3-3-3	R: 4-X-6
Gill rakers	U: 7-X-9	L: 14-X-17

MERISTICS *S. macrochir* Günther 1880

Vertebrae	Total: 29-29-29	
	Precaudal: X-X-X	
	Caudal: X-X-X	
Branchiostegal rays	7-7-7	
Caudal fin	X, 8+7, X	
Pelvic fin	Thoracic	
	S: 1-1-1	R: 5-5-5
Dorsal fin	S: 15-15-16	R: 8-X-10
Pectoral fin	R: 22-X-23	
Anal fin	S: 3-3-3	R: 5-5-5
Gill rakers	U: 7-X-8	L: 13-X-14

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N ^a
Ecology	Benthals ^b
ELH pattern	Oviparous; gelatinous, pelagic egg masses; ^c pelagic larvae
Spawning	Season: Jan-June (California); ^d spring (Oregon) ^c

EARLY LIFE HISTORY DESCRIPTION**LARVAE** - Genus^c

Preanal length <50%, with development increasing to 50-75% SL

Length at flexion 6.0-7.5 mm SL

Length at transformation 14-20 mm SL^f

Sequence of fin development Pectorals; caudal; dorsal, anal, and pelvics

Pigment - Genus

- Postanal band that disappears with development
- Pectoral fins pigmented at fringes
- Body lacks pigment from ~6-12 mm SL

Diagnostic characters

- <10.0 mm SL: Not separable to species
 - >10.0 mm SL: *S. alascanus* less robust, shorter, and with fewer pectoral rays than *S. altivelis*
- Distinguished from *Melamphaes lugubris* (p. 254) at sizes <5 mm SL by
- Postanal pigment band more posterior, beginning about myomere 20
 - Pelvic fin not precocious in *Sebastolobus* spp.

^a *S. macrochir* found only in the Bering Sea.

^b *S. alascanus* epi-, meso-, and bathybenthal (26-1524 m); *S. altivelis* meso- and bathybenthal (305-1524 m); *S. macrochir* mesobenthal.

^c Percy 1962

^d Moser 1974

^e Larvae of *S. macrochir* are unknown.

^f Juveniles are pelagic to 42-56 mm SL.




Figure A, NWAFC original (B. Vinter); B–G, Moser et al. 1977.

MERISTICS

Vertebrae	Total: 61-63-66 Precaudal: 29-31-33 Caudal: 31-33-34
Branchiostegal rays	6-6-6
Caudal fin	X, 7+7, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 17-22-30 R: 16-18-21
Pectoral fin	R: 14-14-14
Anal fin	S: 2-X-3 R: 15-19-23
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathybenthal, 0 (juveniles) - 2740 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Dec-Apr (California); ^a Sept-Apr (Oregon-Brit.Col.); ^b fall-summer (Bering Sea) ^b Area: Pelagic (175-1450 m) ^b Mode: Migration: To deeper water ^c
Fecundity	Range/function: 100,000 ^a -1,300,000 ^d
Age at first maturity	5-7 yr ^d
Longevity	55 yr ^e

^aPhillips and Imamura 1954^bMason et al. 1983^cKendall and Matarese 1987^dAlton and Webber 1976^eMcFarlane and Beamish 1983

Ref: Ahlstrom and Stevens 1976, Kendall and Matarese 1987.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	2.05-2.10 mm
No. of oil globules	None
Yolk	Homogeneous
Envelope	Smooth, unsculptured
Hatch size	6 mm SL
Incubation time/temp.	
Pigment	• Unpigmented embryo

Diagnostic characters

- Narrow perivitelline space
- Unpigmented embryo with high myomere count and long gut
- No oil globule
- Large size (2.05-2.10 mm)

LARVAE


Preanal length	50-60% SL
Length at flexion	~12.0 mm SL
Length at transformation	
Sequence of fin development	Caudal, pectorals, 2nd dorsal (rays) and anal, 1st dorsal (spines), pelvics
Pigment	• Initially unpigmented • Preflexion larvae develop dorsal and ventral midline pigment • Flexion and later larvae are heavily and uniformly pigmented

Diagnostic characters (see Table 3)

- Slender, heavily pigmented larvae (flexion)
- Long preanal length (50-60% SL)
- >15 mm SL: long pigmented pectoral fins
- Separate dorsal fins (larger larvae)

Larvae of *Erilepis zonifer* (skilfish), an anoplopomatid from the study area, are unknown. The following meristics distinguish them from *A. fimbria*.

Total vertebrae	45-46
Dorsal fin rays	XII-XVI, 15-19
Pectoral fin rays	16-19
Anal fin rays	III, 11-14



Figures A–D, Kendall and Matarese 1987; E–F, Ahlstrom and Stevens 1976.

Hexagrammidae

The greenlings, endemic to the North Pacific Ocean, are composed in our area of nine species in five genera. Most adult greenlings are demersal and occur nearshore; however, larvae commonly occur off the continental shelf. Eggs are demersal and are usually guarded in nests. Development from hatching to juvenile is direct without any marked transformation. An epipelagic juvenile stage occurs in most species. Larvae are heavily pigmented with scattered melanophores over most of the body, especially dorsally. Characters that will distinguish hexagrammids from other heavily pigmented larvae are presented in Table 3. Identification is accomplished using a combination of pigment and meristic characters along with geographic occurrence. *Oxylebius pictus* and *Zaniolepis* spp. larvae have lower vertebral counts, larger pectoral fins with heavier pigment, and differ from other genera in a number of morphological features (e.g., larger heads, longer preanal length, and deeper bodies). *Ophiodon elongatus* larvae can be separated by the presence of a pointed snout and a large terminal mouth. *Pleurogrammus monopterygius* larvae lack pigment on the snout which serves to distinguish them from larvae of *Hexagrammos* spp. Differences among the larvae of the various species of *Hexagrammos* spp. are discussed in Tables 34-35 and illustrated on page 351.

Table 32
Meristic characters of Northeast Pacific hexagrammids. For all species the normal count for branchiostegal rays is 6 and for pelvic fin rays is 1,5 (Kendall and Vinter 1984).

Species	Fin rays											
	Vertebrae			Dorsal			Caudal					Gill rakers
	Precaudal	Caudal	Total	First (spines)	Second (soft rays)	Total anal*	Pectoral	Dorsal		Ventral		
								Secondary	Principal	Principal	Secondary	
<i>Oxylebius pictus</i>	13-15	23-25	36-40	15-17	13-16	14-17	14-17	9	7	6	9	11-14
<i>Zaniolepis frenata</i>	14-15	26-28	40-43	21	12	18-19						
<i>Zaniolepis latipinnis</i>	14	28	42	21-22	11-12	18-20	14	6-8	7	6	7-9	11-12
<i>Ophiodon elongatus</i>	23-24	33-35	56-59	25-28	19-21	21-25	16-18	13-15	7	7	12-14	19-28
<i>Pleurogrammus monopterygius</i>	26-28	32-35	58-63	21-24	24-30	23-28	23-28	16-19	8	11	16-20	22-27
<i>Hexagrammos decagrammus</i>	20-22	33-35	52-57	21-23	22-26	23-26	18-20	12-16	7	9	12-14	15-20
<i>Hexagrammos lagocephalus</i>	20-23	32-34	52-57	20-23	20-25	21-24	18-21	17-22	7	10	15-19	14-18
<i>Hexagrammos octogrammus</i>	18-19	32-35	50-54	18-20	22-25	23-26	18-19	15-17	7	8	14-15	14-17
<i>Hexagrammos stelleri</i>	20-22	31-34	51-56	22-25	18-22	22-25	18-20	16-17	7	8	14-15	16-20

*Anal spines are very weak in *Pleurogrammus* and *Hexagrammos*, therefore only total anal fin elements are given for these taxa. Counts for anal spines in specific taxa are given on the individual text pages.

Table 33
Larval characters that allow distinction between *Oxylebius pictus* and *Zaniolepis* spp. (Kendall and Vinter 1984).

	<i>Oxylebius pictus</i>	<i>Zaniolepis</i> spp.
3-7 mm SL		
Presence of pigment		
On isthmus	No	Yes
Laterally above gut	Less	More
On ventral midline of trunk	More	Less
On tip of snout and onto tip of palate	No	Yes
On internal surface of pectoral fin base	Yes	No
On lower lip	No	Sometimes
Morphology		
Eye size	Smaller (<20% head length)	Larger (>30% head length)
Preanal finfold (5 to 8 mm)	No	Yes
Pectoral fin length	Reaches anus by 8.0 mm	Reaches anus by 5.5 mm
Notochord flexion	7-9 mm	5-7 mm
Dorsal indentation on eye	No	Yes
>7 mm SL		
Spiny scales cover body	No	Yes
Body depth	Stout (<30% SL)	Slender (<25% SL)

MERISTICS

Vertebrae	Total: 36-X-40
	Precaudal: 13-13-15
	Caudal: 23-24-25
Branchiostegal rays	6-6-6
Caudal fin	9, 7+6, 9
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 15-X-17 R: 13-X-17
Pectoral fin	R: 14-X-17
Anal fin	S: 3-X-4 R: 12-X-14
Gill rakers	U: 2-X-5 L: 7-X-8

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Nearshore shelf demersal, intertidal to 49 m
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: May-Aug (Puget Sound); ^a Oct-July (Calif.) ^b Area: Demersal, on or near rocks ^b Mode: Eggs guarded by male; may have three breeding cycles/season ^b Migration:
Fecundity	Range/function: 12,000-28,000/ $F=0.0338 \times L^{2.114}$, L=TL mm (Shilshole break-water, Washington); $F=0.003 \times L^{2.628}$, L=TL mm (Monterey, Calif.) ^b
Age at first maturity	3 yr (females) ^b 2 yr (males) ^b
Longevity	>8 yr ^c

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	4-5 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	54-64% SL
Length at flexion	7-9 mm SL
Length at transformation	16-20 mm SL
Sequence of fin development	Pectorals, caudal, 2nd dorsal (rays) and anal, 1st dorsal (spines) and pelvics

Pigment

- Pectoral fin
- Ventral midline
- Dorsolateral on gut
- Flexion larvae with increased pigment on anterior body

Diagnostic characters


- See Table 33 for characters that allow separation from *Zaniolepis* spp.
- Large pigmented pectoral fins

^aPatten 1980

^bDeMartini 1976

^cFitch and Lavenberg 1975

Ref: Kendall and Vinter 1984.



Figures A–D, Kendall and Vinter 1984.

***Z. frenata* Eigenmann and Eigenmann 1889**

MERISTICS

Vertebrae	Total: 40-X-43
	Precaudal: 14-14-15
	Caudal: 26-28-28
Branchiostegal rays	6-X-7
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 21-21-21 R: 12-12-12
Pectoral fin	R: X-X-X
Anal fin	S: 3-3-3 R: 15-X-16
Gill rakers	U: X-X-X L: 10-X-11

***Z. latipinnis* Girard 1857**

MERISTICS

Vertebrae	Total: 42-42-42
	Precaudal: 14-14-14
	Caudal: 28-28-28
Branchiostegal rays	6-X-7
Caudal fin	6-8, 7+6, 7-9
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 21-X-22 R: 11-X-12
Pectoral fin	R: 14-14-14
Anal fin	S: 3-3-3 R: 15-X-17
Gill rakers	U: 3-3-3 L: 8-X-9

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N ^a
Ecology	Benthalth ^b
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Fall-winter (California) ^c Area: Mode: Migration:
Fecundity	Range/function: 350-6530 (may produce three clutches/season) ^c
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.40-1.56 mm ^d
No. of oil globules	30-50 (red, orange) ^d
Oil globule diameter	0.04-0.12 mm ^d
Yolk	
Envelope	Finely striated ^d
Hatch size	2.5 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	~50% SL
Length at flexion	5-7 mm SL
Length at transformation	
Sequence of fin development	Pectorals, caudal, 2nd dorsal (rays) and anal, 1st dorsal (spines) and pelvics

Pigment

- Laterally on gut
- Ventral midline
- Flexion larvae with increased pigment on body^d

Diagnostic characters

- See Table 33 for characters that allow separation from *Oxylebius pictus*
- Spiny scales


^a*Z. frenata* only to Oregon, 42-46°N.

^b*Z. frenata* epi- and mesobenthic (55-244 m); *Z. latipinnis* nearshore shelf (37-201 m).

^c*Z. latipinnis* only (Goldberg 1980a).

^dData are from unfertilized, hydrated eggs of *Z. frenata* (W. Watson, Marine Ecological Consultants, 531 Encinitas Blvd., Suite 110, Encinitas, CA 92024, pers. commun., 6 Feb. 1988).

Ref: Kendall and Vinter 1984.



Figures A–C, Kendall and Vinter 1984.

MERISTICS

Vertebrae	Total: 56-X-59
	Precaudal: 23-23-24
	Caudal: 33-34-35
Branchiostegal rays	6-X-7
Caudal fin	13-15, 7+7, 12-14
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 25-X-28 R: 19-X-21
Pectoral fin	R: 16-X-18
Anal fin	S: 3-3-3 R: 21-X-25
Gill rakers	U: 5-X-8 L: 16-X-19

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesobenthic, intertidal to 475 m
ELH pattern	Oviparous; demersal, adhesive, attached eggs; pelagic larvae
Spawning	Season: Dec-Apr ^a Area: Intertidal to 19 m below low tide; ^a nest in rocky areas with high current velocities ^b Mode: One or more nests guarded by male ^c Migration: To shallow water ^d
Fecundity	Range/function: 60,000-500,000 ^e / N=0.0002824 × L ^{3.001} ^f
Age at first maturity	2-3 yr ^g
Longevity	20 yr ^h

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	2.24-3.23 mm
No. of oil globules	One
Oil globule diameter	
Yolk	
Envelope	Opaque, thick
Hatch size	7-10 mm SL (9.3 mm SL)
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	~43-55% SL
Length at flexion	11-15 mm SL
Length at transformation	30 mm SL; larvae remain pelagic to 52 mm SL
Sequence of fin development	Caudal and pectorals, 2nd dorsal (rays) and anal, 1st dorsal (spines), pelvics

Pigment

- Head and snout
- Laterally on surface of gut
- Along dorsal midline and ventrolateral body
- Rather uniformly pigmented (flexion)

Diagnostic characters (see Table 3)

- Distinguished from other hexagrammids by
- Pointed snout, protruding lower jaw
 - Large terminal mouth with gape directed upward
 - Longer gut
 - Distribution of pigment on tail

^aLaRiviere et al. 1981

^bGiorgi 1981

^cJewell 1968

^dMiller and Geibel 1973





^ePhillips 1959

^fHart 1967; N=egg number.

^gFrey 1971

^hFitch and Lavenberg 1971

Ref: Kendall and Vinter 1984.



Figures A–D, Kendall and Vinter 1984.

MERISTICS

Vertebrae	Total: 58-X-63 Precaudal: 26-X-28 Caudal: 32-X-35
Branchiostegal rays	5-X-7
Caudal fin	16-19, 8+11, 16-20
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 21-X-24 R: 24-X-30
Pectoral fin	R: 23-X-28
Anal fin	R: 23-X-28
Gill rakers	U: 6-X-8 L: 16-X-19

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthal, intertidal to 575 m
ELH pattern	Oviparous; demersal, attached eggs; pelagic larvae
Spawning	Season: June-Sept (Bering Sea) ^a Area: Mode: Nests guarded by males ^b Migration:
Fecundity	Range/function: 3653-18,694, ^c 43,000 ^a
Age at first maturity	3-4 yr ^a
Longevity	11 yr ^a

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	2.5-2.8 mm
No. of oil globules	Many in group
Oil globule diameter	1.38-1.40 mm
Yolk	
Envelope	Smooth
Hatch size	~8 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	33-50% SL
Length at flexion	14-19 mm SL
Length at transformation	
Sequence of fin development	Caudal, pectorals, dorsal and anal, pelvics
Pigment	<ul style="list-style-type: none"> • Midbrain and internally on hindbrain • Dorsolateral surface of gut • Postanal: Dorsal midline along body length to last myomere; ventral midline extends only from mid-body to last myomere • Internal above and below notochord

Diagnostic characters (see Table 3)





- Distinguished from other hexagrammids by
- Pigment pattern (unpigmented snout)
 - Morphology: Larger eye, shorter snout

^aGorbunova 1962

^bKendall and Vinter 1984

^cLee 1985

Ref: Kendall and Vinter 1984.



Figures A–D, Kendall and Vinter 1984.

Table 34
Guide for identifying Northeast Pacific species of
Hexagrammos based on meristic characters
 (Kendall and Vinter 1984).

Based on principal ventral caudal fin rays*	
Principal ventral caudal rays	8
precaudal vertebrae	20-22 = <i>H. stelleri</i>
precaudal vertebrae	18-19 = <i>H. octogrammus</i>
Principal ventral caudal rays	9 = <i>H. decagrammus</i>
Principal ventral caudal rays	10 = <i>H. lagocephalus</i>
Based on extremes of meristic characters	
Dorsal fin spinous rays	18-19 = <i>H. octogrammus</i>
	20-23 = indeterminate
	24-25 = <i>H. stelleri</i>
Dorsal fin soft rays	18-19 = <i>H. stelleri</i>
	20-25 = indeterminate
	26 = <i>H. decagrammus</i>
Total anal fin elements	21 = <i>H. lagocephalus</i>
	22-26 = indeterminate
Precaudal vertebrae	18-19 = <i>H. octogrammus</i>
	20-23 = indeterminate
Caudal vertebrae	31 = <i>H. stelleri</i>
	32-35 = indeterminate
Total vertebrae	50 = <i>H. octogrammus</i>
	51-57 = indeterminate

*Prior to completion of caudal fin development, the principal ventral caudal fin ray count can be one less in larvae.

Table 35
Pigmentation characteristics that distinguish larvae <30 mm of the four species of
Hexagrammos in the Northeast Pacific
 (Kendall and Vinter 1984, in part).^a


Species	Pigment area			
	Postanal ventral midline	Isthmus	Notochord tip	Internal pectoral fin base ^b (>17 mm)
<i>H. stelleri</i>	Absent until ~15 mm, then starts to form posteriorly (on caudal peduncle), later (>23 mm) forms along entire base of anal fin.	Absent at hatching. Anterior half gradually becomes pigmented starting at 10 mm.	Absent	Present
<i>H. decagrammus</i>	Absent until ~13 mm, then starts to form along anal fin base near its origin. By 18 mm along anal fin base to caudal peduncle. Spots more numerous and smaller than on <i>H. lagocephalus</i> .	About four equal-sized and -spaced spots form on each side making a "V." Present from ~9 mm on.	Absent at hatching but forms by ~10 mm.	Inconsistently present
<i>H. lagocephalus</i>	Present throughout development. Particularly dense on caudal peduncle. Spots along anal fin base tend not to touch each other. Spots uneven in spacing and size to create irregular line. Few spots anterior to origin of anal fin.	An anterior medial spot and about five equal-sized and -spaced spots present from ~9 mm on.	Present	Absent
<i>H. octogrammus</i>	Present throughout development. Extends from anus to base of caudal fin. Dense on caudal peduncle. Spots tend to touch each other to create a continuous straight line.	Present; anteriormost and posteriormost spots larger than others. Anterior medial spot seen at 10 mm, up to five on each side of isthmus seen at 12 mm. Spacing uneven.	Present	Present

^aPigmentation characteristics discussed in this table were based on larvae collected in the Gulf of Alaska. Variation in some of these features is known to occur in larvae from other areas.


^bOne to a few spots occur near the dorsal edge of the internal surface of the pectoral fin base.

VENTRAL VIEW OF *HEXAGRAMMOS* SPP. LARVAE (16-18 mm SL)
DEPICTING PIGMENT ON THE LOWER JAW, ISTHMUS, AND POSTANAL VENTRAL MIDLINE


A *Hexagrammos stelleri*




B *Hexagrammos decagrammus*



C *Hexagrammos lagocephalus*



D *Hexagrammos octogrammus*



MERISTICS

Vertebrae	Total: 52-X-57 Precaudal: 20-21-22 Caudal: 33-X-35
Branchiostegal rays	6-6-6
Caudal fin	12-16, 7+9, 12-14
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 21-X-23 R: 22-X-26
Pectoral fin	R: 18-X-20
Anal fin	R: 23-X-26
Gill rakers	U: 3-X-5 L: 9-X-14

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Intertidal, nearshore, 0-46 m
ELH pattern	Oviparous; demersal, attached eggs; pelagic larvae
Spawning	Season: Oct-Nov ^a Area: Mode: Egg masses on rocks ^a Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	7-9 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	Much <50% before flexion
Length at flexion	12-18 mm SL
Length at transformation	Epipelagic to 50 mm SL
Sequence of fin development	Caudal, pectorals, 2nd dorsal (rays) and anal, 1st dorsal (spines), pelvics
Pigment	<ul style="list-style-type: none"> • Heavily pigmented • Postanal ventral midline absent until ~13 mm SL, then starts near anal fin origin; reaches caudal peduncle by 18 mm SL

Diagnostic characters (see Table 3 and tables this section)

- Principal caudal fin ray count in juveniles (7+9)


^aMarliave 1975a

Ref: Kendall and Vinter 1984.


A




B




C



D



E



Figures A–E, Kendall and Vinter 1984.

MERISTICS

Vertebrae	Total: 52-X-57 Precaudal: 20-20-23 Caudal: 32-33-34
Branchiostegal rays	6-6-6
Caudal fin	17-22, 7+10, 15-19
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 20-X-23 R: 20-X-25
Pectoral fin	R: 18-X-21
Anal fin	R: 21-X-24
Gill rakers	U: 4-X-5 L: 9-X-11

LIFE HISTORY

Range	Cent. California, 34-38°N, to Bering Sea, 54-66°N
Ecology	Intertidal, nearshore
ELH pattern	Oviparous; demersal, adhesive, attached eggs; pelagic larvae
Spawning	Season: June-Aug (Aleutian Is.); ^a June-Sept (west. Pacific) ^b Area: On rocks or algal holdfasts in areas of strong currents ^a Mode: Intermittent, males guard nests ^c Migration:
Fecundity	Range/function: 14,400-103,000 ^b
Age at first maturity	3-4 yr ^b
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	7-9 mm SL
Incubation time/temp.	30 d/6-10°C
Pigment	

Diagnostic characters

LARVAE

Preanal length	Much <50% before flexion
Length at flexion	~16 mm SL
Length at transformation	
Sequence of fin development	Caudal, pectorals, 2nd dorsal (rays) and anal, 1st dorsal (spines), pelvics
Pigment	<ul style="list-style-type: none"> • Heavily pigmented • Postanal ventral midline present throughout development, particularly heavy on caudal peduncle; spots create irregular line

Diagnostic characters (see Table 3 and tables this section)

- Principal caudal fin ray count in juveniles (7+10)


^a Simenstad 1971

^b Gorbunova 1962


^c Kendall and Vinter 1984

Ref: Kendall and Vinter 1984.


A




B



C



D



Figures A–D, Kendall and Vinter 1984.

MERISTICS

Vertebrae	Total: 50-X-54
	Precaudal: 18-X-19
	Caudal: 32-X-35
Branchiostegal rays	6-6-6
Caudal fin	15-17, 7+8, 14-15
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 18-X-20 R: 22-X-25
Pectoral fin	R: 18-X-19
Anal fin	R: 23-X-26
Gill rakers	U: 4-X-5 L: 11-X-12

LIFE HISTORY

Range	Brit. Col., 48°30'-55°N, to Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal
ELH pattern	Oviparous; demersal, attached eggs; pelagic larvae
Spawning	Season: Area: Mode: Nests guarded by males ^a Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS^b

Diameter	1.75-2.10 mm
No. of oil globules	Many
Oil globule diameter	0.8 mm
Yolk	
Envelope	
Hatch size	7-9 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	Much <50% before flexion
Length at flexion	~16 mm SL
Length at transformation	
Sequence of fin development	Caudal, pectorals, 2nd dorsal (rays) and anal, 1st dorsal (spines), pelvics

Pigment

- Most heavily pigmented *Hexagrammos* sp.
- Postanal ventral midline present throughout development, forms continuous line from anus to base of caudal fin

Diagnostic characters (see Table 3 and tables this section)


- Principal caudal fin ray count in juveniles (7+8)

^aKendall and Vinter 1984


^bData are from Gorbunova (1962) as cited in Washington et al. (1984b).

Ref: Kendall and Vinter 1984.


A




B




C



D



E



Figures A–E, Kendall and Vinter 1984.

MERISTICS

Vertebrae	Total: 51-X-56 Precaudal: 20-X-22 Caudal: 31-X-34
Branchiostegal rays	6-6-6
Caudal fin	16-17, 7+8, 14-15
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 22-X-25 R: 18-X-22
Pectoral fin	R: 18-X-20
Anal fin	R: 22-X-25
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Oregon, 42-46°N, to Chukchi Sea, north of 66°N
Ecology	Nearshore shelf demersal, intertidal to 175 m
ELH pattern	Oviparous; demersal, attached eggs; pelagic larvae
Spawning	Season: Feb ^a or Apr ^b (British Columbia); reported summer to fall at extremes of range in Pacific Ocean ^c Area: Mode: May be deposited on rocks ^a Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	7-9 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	Much <50% before flexion
Length at flexion	~15 mm SL
Length at transformation	
Sequence of fin development	Caudal, pectorals, 2nd dorsal (rays) and anal, 1st dorsal (spines), pelvics

Pigment

- Most lightly pigmented *Hexagrammos* sp.
- Postanal ventral midline absent until ~15 mm SL, then starts on caudal peduncle; by ~23 mm SL all along anal fin base

Diagnostic characters (see Table 3 and tables this section)


- Principal caudal fin ray count in juveniles (7+8)

^aClemens and Wilby 1961

^bHart 1973

^cGorbunova 1962

Ref: Kendall and Vinter 1984.



Figures A–E, Kendall and Vinter 1984.

COTTIDAE

The sculpins are represented by over 100 species within 45 genera in the study area. Most species are benthic as juveniles and adults and planktonic as larvae. Adults commonly occur in nearshore and intertidal regions throughout the study area. Cottids generally produce demersal eggs that are guarded in shallow water. Although larval cottids are commonly collected during ichthyoplankton surveys, the early life histories of most species are poorly known and larval descriptions are available for fewer than half the species. The cottid section is arranged according to the phenetic groups first described by Richardson (1981a) and recently updated by Washington et al. (1984a).¹ Preceding the taxonomic sections, the groups are briefly described, representative taxa from each group are illustrated, and a table of meristic characters is provided (Table 36). Betsy Washington provided vertebral counts for several genera based on her osteological studies of the cottids.² In most cases (except for *Clinocottus acuticeps* where $n=3$), counts were taken from one specimen.

¹Psychrolutidae (sensu J. Nelson 1984) is included within Cottidae.

²B.B. Washington, NMFS Natl. Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560, pers. commun., 8 Dec. 1986.

Table 36
Larval information, numbers of species, and meristic characters of genera of cottids from the Northeast Pacific Ocean.
Counts are from Washington et al. (1984a) unless otherwise noted.

Genera	Larval ^a group	Number of species		Vertebrae	Fins				
		Total ^b	With larval descriptions		Dorsal				
					Spines	Soft rays	Anal	Pectoral	Pelvic
<i>Archaulus</i>		1			IX-X	28-29	22-23	16	I, 3
<i>Artediellichthys</i>		1			VII-IX	12-13	9-11	21-23	I, 3
<i>Artedielliscus</i>		1							
<i>Artediellus</i>		5		28-30	VI-IX	11-14	10-14	20-24	I, 3
<i>Artedius</i>	Art ^c	6	4	30-35	VII-X	14-18	10-14	13-17	I, 2-3
<i>Ascelichthys</i>	Myo	1		33-36	VII-X	17-19	13-16	16-18	0
<i>Asemichthys</i>		1		33-35	IX-XI	14-16	15-16	16-18	I, 3
<i>Blepsias</i>	Hem	2	1	37-39	VI-IX	20-24	18-21	11-17	I, 3
<i>Chitonotus</i>	Myo	1		35-36	VIII-XI	14-17	14-17	16-18	I, 2-3
<i>Clinocottus</i>	Art	4	4	31-34	VII-X	13-17	9-13	12-15	I, 3
<i>Cottus</i>	Lep	2	1	34-39	VII-XI	16-21	12-18	13-17	I, 4
<i>Dasycottus</i>	Mal	1		34-35	VIII-XI	13-16	12-16	22-26	I, 3
<i>Enophrys</i>	Myo	3	1	29-35	VII-IX	9-15	8-13	15-19	I, 2-3
<i>Eurymen</i>		1		38	VIII	21-23	15-17	25-26	I, 3
<i>Gilbertidia</i> ^d	Psy	1		33-35	VII-VIII	18-19	12-15	14-17	I, 3
<i>Gymnocanthus</i>	Myo	4	1-2	35-40	IX-XII	13-18	14-20	15-21	I, 3
<i>Hemilepidotus</i>	Hem-Sco	6	5	35-39	X-XII	17-22	13-19	15-19	I, 3-4
<i>Hemitriperus</i>	Hem	2	1	38-41	XI-XIX	11-14	12-15	18-22	I, 3
<i>Icelinus</i>	Myo	5	1	34-39	IX-XII	14-18	11-17	14-19	I, 2
<i>Icelus</i>	Myo	6		37-42	VII-X	17-25	13-20	15-20	I, 3
<i>Jordania</i>		1		46-48	XVII-XVIII	15-18	22-24	13-15	I, 4-5
<i>Leptocottus</i>	Lep	1		35-39	VI-VIII	15-20	15-20	17-20	I, 4
<i>Malacocottus</i>	Mal	2-3	1	30-33	VIII-X	12-15	9-13	19-23	I, 3
<i>Megalocottus</i>		2			VIII-X	12-15	11-13	16-18	I, 3
<i>Microcottus</i>		1		32-34	VII-IX	12-14	10-12	14-17	I, 3
<i>Myoxocephalus</i>	Myo	9-10	1-2	34-46	VIII-XII	10-20	8-17	14-19	I, 3
<i>Nautichthys</i>	Hem	3	1	35-41	VII-X	19-30	14-21	13-17	I, 3
<i>Oligocottus</i>	Art	3	2	33-37	VII-X	15-20	12-15	12-15	I, 3
<i>Paricelinus</i>	Myo	1		42-43	XII-XIII	19-20	23-24	14-15	I, 5
<i>Phallocottus</i>		1			X-XII	22-24	22-25	14-16	I, 3
<i>Porocottus</i>		2		34-38	VIII-X	13-18	11-18	13-19	I, 3
<i>Psychrolutes</i>	Psy	2	1	33-37	VIII-XII	12-20	10-14	19-26	I, 3
<i>Radulinus</i>	Myo	2	2	38-40	VIII-XI	20-23	21-25	17-20	I, 3
<i>Rhamphocottus</i>	Rha	1		26-28	VII-IX	12-14	6-9	14-18	I, 3-4
<i>Scorpaenichthys</i>	Hem-Sco	1		35-37	VIII-XII	15-19	11-14	14-16	I, 4-5
<i>Sigmistes</i>		2		34-36	VIII-X	19-26	14-20	13-15	I, 3
<i>Stelgistrum</i>		2		36	VIII-IX	17-19	12-14	14-16	I, 3
<i>Sternias</i>		1		44-46	X-XI	22-24	22-24	16-18	I, 3
<i>Silegicottus</i>		1			IX	19	17	18	I, 3
<i>Synchirus</i>	Myo	1		38-39	VIII-X	19-21	18-21	21-24	I, 3
<i>Taurocottus</i>		1			XI	15-16	12-13	19	I, 3
<i>Thecopterus</i>		1			X	14	11	20	I, 2
<i>Thyriscus</i>		1		38-39	X	21	17	15	I, 3
<i>Triglops</i>		6	1?	45-54	IX-XIII	21-32	20-32	15-22	I, 3
<i>Zesticelus</i>	Myo	1		25-26	V-VII	10-13	8-11	19-21	I, 2-3

^a Art - *Artedius*

Hem - *Hemitriperus*

Hem-Sco - *Hemilepidotus-Scorpaenichthys*

Lep - *Leptocottus*

Mal - *Malacocottus*

Myo - *Myoxocephalus*

Psy - *Psychrolutes*

Rha - *Rhamphocottus*

^bLimited to study area.

^c*Artedius meanyi* is placed in *Myoxocephalus* group.

^dB.B. Washington, NMFS Natl. Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560, pers. commun., 8 Dec. 1986.

COTTIDAE

PHENETIC GROUPS

Rhamphocottus (Rhamphocottus) Larvae are extremely deep-bodied with a long snout-anus length. Melanophores develop uniformly over the body except on the caudal peduncle and ventral gut surface. Small prickles develop all over the body by 9-10 mm, and only one preopercular spine develops.

Hemilepidotus-Scorpaenichthys (Hemilepidotus, Scorpaenichthys) Larvae are relatively long and slender at hatching with moderately long guts (40-60% SL) and rounded snouts. They become increasingly deep-bodied. Larvae are relatively heavily pigmented. Four prominent preopercular spines develop.


Myoxocephalus (Artemius meanyi, Ascelichthys, Chitonotus, Enophrys, Gymnocanthus, Icelinus, Icelus, Myoxocephalus, Paricelinus, Radulinus, Synchronus, Triglops) Larvae are generally slender-bodied with pointed snouts. Pigment is variable, but most members have heavy pigment on dorsal surface of the gut, nape, and along the postanal ventral midline. Larvae develop four preopercular spines and a distinct bony preopercular shelf. Parietal, nuchal, supracleithral, posttemporal, and occasionally postocular spines develop.

Artemius (Artemius other than A. meanyi; Clinocottus, Oligocottus) Larvae are stubby-bodied with a slightly humped appearance at the nape. Snouts are rounded and guts trail distinctively below the ventral body midline. Some species have gut diverticula. Pigment is relatively light, occurring on the nape, over the gut, and along the postanal ventral midline. Larvae develop a unique preopercular spine pattern with 6-24 spines (enlarged on illustrations).

PHENETIC GROUPS


RHAMPHOCOTTUS

A *Rhamphocottus richardsoni*




HEMILEPIDOTUS – SCORPAENICHTHYS

B *Hemilepidotus spinosus*




MYOXOCEPHALUS

C *Ascelichthys rhodorus*




D *Myoxocephalus G*



ARTEDIUS

E *Artedius harringtoni*



Figures A–B, E, Richardson and Washington 1980; C, Matarese and Marliave 1982; D, NWAFC original (B. Vinter).

COTTIDAE

Psychrolutes (*Gilbertidia*, *Psychrolutes*) Larvae are tadpole-shaped with large rounded heads tapering toward the tail. They possess an outer layer of flabby skin. The head, nape, gut, and pectoral fins are pigmented. No postanal ventral midline melanophores are present, but pigment is added laterally with development. Head and preopercular spines are absent.

Malacocottus (*Dasycottus*, *Malacocottus*) Similar to *Psychrolutes* group but larvae develop 4-5 preopercular spines.


Leptocottus (*Cottus*, *Leptocottus*) Larvae are relatively slender with rounded snouts and moderately short guts. Pigment is light on the postanal body. Other pigment occurs on nape, over gut, and widely spaced along the postanal ventral midline. Four weak preopercular spines develop but other head spines are lacking.

Hemitripteris (*Blepsias*, *Hemitripteris*, *Nautichthys*) Newly hatched larvae are elongate and slender, becoming deeper with development. *Nautichthys* larvae have long pigmented pectoral fins. Pigmentation is heavy, covering the body except for the caudal peduncle. Pigment extends into the dorsal and ventral finfolds. Larvae develop four prominent preopercular spines.

PHENETIC GROUPS


PSYCHROLUTES

A *Psychrolutes paradoxus*




MALACOCOTTUS

B *Malacocottus zonurus*




LEPTOCOTTUS

C *Leptocottus armatus*



HEMITRIPTERUS

D *Hemitripterus villosus*



Figures A–B, Richardson and Bond 1980; C, Richardson and Washington 1980; D, Okiyama and Sando 1976.

MERISTICS

Vertebrae	Total: 26-26-28 Precaudal: 12-12-12 ^a Caudal: 15-15-15 ^a
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic S: 1-1-1 R: 3-3-4
Dorsal fin	S: 7-8-9 R: 12-12-14
Pectoral fin	R: 14-15-18
Anal fin	R: 6-7-9
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthic, intertidal to 274 m
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Aug-Oct (California); ^b winter (British Columbia) ^c Area: Nearshore (20 m) ^d Mode: Egg masses guarded by females ^d Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	2.5-2.8 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	Orange; ^c white ^d
Hatch size	6-7 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	>50% SL
Length at flexion	6.9-10.0 mm NL
Length at transformation	14-15 mm SL
Sequence of fin development	Dorsal, anal, and pectorals followed by caudal and pelvics

Pigment

- Larvae are uniformly covered with melanophores except for caudal peduncle and the ventral surface of gut

Diagnostic characters (see Table 3)

- Morphology: Deep-bodied, very long snout-to-anus length
- Uniformly heavily pigmented
- Presence of pigmented preanal finfold
- Prickles develop over most of body by 9-10 mm SL
- Spines: Only one preopercular (small spiny projections appear along the preopercular margin); others include parietal, nuchal, supracleithral, posttemporal, and postocular


^aB.B. Washington, NMFS Natl. Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560, unpubl.

^bFitch and Lavenberg 1975

^cHart 1973

^dGarrison and Miller 1982

Ref: Blackburn 1973, Marliave 1975a, Richardson and Washington 1980, Saruwatari et al. 1987, Washington et al. 1984b.



Figures A–C, Richardson and Washington 1980.

Table 37


Selected postanal pigmentation characters useful in distinguishing preflexion and flexion *Hemilepidotus* larvae (Matarese and Vinter 1985). Larvae of *H. papilio* are unknown. All lengths are mm SL.

Taxon	Pigmentation characters							Diagnostic
	Dorsal midline	Dorso-lateral	Above notochord (internal)	Below notochord (internal)	Ventro-lateral	Ventral midline	Caudal region	
<i>Hemilepidotus spinosus</i>	By 5 mm, a continuous line from head to posteriormost myomere, becoming heavier	By 6 mm, becoming heavier	By 8-9 mm, along length of body	Not obvious	By 6 mm, becoming heavier	From anus to posteriormost myomere, >15 melanophores	Sparse, ventral midline continuous	Lateral, ventral midline
<i>H. hemilepidotus</i>	Until 7 mm, unpigmented area between myomeres 4-11 becoming moderately heavy	By 11 mm, moderate	By 6-7 mm, along length of body	By 8-9 mm, begins posterior to anus	By 11 mm, moderate	Begins 9 myomeres after anus, <15 melanophores	None	Lateral, lack of caudal pigment
<i>H. jordani</i> ^a	Similar to <i>H. hemilepidotus</i> but not as heavy	None	Similar to <i>H. hemilepidotus</i>	Similar to <i>H. hemilepidotus</i>	Some internal only	By flexion, a few internal melanophores	None	Lack of lateral and caudal pigment
<i>H. zapus</i>	Similar to <i>H. hemilepidotus</i> but not as heavy	None	By 6 mm, along length of body	By 8 mm, incomplete; begins posterior to anus	Some internal only	Begins 11 myomeres after anus, <15 melanophores	Ventral midline continuous, above and below urostyle	Urostyle
<i>H. gilberti</i> ^b	Pigment begins 8-9 myomeres after anus	None	None	None	None	Begins 10 myomeres after anus, <15 melanophores	None	Unpigmented area along dorsal midline, lack of lateral pigment

^a Preflexion larvae of *H. hemilepidotus* and *H. jordani* cannot presently be separated in samples from areas where they co-occur.

^b Specimens were not available; description based on figures in Hattori (1964).

POSTANAL PIGMENT PATTERNS OF *HEMILEPIDOTUS* SPP. LARVAE




Figures A–C, Matarese and Vinter 1985.

Table 38


Characters useful in distinguishing pelagic juveniles of *Hemilepidotus* spp. (Peden 1978, in part). Specimens of *H. papilio* and *H. gilberti* have not been collected from the study area and are not included here.

Character	<i>H. hemilepidotus</i>	<i>H. jordani</i>	<i>H. spinosus</i>	<i>H. zapus</i>
Pectoral fin rays	16(15-17)	18(17-19)	15-16(14-16)	16(15-17)
Total soft fin rays; dorsal, anal, and both pectoral fins	63-68	71-78	63-66	67-76
Lateral line pores	65(59-68)	64(59-68)	63(57-67)	52(47-58)
Vertebrae	35-37	37-39	36	37-38
Horizontal rows in ventral scale band	6 or 7	~8	~4	8 or 9
Gill membranes fused to isthmus so as to form a free fold posteriorly	Yes	Yes	No free fold posteriorly	Yes
Horizontal rows in dorsal scale band	≤5	≤5	≥6	≤5


PELAGIC JUVENILES

A *H. hemilepidotus*


19.0 mm SL

B *H. jordani*

18.4 mm SL

C *H. spinosus*

19.0 mm SL

D *H. zapus*

22.8 mm SL

Figure A, C, Richardson and Washington 1980; B, NWAFC original (B. Vinter); D, Matarese and Vinter 1985.

MERISTICS

Vertebrae	Total: 36-X-38
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 4-4-4
Dorsal fin	S: 11-X-12 R: 17-X-22
Pectoral fin	R: 15-X-17
Anal fin	R: 14-X-19
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal
ELH pattern	Oviparous, eggs probably demersal, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE

Preanal length <50% SL increasing with development to 50% SL
Length at flexion ~7-12 mm SL
Length at transformation
Sequence of fin development
Pigment
• Head and dorsal gut
• With development, nape, upper body, and along dorsal midline


Diagnostic characters (see Tables 3 and 37)

- Unpigmented area along dorsal midline in preflexion larvae
- Lack of lateral pigment throughout larval period

A


Pigment
on crown

Lack of postanal melanophores
in preflexion larvae



7.1 mm SL


B



11.6 mm SL


Nape and upper body pigment

C



19.2 mm SL

D



32.5 mm SL

Figures A–D, Hattori 1964. Identification of Figure B is questionable as only 30 myomeres are illustrated.

MERISTICS

Vertebrae	Total: 35-36-37 Precaudal: 12-12-12 ^a Caudal: 24-24-24 ^a
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic S: 1-1-1 R: 3-4-4
Dorsal fin	S: 10-11-13 R: 18-19-20
Pectoral fin	R: 15-16-17
Anal fin	R: 13-15-16
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Cent. California, 34-38°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthic, intertidal to 275 m
ELH pattern	Oviparous; demersal, adhesive, guarded eggs; pelagic larvae
Spawning	Season: Oct-Jan ^b Area: Shallow water on rocks or pilings in areas with high current velocities ^b Mode: Guarded by female, male, or both ^b Migration:
Fecundity	Range/function: 59,000-126,000 ^b
Age at first maturity	4 yr ^b
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.5-1.6 mm
No. of oil globules	One
Oil globule diameter	0.31-0.56 mm
Yolk	
Envelope	
Hatch size	5-6 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	34% SL increasing with development to >50% SL
Length at flexion	<9 mm SL
Length at transformation	>19-23 mm SL
Sequence of fin development	
Pigment	<ul style="list-style-type: none"> • Dorsal pigment on head and gut • Break in dorsal midline pigment above gut (preflexion) • Short posteriorly placed ventral midline series • Lateral pigment on postanal body

Diagnostic characters (see Tables 3 and 37-38)


- Pigment
 - Lack of pigment around urostyle
 - Presence of dorso- and ventrolateral pigment

^aB.B. Washington, NMFS Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560, unpubl.


^bGarrison and Miller 1982

Ref: Richardson and Washington 1980, Washington et al. 1984b.


A



B




C



A notch after the 3rd dorsal spine, not shown here, is visible on all our specimens of this size

D



Figures A–D, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 37-38-39
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-4-4
Dorsal fin	S: 10-11-12 R: 17-21-22
Pectoral fin	R: 17-18-19
Anal fin	R: 16-17-18
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	SE Alaska, 55-59°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthal, 25-525 m
ELH pattern	Oviparous, eggs probably demersal, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE^a

Preanal length
Length at flexion
Length at transformation
Sequence of fin
development
Pigment

- Similar to *H. hemilepidotus* except generally less pigmented (see p. 374)
- Ventrolateral pigment internal only


Diagnostic characters (see Tables 3 and 37-38)

- Lack of lateral and caudal pigment
- Distinguished from *H. hemilepidotus* postflexion larvae by
- Less dorso- and ventrolateral pigment (see figure, p. 371)
- Delayed development of postocular and parietal spines


^a Preflexion larvae of *H. hemilepidotus* and *H. jordani* cannot presently be separated in samples from areas where they co-occur.

Ref: Matarese and Vinter 1985.

A Postocular and parietal spines
not developed by this size



B



Figures A–B, NWAFC originals (B. Vinter).

COTTIDAE

(*Hemilepidotus-Scorpaenichthys* Group) *Hemilepidotus spinosus* (Ayres 1855)

MERISTICS

Vertebrae	Total: 35-36-37
	Precaudal: 12-12-12
	Caudal: 23-24-25
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 4-4-4
Dorsal fin	S: 10-11-11 R: 18-20-20
Pectoral fin	R: 14-15-16
Anal fin	R: 14-15-16
Gill rakers	U: 2-2-2 L: 5-X-8

LIFE HISTORY

Range	S. California, 32-34°N, to SE Alaska, 55-59°N
Ecology	Nearshore shelf demersal, intertidal to 97 m
ELH pattern	Oviparous, eggs probably demersal, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	~5 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters


LARVAE

Preanal length	38% SL increasing with development to >50% SL
Length at flexion	7.6-10.0 mm SL
Length at transformation	19 mm SL
Sequence of fin development	
Pigment	

- Dorso- and ventrolateral at 6 mm SL becoming heavier with development
- Preflexion larvae (>5 mm SL) have continuous line of dorsal midline pigment extending from head to posteriormost myomere

Diagnostic characters (see Tables 3 and 37-38)

- Pigment: Lateral, ventral midline
- Preflexion: Continuous pigment along dorsal and ventral body midline
- Flexion and larger: Heavy concentration of lateral pigment



Figures A–D, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 37-38-38
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 4-4-4
Dorsal fin	S: 11-11-12 R: 18-20-22
Pectoral fin	R: 15-15-17
Anal fin	R: 16-17-17
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Aleutian Is., 51-55°N, to Chukchi Sea, north of 66°N
Ecology	Nearshore shelf demersal
ELH pattern	Oviparous, eggs probably demersal, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	~4.3 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE


Preanal length	<50% SL
Length at flexion	Complete at 13 mm SL
Length at transformation	>22.8 mm SL
Sequence of fin development	
Pigment	

- Similar to *H. hemilepidotus* except generally less pigmented (see p. 374)
- Preflexion larvae have pigment above and below urostyle

Diagnostic characters (see Tables 3 and 37-38)

- Pigment above and below urostyle

See also pelagic juvenile figure of *H. hemilepidotus* and *H. jordani* (p. 371)



Figures A–D, Matarese and Vinter 1985.

MERISTICS

Vertebrae	Total: 35-36-37 Precaudal: 13-15-16 Caudal: 20-21-23
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic S: 1-1-1 R: 4-5-5
Dorsal fin	S: 8-11-12 R: 15-18-19
Pectoral fin	R: 14-15-16
Anal fin	R: 11-13-14
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59°N
Ecology	Nearshore shelf demersal, intertidal to 76 m
ELH pattern	Oviparous; demersal, adhesive, attached eggs; pelagic larvae
Spawning	Season: Nov-Mar (California); ^a Jan-May (British Columbia) ^b Area: In rocky crevices ^c or on algae ^d Mode: Migration:
Fecundity	Range/function: 49,000-98,000 (may spawn twice in one season) ^a
Age at first maturity	3-4 yr (females) ^a 2-3 yr (males) ^a
Longevity	13 yr ^a

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.4-1.9 mm
No. of oil globules	One large with 1-4 smaller ones
Oil globule diameter	0.2-0.3 mm ^e
Yolk	Homogeneous
Envelope	Thick, translucent
Hatch size	4-6 mm SL; ^f 3.1-4.8 mm SL ^e (yolk absorbed by 6.5 mm SL)
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	45-50% SL
Length at flexion	~7.5-8.7 mm SL
Length at transformation	14 mm SL, but remains pelagic until 35 mm SL
Sequence of fin development	
Pigment	• Dense pigment covering body except for caudal peduncle

Diagnostic characters (see Table 3)

- Develops bony bumps on areas where head spines occur in *Hemilepidotus* spp.
- Becomes increasingly deep-bodied with development
- Uniformity of pigment

^aO'Connell 1953

^bPillsbury 1957

^cFeder et al. 1974


^dBurge and Schultz 1973

^eWang 1981

^fRichardson and Washington 1980


Ref: O'Connell 1953, Richardson and Washington 1980, Wang 1981, Washington et al. 1984b.

A



5.3 mm NL


B



8.6 mm NL

Dense, uniform pigment
except at caudal peduncle


C



10.4 mm SL

Deep body

D



13.8 mm SL

Figures A–D, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 33-33-35 ^a Precaudal: 11-11-11 ^b Caudal: 24-24-24 ^b
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic S: 1-1-1 R: 2-3-3 ^c
Dorsal fin	S: 9-10-10 R: 14-16-17
Pectoral fin	R: 14-15-16
Anal fin	R: 10-12-12
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	N. California, 38-42°N, to SE Alaska, 55-59°N
Ecology	Nearshore shelf demersal, intertidal to 82 m
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	~3 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	<50% SL
Length at flexion	6.2-9.4 mm SL
Length at transformation	13-19 mm SL
Sequence of fin development	Caudal, 2nd dorsal (rays) and anal, 1st dorsal (spines) and pectorals, pelvics (pectorals and pelvics formed by 11.5 mm SL)

Pigment

- Low number (<15) ventral midline melanophores
- Dorsal and anal finfolds
- Base of cleithrum

Diagnostic characters

- Short gut (33% SL)
- Pointed snout
- Pigment (see above)
- Four preopercular spines, other spines (as for *Myoxocephalus* group)
- Pelvic fin ray count I,2; visible in late-stage larvae and juveniles

^aTotal myomere count from Figures A and B = 36.

^bB.B. Washington, NMFS Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560, unpubl.

^cI,2 is the most common count. The first ray is greatly thickened and broadly branched distally, probably leading to the erroneous counts of I,3 in the literature (B.B. Washington, pers. commun., 8 Dec. 1986).

Ref: Richardson and Washington 1980 (as *Icelinus* spp.), Washington 1986, Washington et al. 1984b.

A




Figure A, Washington 1986; B–E, Richardson and Washington 1980. Total myomere counts for Figures B and C are 36.

COTTIDAE**(Myoxocephalus Group)*****Ascelichthys rhodorus* Jordan and Gilbert 1881****MERISTICS**

Vertebrae	Total: 33-36-36 Precaudal: 10-11-12 Caudal: 24-25-25
Branchiostegal rays	6-6-7
Caudal fin	10-13, 6+7, 8-11
Pelvic fin	Absent
Dorsal fin	S: 7-9-10 R: 17-18-19
Pectoral fin	R: 16-17-18
Anal fin	R: 13-15-16
Gill rakers	U: 0-X-3 L: 3-X-5

LIFE HISTORY

Range	Cent. California, 34-38°N, to SE Alaska, 55-59°N
Ecology	Intertidal, nearshore shelf demersal
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Mar (British Columbia) ^a Area: Cobble beach, under boulders ^a Mode: Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.7-2.0 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous, transparent blue developing to purple
Envelope	Smooth
Hatch size	6 mm SL
Incubation time/temp.	24 d/10°C
Pigment	

Diagnostic characters**LARVAE**

Preanal length	<50% SL
Length at flexion	8.8-11.0 mm SL
Length at transformation	12-13 mm SL
Sequence of fin development	
Pigment	<ul style="list-style-type: none"> • Heavy pigment on head and gut • 20-30 ventral midline melanophores prior to flexion, 15-20 in postflexion larvae


Diagnostic characters

- Moderately slender form
- Pointed snout
- Four preopercular spines
- Ventral midline melanophores (the number at various developmental stages)


^aMatarese and Marliave 1982

Ref: Matarese and Marliave 1982.


A




B



C



D



Figures A–D, Matarese and Marliave 1982.

MERISTICS

Vertebrae	Total: 35-35-36
	Precaudal: 10-11-11
	Caudal: 24-25-25
Branchiostegal rays	6-6-6
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 2-3-3
Dorsal fin	S: 8-10-11 R: 14-16-17
Pectoral fin	R: 16-17-18
Anal fin	R: 14-16-17
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Nearshore shelf demersal, intertidal to 142 m
ELH pattern	Oviparous; adhesive, demersal eggs; pelagic larvae
Spawning	Season: Winter-spring (Calif.) ^a Area: Mode: Internal fertilization ^b Migration:
Fecundity	Range/function: 450-1900 (\bar{x} 1043) may produce three clutches/season ^a
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.02-1.05 mm
No. of oil globules	One large, 5-8 small
Oil globule diameter	0.3 mm (large)
Yolk	
Envelope	Salmon-colored
Hatch size	2.9-3.0 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	<50% SL
Length at flexion	
Length at transformation	Begins ~16.6 mm SL
Sequence of fin development	Caudal first, pelvics last
Pigment	

- Dorsolateral gut pigment generally light according to Richardson and Washington (1980),^c but specimens we collect have heavier pigment (compare Figures A and B with C and D)
- Our specimens have heavy crown pigment

Diagnostic characters

- Number of ventral midline melanophores >40 in preflexion larvae (<6 mm SL) and >20 in others
- Upper two preopercular spines larger than lower two


^aGoldberg 1980b

^bMisitano 1980

^cRichardson suspects Figures C and D are incorrect; series requires reevaluation (S.L. Richardson, deceased, pers. commun., Oct. 1984). Washington suspects they are *Icelinus*, probably *I. quadriseriatus*, which occurs outside our study area (B.B. Washington, NMFS Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560, pers. commun., 8 Dec. 1986).


Ref: Misitano 1980, Richardson and Washington 1980, Washington et al. 1984b.

A




> 40 ventral midline melanophores in preflexion larvae

B



C See footnote c.



D See footnote c.




Gap between 3rd and 4th dorsal spines

E



F



Figures A–B, NWAFC originals (B. Vinter); C–F, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 29-31-31
	Precaudal: 11-11-12
	Caudal: 18-20-20
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 7-8-9 R: 9-12-13
Pectoral fin	R: 15-16-17
Anal fin	R: 8-9-10
Gill rakers	U: 0-X-1 L: 4-X-6

LIFE HISTORY

Range	Cent. California, 34-38°N, to Gulf of Alaska, 54-60°N
Ecology	Nearshore shelf demersal, 0-20 m
ELH pattern	Oviparous; adhesive, demersal, guarded eggs; pelagic larvae
Spawning	Season: Feb-May (Puget Sound) ^a Area: On rocks or pilings nearshore ^a Mode: Internal fertilization likely, ^b polygamous male guards nests ^c Migration:
Fecundity	Range/function: 18,800-31,900 (spawn twice each season) ^a
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.7-2.0 mm
No. of oil globules	One
Oil globule diameter	0.36 mm
Yolk	Homogeneous, orange
Envelope	Clear, orange-brown
Hatch size	4.9-5.2 mm SL (5 mm SL)
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	48-55% SL
Length at flexion	5.2-7.0 mm SL
Length at transformation	7.6-7.8 mm SL
Sequence of fin development	Caudal first; pectorals and pelvics formed by 9 mm SL

Pigment

- Dorsal surface of gut
- Ventral midline melanophores (≤15)
- Nape

Diagnostic characters


- Only stout-bodied member of *Myoxocephalus* group
- Ventral midline: ≤15 melanophores beginning several myomeres after anus
- Four preopercular spines
- Head spines prominent (e.g., parietal)
- Low vertebral count (29-31)

Eggs and larvae of *E. diceraus* and *E. lucasi* are unknown. The following information may aid in their identification.

	<i>E. diceraus</i>	<i>E. lucasi</i>
Total vertebrae	31-35	32-34
Dorsal fin spines	7-8	7-8
Dorsal fin rays	13-15	12-14
Anal fin rays	11-13	9-11
Pectoral fin rays	15-19	15-18
Pelvic fin rays	2-3	2-3
Range	SE Alaska-Arctic	Brit. Col.-Bering Sea

^aDeMartini 1978
^bAndriashev 1954
^cMisitano 1978

Ref: Misitano 1978, Richardson and Washington 1980, Washington et al. 1984b.




Figures A–C, Richardson and Washington 1980.

Presently we cannot identify to species *Gymnocanthus* larvae from samples collected in our study area. We identify one type, *Gymnocanthus* A, and include illustrations of *G. hertzensteini* for comparison. Generic characters include:

- Patch of pigment on crown
- Heavy dorsal pigment on gut, increasing dorsolaterally with development
- Numerous, closely spaced, ventral midline melanophores in preflexion larvae

The following meristic information may aid in their identification.

Species	Distribution	Vertebrae	Fins				
			Dorsal		Anal rays	Pectoral rays	Pelvic rays
			Spines	Rays			
<i>Gymnocanthus detriscus</i>	Bering Sea	37-39	IX-XI	15-18	15-19	19-20	I,3
<i>Gymnocanthus galeatus</i>	Brit. Col. - Bering Sea	37-40	X-XII	14-17	17-20	19-21	I,3
<i>Gymnocanthus pistilliger</i>	SE Alaska - Bering Sea	35-38	IX-XI	13-16	14-18	15-20	I,3
<i>Gymnocanthus tricuspis</i>	Bering Sea - Arctic	36-40	X-XII	15-17	15-18	17-20	I,3



Figures A–D, Kyushin 1970 (reared from specimens collected near Hokkaido, Japan; B–D, redrawn); E–F, NWAFC originals (B. Vinter).

MERISTICS

Vertebrae	Total: 36-38-40 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
Dorsal fin	S: 1-1-1 R: 3-3-3
Pectoral fin	S: 10-11-12 R: 15-15-17
Anal fin	R: 17-18-20
Gill rakers	R: 15-17-18 U: X-X-X L: X-X-X

LIFE HISTORY

Range	Bering Sea, 54-66°N, to Arctic (throughout)
Ecology	Epi- and mesobenthic, 0-240 m
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Winter-spring ^a Area: Mode: Internal fertilization likely ^a Migration:
Fecundity	Range/function: 2000-3500 (117-158 mm specimens) ^a
Age at first maturity	4 yr ^a
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	2 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length
Length at flexion
Length at transformation
Sequence of fin development
Pigment ^b

- Heavy pigment on crown and dorsolateral surface of gut
- Larvae >12.2 mm SL have pigment along dorsal midline, and larvae >13.9 mm SL develop pigment along lateral line
- Ventral midline melanophores are widely spaced anteriorly and closely spaced posteriorly


Diagnostic characters

- Specimens of other species of *Gymnocanthus* at comparable developmental stages have not been described

^aAndriashev 1954

^bSpecimens have not been collected from our study area. Illustrations of out-of-area specimens are presented for comparison and may differ.

Ref: Andriashev 1954, Khan 1972, Washington et al. 1984b.



Figures A–E (C, dorsal view; E, ventral view), Khan 1972 (Atlantic specimens, redrawn).


Eggs and larvae of *Icelinus* spp. from the study area are unknown. Illustrations of *Icelinus* A and B, both tentatively identified as *I. borealis* by Washington and Richardson (unpubl.),¹ are presented here.

The following meristic characters may aid in their identification.


Species	Distribution	Vertebrae	Fins				
			Dorsal		Anal rays	Pectoral rays	Pelvic rays
			Spines	Rays			
<i>Icelinus borealis</i>	Wash. - Bering Sea	35-36	IX-XI	14-17	11-14	14-17	I,2
<i>Icelinus burchami</i>	S. Calif. - SE Alaska	35-37	X-XI	16-18	12-14	16-19	I,2
<i>Icelinus filamentosus</i>	S. Calif. - Gulf of Alaska	34-37	X-XII	15-17	13-16	16-18	I,2
<i>Icelinus fimbriatus</i>	S. Calif. - Brit. Col.	37	XI	15-17	13-14	17	I,2
<i>Icelinus tenuis</i> *	SSC - Brit. Col.	37-39	IX-XI	16-18	14-17	15-17	I,2

*A partial series of *I. tenuis* larvae has been identified at Los Angeles County Museum (R. Feeney, Los Ang. Cty. Mus. Nat. Hist., 900 Exposition Blvd., Los Angeles, CA 90007, pers. commun., Oct. 1986).


¹B.B. Washington, NMFS Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560.

ICELINUS*Icelinus A***A**Widely spaced postanal
ventral melanophores


7.1 mm SL

*Icelinus B***B**

7.9 mm SL

C

11.9 mm SL

D

14.4 mm SL

Figures A–D, Washington and Richardson, unpubl.

Eggs and larvae of *Icelus* spp. from our study area are unknown. The following meristic characters may aid in their identification.

Species	Distribution	Vertebrae	Fins				
			Dorsal		Anal rays	Pectoral rays	Pelvic rays
			Spines	Rays			
<i>Icelus canaliculatus</i>	Gulf of Alaska - Bering Sea	41	VII-VIII	22-25	18-20	15-19	I,3
<i>Icelus euryops</i>	Gulf of Alaska - Bering Sea		VIII-X	20-23	15-19	16-18	I,3
<i>Icelus scutiger</i> *	Gulf of Alaska - Bering Sea	39-41	IX-X	18-21	17-19	17-19	I,3
<i>Icelus spatula</i>	Gulf of Alaska - Arctic	39-41	VII-X	17-22	13-18	16-20	I,3
<i>Icelus spiniger</i>	Brit. Col. - Bering Sea	40-42	VIII-X	19-24	15-20	17-20	I,3
<i>Icelus uncinalis</i>	Bering Sea	37-40	IX-X	18-20	14-16	17-18	I,3

*Placed in the genus *Rastrinus* by D.W. Nelson (1984).

Presently we cannot identify to species *Myoxocephalus* larvae from samples we collect in our study area. We collect at least two types based on the following pigmentation characters:





***Myoxocephalus* B** Ventral midline melanophore series (>40 spots), dorsal pigment on gut, and crown pigment; with development, dorsolateral pigment appears covering about 2/3 body length, small ventrolateral patch appears at midbody. Ventrolateral body from over gut to about six myomeres after anus is unpigmented.

***Myoxocephalus* G (probably *M. polyacanthocephalus*)** Preflexion pigment pattern similar to flexion pattern of the other *Myoxocephalus* type; crown, gut, and ventral midline series with dorsolateral and lateral pigment covering 2/3 body length. Body pigment becomes more intense and crown pigment covers entire head with development. No unpigmented area on lateral body over gut and anus.

The following meristic characters may aid in their identification.

Species	Distribution	Vertebrae	Fins				
			Dorsal		Anal rays	Pectoral rays	Pelvic rays
			Spines	Rays			
<i>Myoxocephalus axillaris</i> *	Bering Sea - Arctic		VIII-X	15-17	11-13	14-16	I,3
<i>Myoxocephalus brandti</i>	Bering Sea		IX	15	12-13	16-17	I,3
<i>Myoxocephalus jaok</i>	Gulf of Alaska - Chukchi Sea	35-38	VIII-XI	13-17	12-16	17-19	I,3
<i>Myoxocephalus niger</i>	Bering Sea	36-39	VIII-X	14-18	10-12	16-18	I,3
<i>Myoxocephalus polyacanthocephalus</i>	Wash. - Bering Sea	34-37	IX-X	10-15	8-13	16-19	I,3
<i>Myoxocephalus quadricornis</i>	Bering Sea - Arctic	38-42	VII-X	12-16	13-17	14-18	I,3
<i>Myoxocephalus scorpioides</i> *	Bering Sea - Arctic	35-38	VIII-X	13-18	10-14	14-17	I,3
<i>Myoxocephalus scorpius</i> *	SE Alaska - Arctic	38-46	VIII-XII	12-20	10-16	16-19	I,3
<i>Myoxocephalus stelleri</i>	Bering Sea - Chukchi Sea		VIII-X	15-16	11-14	16-18	I,3
<i>Myoxocephalus verrucosus</i> *	Brit. Col. - Arctic		IX-XII	15-18	13-15	16-19	I,3

*According to Neelov (1979), *M. scorpioides* = *M. axillaris*, and *M. scorpius* in the study area is *M. verrucosus*.

MYOXOCEPHALUS*Myoxocephalus* B**A****B***Myoxocephalus* G**C****D***M. polyacanthocephalus*

Figures A–C, NWAFC originals (B. Vinter); D, Richardson 1981a.

MERISTICS

Vertebrae	Total: 38-X-46
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 8-X-12 R: 12-X-20
Pectoral fin	R: 16-X-19
Anal fin	R: 10-X-16
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Brit. Col., 48°30'-55°N, to Arctic, not specific
Ecology	Epi-, meso-, and bathybenthal, 0-550 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Winter ^b Area: Mode: Eggs guarded by males ^b Migration:
Fecundity	Range/function: 2700 ^b
Age at first maturity	3-4 yr ^b
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	2.0-2.5 mm; ^b 1.8-2.5 mm ^c
No. of oil globules	Several
Oil globule diameter	0.4-0.5 mm
Yolk	
Envelope	
Hatch size	7.4-8.6 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	
Length at flexion	9-15 mm SL
Length at transformation	17-20 mm SL
Sequence of fin development	
Pigment	

- Crown, dorsolateral gut surface, ventral midline starting at myomere 16
- Develops midbody patch at flexion with dorsal pigment extending anteriorly


Diagnostic characters

^a According to Neelov (1979), specimens from our area are *M. verrucosus*. Illustrations of Atlantic specimens of *M. scorpius* are presented for comparison. Specimens from our area may differ.

^b Andriashev 1954

^c Washington et al. 1984b

Ref: Andriashev 1954, Washington et al. 1984b.



Figures A–B, E, Fahay 1983 (A, after Rass 1949; B, E, after Khan 1972); C–D (C, ventral view), Khan 1972 (B–E, Gulf of St. Lawrence specimens).

MERISTICS

Vertebrae	Total: 42-42-43
	Precaudal: 12-12-12 ^a
	Caudal: 31-31-31 ^a
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 12-13-13 R: 19-19-20
Pectoral fin	R: 14-15-15
Anal fin	R: 23-24-24
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	S. California, 32-34°N, to Brit. Col., 48°30'-55°N
Ecology	Nearshore shelf demersal, 0-183 m
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	<5.6 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE





Preanal length	35-36% SL, increasing with development
Length at flexion	
Length at transformation	~25 mm SL
Sequence of fin development	
Pigment	
	• Gut pigmented all over
	• Ventral midline melanophores >30, decreasing to 15-20 by postflexion stage
	• Melanophores near tail tip

Diagnostic characters

- Pigment pattern (tail pigment)
- Pointed snout and slender body
- Four preopercular spines
- Myomeres (42-43)
- Spiny scales develop in postflexion larvae

^aB.B. Washington, NMFS Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C.
20560, unpubl.

Ref: Richardson and Washington 1980.



Figures A–D, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 38-39-40
	Precaudal: 12-X-13
	Caudal: 27-27-27
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 8-9-10 R: 20-22-23
Pectoral fin	R: 17-18-20
Anal fin	R: 21-23-25
Gill rakers	U: X-X-1 L: 7-X-8

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesobenthal, 18-283 m
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous
Envelope	
Hatch size	≤4.7 mm NL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	44-54% SL
Length at flexion	7.2-10.9 mm NL
Length at transformation	≥14-15 mm SL
Sequence of fin development	
Pigment	

- Heavy over body except dorsolaterally above body midline over gut and dorsally and laterally on tail tip


Diagnostic characters (see Table 3)

- Heavy pigmentation
- Preopercular spines not prominent
- Gut distinctively coiled

Distinguished from *Scorpaenichthys marmoratus* (p. 382) by


- Series of melanophores along lateral line

A



4.7 mm NL

B




Mediolateral series

Pointed head


9.6 mm NL

C



12.6 mm SL

D



14.4 mm SL

Figures A–D, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 39-40-40
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 8-9-11 R: 20-20-22
Pectoral fin	R: 18-19-20
Anal fin	R: 21-22-23
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	S. California, 32-34°N, to Brit. Col., 48°30'-55°N
Ecology	Nearshore shelf demersal, 15-146 m
ELH pattern	Oviparous, eggs probably demersal, pelagic larvae
Spawning	Season: Area: Mode: Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous
Envelope	
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	~58% SL
Length at flexion	~8.7 mm SL
Length at transformation	
Sequence of fin development	
Pigment	
• Generally heavily pigmented over anterior 3/4 body	

Diagnostic characters (see Table 3)

- Distinguished from *R. asprellus* by
- Greater portion of tail unpigmented
 - Morphological differences
 - Preanal length longer
 - Greater body depth




Figure A, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 38-39-39 Precaudal: 12-12-12 Caudal: 26-26-26
Branchiostegal rays	6-6-6 ^a
Caudal fin	8-9, 6+5, 7-8
Pelvic fin	Thoracic S: 1-1-1 R: 3-3-3
Dorsal fin	S: 8-10-10 R: 19-20-21
Pectoral fin	R: 21-22-24
Anal fin	R: 18-20-21
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	S. California, 32-34°N, to SE Alaska, 55-59°N
Ecology	Intertidal, nearshore shelf demersal
ELH pattern	Oviparous; demersal, attached, adhesive eggs; pelagic larvae
Spawning	Season: Jan-Feb through Apr (British Columbia), larvae collected in spring; ^b May ^c Area: Shallow, rocky subtidal areas (extreme nearshore), larvae collected 0 and 4 m from shore ^b Mode: Internal fertilization, eggs laid in masses on <i>Laminaria</i> holdfasts (15-20 eggs/mass) ^b Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous, pink (wild)
Envelope	
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE^d

Preanal length	37-42% SL
Length at flexion	6.5-8.5 mm SL
Length at transformation	
Sequence of fin development	Pectorals and caudal, dorsal and anal, pelvics
Pigment	<ul style="list-style-type: none"> • Gut pattern • High number of postanal ventral melanophores (\bar{x} 51 in preflexion, 47 in flexion)

Diagnostic characters





- Pigment on gut
- Accessory or inner-shelf spines associated with preopercular spines in preflexion larvae (not present in other members of *Myoxocephalus* group)
- Pectoral fin rays develop toward each other ventrally
- Spines: Nasal spines develop at 12-14 mm SL

^a According to Marliave et al. (1985), branchiostegal rays = 7.

^b Marliave et al. 1985

^c Hart 1973

^d Tentative placement in *Myoxocephalus* group.



Figures A–B, Marliave et al. 1985; C–D, NWAFC originals (B. Vinter).

Presently we cannot identify to species *Triglops* spp. larvae from samples we collect in our study area. Illustrations of *Triglops* sp. (designated as B here) (Richardson and Washington 1980) and *Triglops* A (Washington and Richardson, unpubl.¹) are presented here. Generic characters include:


- High myomere count (>45)
- Pointed snout
- Heavy dorsolateral pigmentation on gut
- Postanal ventral midline melanophores, probably becoming embedded in postflexion larvae (except *Triglops* B as figured by Richardson and Washington [1980]).

The following meristic information may aid in their identification.


Species	Distribution	Vertebrae	Fins				
			Dorsal		Anal rays	Pectoral rays	Pelvic rays
			Spines	Rays			
<i>Triglops forficata</i>	Gulf of Alaska - Bering Sea	52-54	IX-XI	27-32	27-32	20-22	I,3
<i>Triglops jordani</i>	Bering Sea		IX-XI	24-28	23-29	19-21	I,3
<i>Triglops macellus</i>	Wash. - Bering Sea	51	X-XI	27-31	27-31	15-17	I,3
<i>Triglops metopias</i>	SE Alaska - Bering Sea	48-49	X-XI	23-27	22-27	18-22	I,3
<i>Triglops pingeli</i>	Wash. - Arctic	46-48	IX-XIII	22-28	20-28	16-19	I,3
<i>Triglops scepticus</i>	SE Alaska - Bering Sea	45-46	X-XII	21-23	22-24	17-19	I,3

¹B.B. Washington, NMFS Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560.


TRIGLOPS

Triglops A (possibly *T. scepticus* based on counts)**A**


Ventral midline melanophores 6.5 mm SL

B

16.9 mm SL

Triglops B**C**

6.9 mm NL

Lack of postanal
body pigment**D**

15.4 mm SL

Figures A–B, Washington, unpubl.; C–D, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 32-33-35
	Precaudal: 11-X-12 ^a
	Caudal: 22 ^a -X-23
Branchiostegal rays	6-6-6
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 8-9-10 R: 16-17-18
Pectoral fin	R: 14-15-16
Anal fin	R: 12-13-14
Gill rakers	U: 1-1-1 L: 4-X-5

LIFE HISTORY

Range	Cent. California, 34-38°N, to Aleutian Is., 51-55°N
Ecology	Nearshore shelf demersal, intertidal to 55 m
ELH pattern	Oviparous; demersal, adhesive, attached eggs; pelagic larvae
Spawning	Season: Feb-May ^b Area: Intertidal, under rocks ^b Mode: Polygamous males, eggs laid in nests ^b Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

^aB.B. Washington, NMFS Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560, unpubl.

^bMarliave 1975a

^cPigment patterns within the genus *Artedius* appear to be highly variable. Reared specimens vary between hatches and stages (J. Marliave, Vancouver Public Aquarium, P.O. Box 3232, Vancouver, B.C., Canada V6B 3X8, pers. commun., 16 Oct. 1986). Wild specimens collected from Yaquina Bay, Oregon, show variations from the pigment patterns described in Washington (1986) (B. Mundy, NMFS Southwest Fish. Cent., Honolulu Lab., 2570 Dole St., Honolulu, HI 96822, pers. commun., 1 Oct. 1986).

Ref: Marliave 1975a; Richardson and Washington 1980 (see *Artedius* 2); Washington 1981, 1986.

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous; blue, purple, or gray
Envelope	
Hatch size	3.5-3.8 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	<50% SL
Length at flexion	5.9-6.8 mm NL
Length at transformation	12-14 mm SL
Sequence of fin development	Caudal, dorsal and anal rays, dorsal spines, pectorals, pelvics

Pigment^c

- Lack of head pigment
- Usually relatively light nape pigment present
- 13-19 postanal ventral midline melanophores

Diagnostic characters

- Distinguished from other cottids by
- Body shape stubby, humped
 - Presence of gut diverticula
 - Preopercular spines (postflexion, 18-22)
 - Parietal spines (postflexion)


Distinguished from other *Artedius* spp. with gut diverticula by

	Preflexion		Flexion	
	No. of postanal ventral melanophores	Presence of nape pigment	Presence of brain pigment	No. of preopercular spines
<i>A. fenestralis</i>	13-19	Irreg.	No	18-22
<i>A. lateralis</i>	22-32	No	Yes	14-16
<i>Artedius</i> 3	9-13	Yes	No	22-24

A




B




Heavy dorsolateral pigment on gut surface
Occurrence of nape pigment is irregular

C



Parietal spines

D



E




Figure A, Washington 1986; B–E, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 32-33-34 Precaudal: 11-X-12 Caudal: 22-X-23
Branchiostegal rays	7-7-7
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic S: 1-1-1 R: 3-3-3
Dorsal fin	S: 9-9-9 R: 15-17-18
Pectoral fin	R: 13-14-15
Anal fin	R: 10-13-14
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	S. California, 32-34°N, to Aleutian Is., 51-55°N
Ecology	Nearshore shelf demersal, intertidal to 21 m
ELH pattern	Oviparous, eggs unknown, pelagic larvae
Spawning	Season: Possibly spring (California) ^a Area: Mode: Internal fertilization likely ^b Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous
Envelope	
Hatch size	<3 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	<50% SL
Length at flexion	5.2-6.4 mm NL
Length at transformation	~13.6 mm SL (12-14 mm SL)
Sequence of fin development	Caudal, dorsal and anal rays, dorsal spines, pectorals, pelvics
Pigment (see <i>A. fenestralis</i> , p. 414, footnote c)	<ul style="list-style-type: none"> • Lack of head pigment • Presence of nape pigment • 21-33 postanal ventral midline melanophores

Diagnostic characters

- Distinguished from other *Artedius* spp. (*A. fenestralis*, *A. lateralis*, and *Artedius* 3) by
- Pigment (see above)
 - Absence of gut diverticula
 - Humped appearance in nape region
 - Preopercular spines: 18-22 (late flexion, postflexion)
 - Seven branchiostegal rays


^aBurge and Schultz 1973

^bNWAFRC unpubl.


Ref: Richardson and Washington 1980; Washington 1981, 1986.

A


B



C



D



E




Figure A, Washington 1986; B-E, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 32-33-34
	Precaudal: 11-11-11
	Caudal: 21-22-23
Branchiostegal rays	6-6-6
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 7-9-10 R: 15-16-17
Pectoral fin	R: 14-15-16
Anal fin	R: 12-13-14
Gill rakers	U: 1-X-2 L: 6-X-9

LIFE HISTORY

Range	South of southern California to Aleutian Is., 51-55°N
Ecology	Nearshore shelf demersal, intertidal to 13 m
ELH pattern	Oviparous; demersal, adhesive, attached eggs; pelagic larvae
Spawning	Season: Winter-spring (Brit. Col.); ^a June (Puget Sound) ^b
	Area: Demersal, on underside of rocks ^a
	Mode:
	Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	0.98-1.16 mm (1.07 mm)
No. of oil globules	One
Oil globule diameter	0.22 mm
Yolk	Homogeneous; red, yellow, or orange
Envelope	Colorless, 0.031 mm thick
Hatch size	3.9-4.5 mm SL (4.1 mm SL)
Incubation time/temp.	16 d/15°C
Pigment	

Diagnostic characters

LARVAE

Preanal length	<50% SL
Length at flexion	5.0-6.3 mm SL
Length at transformation	8-10 mm SL
Sequence of fin development	Caudal, dorsal and anal rays, dorsal spines, pectorals, pelvics
Pigment (see <i>A. fenestralis</i> , p. 414, footnote c)	
	<ul style="list-style-type: none"> • Lack of head and nape pigment is preflexion larvae • Number (22-32) and size variation of postanal ventral midline melanophores • Postflexion larvae have a marked increase in head pigment

Diagnostic characters






- Presence of gut diverticula
 - Pigment (see above)
- Distinguished from other *Artedius* spp. with gut diverticula (*A. fenestralis* and *Artedius* 3) by
- See *A. fenestralis*

^aMarliave 1977
^bNWAFC, unpubl.

Ref: Marliave 1975a; Richardson and Washington 1980; Washington 1981, 1986; Washington et al. 1984b.

A

B Lack of head and nape pigment



Figures A–E, Washington 1986.

MERISTICS ***A. corallinus* (Hubbs 1926)**

Vertebrae	Total: 31-32-33
	Precaudal: X-X-X
	Caudal: 41-X-43
Branchiostegal rays	6-6-6
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 8-9-9 R: 15-16-16
Pectoral fin	R: 14-15-16
Anal fin	R: 12-13-13
Gill rakers	U: X-X-X L: X-X-X

MERISTICS ***A. notospilotus* Girard 1856**

Vertebrae	Total: 32-33-34
	Precaudal: 12-12-12
	Caudal: 21-21-21
Branchiostegal rays	6-6-6
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 9-9-10 R: 14-15-16
Pectoral fin	R: 14-16-17
Anal fin	R: 11-12-13
Gill rakers	U: 2-2-2 L: 8-X-10

LIFE HISTORY

Range	South of southern California to Washington, 46-48°30'N
Ecology	Intertidal, nearshore shelf demersal
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	≤2.9 mm NL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE (*Artedius* 3)^a

Preanal length	45% SL
Length at flexion	5.6-6.9 mm NL
Length at transformation	
Sequence of fin development	Caudal, dorsal and anal rays, dorsal spines and pectorals, pelvics

- Pigment** (see *A. fenestralis*, p. 414, footnote c)
- Nape pigment
 - 9-13 evenly spaced postanal ventral midline melanophores; begin about 3-4 myomeres posterior to anus

Diagnostic characters

Distinguished from other *Artedius* spp. with gut diverticula (*A. fenestralis* and *A. lateralis*) by


- Low number of postanal ventral midline melanophores in small larvae (9-13)
- Pigment: Lack of head pigment and presence of nape pigment
- Preopercular spines form earlier than other *Artedius* spp. (<4.1 mm NL)
- Number of preopercular spines higher than in other *Artedius* spp.

^a *Artedius* 3 larvae are either *A. corallinus* or *A. notospilotus*. For a complete description see Washington (1986).

Artedius 3

A

B


Gut diverticula and
nape pigment

3.2 mm NL




Artedius type spine
pattern: dorsalmost,
middle, and ventralmost
spines larger than rest

C

9–13 evenly spaced postanal
ventral melanophores

6.8 mm SL

D



7.3 mm SL

Figures A–D, Washington 1986.

COTTIDAE
(*Artedius* Group)

***Clinocottus acuticeps* (Gilbert 1895)**

MERISTICS

Vertebrae	Total: 31-32-33
	Precaudal: 10-X-11 ^a
	Caudal: 21-22-23
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 7-8-9 R: 13-15-17
Pectoral fin	R: 13-14-15
Anal fin	R: 9-12-13
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Cent. California, 34-38°N, to Bering Sea, 54-66°N
Ecology	Intertidal, nearshore shelf demersal
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Spring ^b Area: Vertical rock surfaces under <i>Fucus</i> in upper intertidal zone (laid in monolayer) ^b Mode: Internal fertilization likely ^c Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.0-1.2 mm
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous; brown or purple
Envelope	
Hatch size	3-4 mm NL (3.1-3.3 mm NL)
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	~63% SL
Length at flexion	5.5-7.3 mm NL
Length at transformation	13-14 mm SL
Sequence of fin development	Caudal, dorsal and anal rays, dorsal spines, pectorals, pelvics
Pigment	<ul style="list-style-type: none">Dorsal gut surfaceRelatively few (4-10) ventral midline melanophores

Diagnostic characters

- Distinguished from all other cottids by
- Long gut with hindgut diverticula
 - Flabby appearance
 - Outer bubble of skin

^aB.B. Washington, NMFS Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560, unpubl.

^bMarliave 1981a

^cAndriashev 1954

Ref: Blackburn 1973; Richardson and Washington 1980; Washington 1981, 1986; Washington et al. 1984b.

A




Figure A, Washington 1986; B–E, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 33-33-34
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 8-9-10 R: 14-15-17
Pectoral fin	R: 12-14-15
Anal fin	R: 9-10-12
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Intertidal
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Area: Mode: Internal fertilization likely ^a Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous
Envelope	
Hatch size	~4 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	<50% SL
Length at flexion	6.4-9.6 mm SL
Length at transformation	13-14 mm SL
Sequence of fin development	Caudal, 2nd dorsal (rays) and anal, pectorals, 1st dorsal (spines), pelvics

Pigment

- Presence of head pigment is variable during development
- Light pigment on gut
- High number ventral midline melanophores (15-21)


Diagnostic characters

- Usually lacks head pigment; although variable, other *Clinocottus* spp. have head and snout pigment
- Long trailing gut
- Absence of hindgut diverticula
- 11-14 preopercular spines, tiny parietal spine

^a Andriashev 1954

Ref: Richardson and Washington 1980 (see Cottidae 2); Washington 1981, 1986.

A



Figures A, D–E, Washington 1986; B–C, Richardson and Washington 1980.

COTTIDAE
(Artedius Group)

***Clinocottus globiceps* (Girard 1858)**

MERISTICS

Vertebrae	Total: 32-33-34
	Precaudal: 11-12-12
	Caudal: 20-21-23
Branchiostegal rays	6-6-6 (occasionally 7)
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 8-9-10 R: 13-16-17
Pectoral fin	R: 13-14-15
Anal fin	R: 11-11-12
Gill rakers	U: 1-1-1 L: 5-5-5

LIFE HISTORY

Range	S. California, 32-34°N, to Gulf of Alaska, 54-60°N
Ecology	Intertidal
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Area: Mode: Internal fertilization likely ^a Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.5-2.0 mm
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous
Envelope	
Hatch size	5.1-5.4 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	<50% SL
Length at flexion	6.2-~8.1 mm SL
Length at transformation	12.9-13.5 mm SL
Sequence of fin development	Caudal; dorsal, anal, and pectorals; pelvics
Pigment	<ul style="list-style-type: none">• Heavy on head, nape, dorsolateral surface of gut• Ventral midline melanophores along posteriormost 10 myomeres


Diagnostic characters

- Late flexion: 15-20 preopercular spines and a cluster of spines in parietal region
- Distinguished from other preflexion *Clinocottus* spp. by
- Heavy pigment on head, nape, and gut
- Distinguished from *C. recalvus* by
- 4-8 ventral midline melanophores


^a Andriashev 1954

Ref: Richardson and Washington 1980 (see Cottidae 3); Washington 1981, 1986; Washington et al. 1984b.


A




4-8 discrete spots beginning ~10 myomeres from tail




6.3 mm SL



7.5 mm NL



12.5 mm SL



12.9 mm SL



Dorsalmost spines larger than rest

Figures A-B, F, Washington 1986; C-E, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 32-33-33
	Precaudal: 10-11-12
	Caudal: 20-22-23
Branchiostegal rays	6-6-6
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 8-9-9 R: 14-16-16
Pectoral fin	R: 13-14-15
Anal fin	R: 9-12-13
Gill rakers	U: 1-1-1 L: 4-X-7

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Intertidal, nearshore shelf demersal
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Area: Mode: Internal fertilization likely, ^a males guard nests ^b
Fecundity	Migration:
Age at first maturity	Range/function:
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.25-1.35 mm (1.3 mm)
No. of oil globules	Several
Oil globule diameter	
Yolk	Homogeneous
Envelope	
Hatch size	~4.6 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	
Length at flexion	
Length at transformation	9-11 mm SL
Sequence of fin development	
Pigment	
	<ul style="list-style-type: none">• Heavy on dorsolateral surface of gut• Series of ventral midline melanophores extending into finfold• Heavy pigment on snout, crown, and nape


Diagnostic characters

- Multiple preopercular spines
- Distinguished from other *Clinocottus* spp. by
- Minimal trailing of hindgut

^aAndriashev 1954

^bMorris 1951

Ref: Morris 1951.



Figures A–G (C, dorsal view), Morris 1951 (B–G, redrawn). Redrawn figures are based only on distribution of melanophores. Other pigment cells shown on the original figures are deleted.

MERISTICS

Vertebrae	Total: 33-34-34
	Precaudal: 11-12-12
	Caudal: 21-22-23
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 8-8-9 R: 15-17-18
Pectoral fin	R: 12-14-15
Anal fin	R: 12-13-14
Gill rakers	U: 1-1-1 L: 4-X-5

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Intertidal
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Apr-July (Puget Sound) ^a Area: Intertidal, between rocks, barnacles, or bay mussels (not attached to substrate) ^b Mode: Internal fertilization, multiple spawning (three) ^a Migration:
Fecundity	Range/function:
Age at first maturity	1 yr ^a
Longevity	

^a Atkinson 1939

^b J. Marliave, Vancouver Public Aquarium, P.O. Box 3232, Vancouver, B.C., Canada V6B 3X8, pers. commun., 25 Oct. 1988.

^c Specimens reared in the laboratory have shown a high degree of variability in pigmentation. Postanal ventral melanophores range from 10 to >50. Other characters appear to vary between geographical area, e.g., presence of ventral or lateral gut pigment and the nape bubble (J. Marliave, pers. commun., 25 Oct. 1988).

Ref: Stein 1973; Washington 1981, 1986; Washington et al. 1984b.

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.3-1.5 mm
No. of oil globules	One large, many small
Oil globule diameter	
Yolk	Green, brown, or red ^b
Envelope	
Hatch size	4.2-4.5 mm NL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	37% SL, increasing with development to 48% SL
Length at flexion	7.2-7.6 mm SL
Length at transformation	7.5-10.0 mm SL
Sequence of fin development	Caudal, dorsal and anal rays, dorsal spines, pectorals, pelvics

Pigment^c

- Head and nape
- Dorsolateral surface of gut
- Postanal ventral midline series, with development becoming more closely spaced posteriorly

Diagnostic characters

- Bubble of skin anterior to origin of dorsal finfold
- Distinguished from *O. snyderi* by
- Ventral midline melanophores (>15)
- Bubble of skin pigmented



Eggs and larvae of *O. rimensis* are unknown. The following information will aid in identification.

Total vertebrae	34-37
Precaudal	11-13
Caudal	22-25
Dorsal fin spines	8-10
Dorsal fin rays	16-19
Anal fin rays	13-15
Pectoral fin rays	13-15
Pelvic fin rays	3
Range	SSC-SE Alaska


A

B


Bump similar to *Artemius* but does not develop into a distinct diverticula



C



D



Figures A–D, Washington 1986 (B–D, reared).

MERISTICS

Vertebrae	Total: 34-35-37 Precaudal: 10-11-12 Caudal: 23-24-25
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic S: 1-1-1 R: 3-3-3
Dorsal fin	S: 7-8-9 R: 17-19-20
Pectoral fin	R: 12-14-15
Anal fin	R: 12-14-15
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59°N
Ecology	Intertidal, nearshore shelf demersal
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Winter-spring; fall in more northerly populations ^a Area: Mode: Internal fertilization (spawn twice/season) ^a Migration:
Fecundity	Range/function:
Age at first maturity	<1 yr ^a
Longevity	>2 yr ^a

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.2-1.3 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	4.47 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	<45% SL
Length at flexion	6.2-8.4 mm SL
Length at transformation	11-13 mm SL
Sequence of fin development	Caudal, dorsal and anal rays, dorsal spines, pectorals, pelvics

Pigment

- Head and nape very lightly pigmented
- Dorsolateral surface of gut
- Ventral midline series with <10 evenly spaced melanophores


Diagnostic characters

- Spines
 - Patch of parietal spines
 - 10-12 spines develop along preopercular margin
 - 8-10 accessory spines form anteriorly at the bases of the preopercular spines
- Distinguished from *O. maculosus* by
 - Bubble of skin anterior to origin of dorsal finfold unpigmented and less obvious than *O. maculosus*
 - Ventral midline melanophores (<10)

^aGrossman and DeVlaming 1984

Ref: Richardson and Washington 1980 (see Cottidae 1); Stein 1973; Washington 1981, 1986; Washington et al. 1984b.

A



Figures A–B, D–E, Washington 1986; C, Richardson and Washington 1980 (B–C, reared).

MERISTICS

Vertebrae	Total: 33-34-35
	Precaudal: 13-13-13 ^a
	Caudal: 20-20-20 ^a
Branchiostegal rays	7-7-7
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 7-X-8 R: 18-X-19
Pectoral fin	R: 14-16-17
Anal fin	R: 12-14-15
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Washington, 46-48°30'N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthic, 0-225 m
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Aug (British Columbia) ^b Area: Rocky subtidal areas on solid substrate ^b Mode: Polygamous males guard nest ^c Migration:
Fecundity	Range/function: \bar{x} 130 ^b
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	~2.3 mm
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous; pink
Envelope	White
Hatch size	~6-7 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	
Length at flexion	
Length at transformation	18-20 mm SL (settle) ^d
Sequence of fin development	
Pigment (<i>Psychrolutes</i> Group)	
	• Head, nape, gut, and pectoral fins
	• Lacks postanal ventral melanophores

Diagnostic characters

- Morphology
 - Tadpole shape
 - Large head
 - Outer layer of loose flabby skin
 - Large pigmented pectoral fins
 - No head or preopercular spines
- Distinguished from *Psychrolutes paradoxus* (p. 436) by
- Pectoral fin ray count (14-17)
 - Less body pigment

^aB.B. Washington, NMFS Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560, unpubl.


^bMarliave 1975a

^cJ. Marliave, Vancouver Public Aquarium, P.O. Box 3232, Vancouver, B.C., Canada V6B 3X8, pers. commun., 16 Oct. 1986.

^dJuveniles have a tendency to reenter the water column for feeding, producing a protracted period of ambivalence about settlement (Marliave 1981b).

Ref: Marliave 1975a, Washington et al. 1984b.

A



B




Figure A, Richardson and Bond 1978; B, Richardson 1981a.

COTTIDAE
(Psychrolutes Group)

***Psychrolutes paradoxus* Günther 1861**

MERISTICS

Vertebrae	Total: 36-36-37
	Precaudal: 11-11-11 ^a
	Caudal: 21-21-21 ^a
Branchiostegal rays	7-7-7
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 9-11-12 R: 12-14-17
Pectoral fin	R: 19-21-23
Anal fin	R: 10-12-14
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Washington, 46-48°30'N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthal, 9-219 m
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Winter-spring (British Columbia) ^b
	Area:
	Mode:
	Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	~1.4 mm
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous
Envelope	
Hatch size	~6-7 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	
Length at flexion	
Length at transformation	~13-14 mm SL
Sequence of fin development	
Pigment	

- Initially restricted to head, gut, and upper body (including pectoral fins); with development, increasing to 3/4 body

Diagnostic characters

- Morphology
 - Tadpole shape
 - Large head
 - Outer layer of loose flabby skin
 - Large pigmented pectoral fins
 - No head or preopercular spines
- Distinguished from *Gilbertidia sigalutes* (p. 434) by
- More lateral pigment on head and body with development
 - Pectoral fin ray count (19-23)

Eggs and larvae of *P. phrictus* are unknown (see Cottoid A, p. 438). The following information will aid in identification.

Total vertebrae	33-35
Dorsal fin spines	8-9
Dorsal fin rays	19-20
Anal fin rays	12-14
Pectoral fin rays	22-26
Pelvic fin rays	3
Range	Cent. Calif.-Bering Sea

^aB.B. Washington, NMFS Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560, unpubl.
^bMarliave 1975a

Ref: Blackburn 1973, Marliave 1975a.





Figure A, Richardson and Bond 1978; B, Marliave 1975a (redrawing provided by Marliave); C, Richardson 1981a; D, NWAFC original (B. Vinter).



Figures A–B, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 34-35-35
	Precaudal: 10-X-13 ^a
	Caudal: 23 ^b -X-26
Branchiostegal rays	7-7-7
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 8-10-11 R: 13-15-16
Pectoral fin	R: 22-25-26
Anal fin	R: 12-14-16
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Washington, 46-48°30'N, to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathybenthal, 18-825 m
ELH pattern	Parity unknown, eggs probably demersal, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous
Envelope	
Hatch size	~7.4 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	<50% SL
Length at flexion	
Length at transformation	
Sequence of fin development	Fins complete at 12 mm SL
Pigment	
	<ul style="list-style-type: none">• Heavy spots on head and gut• Double row of small pigment spots occurs along the midline of the ventral surface of gut; more pronounced in specimens >8 mm SL• Pectoral fin pigmented at base

Diagnostic characters

- Similar to *Psychrolutes* group but with four preopercular spines

^aOne specimen with precaudal vertebrae = 13 (B.B. Washington, NMFS Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560, unpubl.).

^bB.B. Washington, unpubl.

Ref: Blackburn 1973.




Figure A, Richardson 1981a; B, Washington et al. 1984a.

MERISTICS

Vertebrae	Total: 30-32-33
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 8-9-9 R: 12-14-15
Pectoral fin	R: 19-20-23
Anal fin	R: 9-11-12
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Washington, 46-48°30'N, to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathybenthal, 75-1980 m
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE


Preanal length
Length at flexion
Length at transformation
Sequence of fin development
Pigment
• Head, nape, entire gut
• Laterally over 1/4 body, with development increas- ing to 3/4 body

Diagnostic characters

- Fifth accessory preopercular spine (sometimes difficult to see)
- Outer layer of loose skin more pronounced than in other genera (genus)

Eggs and larvae of *M. kincaidi* are unknown. The following information may aid in identification.

Total vertebrae	31-33
Dorsal fin spines	8-10
Dorsal fin rays	13-15
Anal fin rays	10-13
Pectoral fin rays	19-21



Figures A–D, Richardson and Bond 1978.

MERISTICS

Vertebrae	Total: 34-37-39 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic S: 1-1-1 R: 4-4-4
Dorsal fin	S: 7-9-11 R: 18-20-21
Pectoral fin	R: 14-15-17
Anal fin	R: 14-17-18
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	S. California, 32-34°N, to Gulf of Alaska, 54-60°N
Ecology	FW or anadromous type
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Jan-Apr; ^a Feb-July ^b Area: Under rocks ^b Mode: Polygamous males, eggs guarded ^b Migration: Downstream in spring to spawn ^b
Fecundity	Range/function: 280-10,980 ^a
Age at first maturity	
Longevity	>7 yr ^b

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.4-1.6 mm
No. of oil globules	One large, some small
Oil globule diameter	0.2-0.3 mm
Yolk	Homogeneous; granular ^a
Envelope	Thick, transparent, orange
Hatch size	4.5-5.0 mm SL; ^a 5.5-6.3 mm SL ^c
Incubation time/temp.	15-16 d/12°C
Pigment	

Diagnostic characters

LARVAE

Prenal length	40% SL
Length at flexion	~7 mm SL
Length at transformation	10 mm SL
Sequence of fin development	Pectorals and pelvics by 10 mm SL

Pigment

- Dorsal and ventral gut surface
- Postanal ventral midline series decreasing in number and becoming more evenly spaced with development

Diagnostic characters

- Morphology: Slender, round snout
- Spines: Four preopercular, without other head spines
- Gut shape unique, posteriorly forked

Larvae of *C. aleuticus* are unknown. Spawning takes place primarily in freshwater and larvae may not occur in coastal marine plankton.


^aWang 1981

^bMorrow 1980


^cRichardson and Washington 1980

Ref: Morrow 1980, Stein 1972, Richardson and Washington 1980, Wang 1981, Washington et al. 1984b.


A



B



C



Figures A–C, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 35-36-39 Precaudal: 10-11-12 Caudal: 24-25-27
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic S: 1-1-1 R: 4-4-4
Dorsal fin	S: 6-7-8 R: 15-18-20
Pectoral fin	R: 17-19-20
Anal fin	R: 15-16-20
Gill rakers	U: 1-X-3 L: 8-X-10

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal, intertidal to 91 m
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Oct-Mar (California) ^a Area: Mode: Migration:
Fecundity	Range/function: 2000-11,000/ ^b $N=0.355 \times L^{1.84}$, N=no. maturing eggs, L=TL mm ^a
Age at first maturity	1 yr ^a
Longevity	5 yr ^c

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.4-1.5 mm (1.43)
No. of oil globules	One, smaller ones
Oil globule diameter	0.3 mm
Yolk	
Envelope	Thick, transparent, bumpy
Hatch size	3.8-5.0 mm SL; 3.9-4.8 TL ^a
Incubation time/temp.	9-14 d/15°C
Pigment	

Diagnostic characters

LARVAE

Preanal length	37-40% SL
Length at flexion	~8 mm SL
Length at transformation	15-20 mm SL
Sequence of fin development	
Pigment	<ul style="list-style-type: none"> • Internal snout pigment • 6-8 bars of pigment on dorsolateral surface of gut • Postanal ventral midline series

Diagnostic characters

- Gut pigment appearing as 6-8 bars
- Internal snout pigment


^aJones 1962

^bWang 1981

^cFitch and Lavenberg 1975


Ref: Jones 1962, Wang 1981, Washington et al. 1984b.

A




B

Snout pigment internal



C



Figures A–C, Richardson and Washington 1980.

MERISTICS

Vertebrae	Total: 37-38-38
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 7-8-9 R: 20-21-22
Pectoral fin	R: 15-16-17
Anal fin	R: 18-19-20
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Brit. Col., 48°30'-55°N, to Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal
ELH pattern	Probably oviparous, eggs probably demersal, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE




Preanal length
Length at flexion ≤12 mm SL
Length at transformation
Sequence of fin
development
Pigment

- Entire body heavily pigmented with small, densely packed melanophores except for caudal peduncle, opercle area, and ventrolateral gut surface

Diagnostic characters (see Table 3)

- Genus
 - Elongate body becoming deeper with development
 - Pigment relatively heavy
 - Strong frontoparietal ridge
- Distinguished from *B. cirrhosus* by
- More pectoral fin rays (15-17)
 - Pigment
 - Larger area of caudal peduncle unpigmented
 - Smaller, denser melanophores
 - Lack of pigment on underside of mouth

Small, densely packed melanophores



Figures A-C, NWAFC originals (B. Vinter).

MERISTICS

Vertebrae	Total: 37-37-39 ^a Precaudal: 13-13-13 ^a Caudal: 26-26-26 ^a
Branchiostegal rays	6-6-6
Caudal fin	X, 6+6, X
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 6-8-8 R: 20-23-24
Pectoral fin	R: 11-12-13
Anal fin	R: 18-19-21
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Cent. California, 34-38°N, to Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal, intertidal to 37 m
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Late winter (British Columbia) ^b Area: Demersal, on rocks ^c Mode: Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous
Envelope	Clear, light brown
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	<50% SL
Length at flexion	≤11 mm SL
Length at transformation	
Sequence of fin development	
Pigment	<ul style="list-style-type: none">• Entire body heavily pigmented except for caudal peduncle, opercle area, and ventrolateral gut surface• Pigment along underside of mouth between dentary bones (chin)

Diagnostic characters (see Table 3)

- Distinguished from *B. bilobus* by
- Low number of pectoral fin rays (11-13)
 - Shorter area of caudal peduncle unpigmented
 - Pigment on underside of mouth
 - See *B. bilobus* for generic characters (p. 448)


^aB.B. Washington, NMFS Systematics Lab., Natl. Mus. Nat. Hist., Wash., D.C. 20560, unpubl.

^bMarliave 1975a


^cClemens and Wilby 1961

Ref: Marliave 1975a, Washington et al. 1984b.

A



B



Figures A–B, NWAFC originals (B. Vinter).

COTTIDAE
(Hemitripteris Group)

***Hemitripteris villosus* (Pallas [1814])**

MERISTICS

Vertebrae	Total: 39-39-41 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic S: 1-1-1 R: 3-3-3
Dorsal fin	S: 16-17-19 R: 11-12-13
Pectoral fin	R: 18-19-20
Anal fin	R: 12-14-15
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Gulf of Alaska, 54-60°N, to Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Fall (Funka Bay, Japan) ^a Area: Rocky sea bottom, 10-30 m ^a Mode: Migration:
Fecundity	Range/function: 2250-11,170/ $E=0.00002147 \times L^{3.374}$, E=ovarian eggs, L=BL mm ^a
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	4.41-4.61 mm
No. of oil globules	Multiple (120-160)
Oil globule diameter	0.05-0.26 mm
Yolk	Yellow/orange
Envelope	Thick, white
Hatch size	14.1-15.2 mm SL; ^a 10.9-11.6 mm SL ^b
Incubation time/temp.	100 d/12°C
Pigment	• Light on yolksac

Diagnostic characters

- Oil globules coalesce to one by late embryonic development

LARVAE

Preanal length	~50% SL
Length at flexion	≤14.4 mm SL (close to hatching)
Length at transformation	~20 mm SL
Sequence of fin development	
Pigment	• Heavily pigmented at hatching (head, 3/4 BL, and dorsolateral surface of gut) • Distinctive finfold pigment

Diagnostic characters (see Table 3)

- *Hemitripteris* group characters
- Finfold pigment
- Prickles, scales
- Newly hatched larvae large and well developed

Eggs and larvae of *H. bolini* are unknown. The following information will aid in identification.


Total vertebrae	38-40
Dorsal fin spines	11-15
Dorsal fin rays	11-14
Anal fin rays	12-14
Pectoral fin rays	20-22
Pelvic fin rays	3
Range	N. Calif. ^c - Bering Sea

^a Kyushin 1968

^b Okiyama and Sando 1976

^c Lea and Quirollo 1986

Ref: Kyushin 1968, Okiyama and Sando 1976.



Figures A–D, Okiyama and Sando 1976 (reared from specimens collected near Hokkaido, Japan).

COTTIDAE
(Hemitripteris Group)

***Nautichthys oculo fasciatus* (Girard 1858)**

MERISTICS

Vertebrae	Total: 40-41-41	
	Precaudal: X-X-X	
	Caudal: X-X-X	
Branchiostegal rays	6-6-6	
Caudal fin		
Pelvic fin	Thoracic	
	S: 1-1-1	R: 3-3-3
Dorsal fin	S: 8-8-9	R: 27-29-30
Pectoral fin	R: 13-14-14	
Anal fin	R: 16-19-21	
Gill rakers	U: X-X-X	L: X-X-X

LIFE HISTORY

Range	Cent. California, 34-38°N, to Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal, 0-110 m
ELH pattern	Oviparous; demersal, adhesive, attached eggs; pelagic larvae
Spawning	Season: Fall-spring ^a Area: Intertidal, in mussel zone ^a Mode: Migration: Females move from subtidal area to mussel zone to deposit eggs ^a
Fecundity	Range/function:
Age at first maturity	1 yr ^b
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	2.0-2.5 mm
No. of oil globules	One
Oil globule diameter	
Yolk	Orange
Envelope	
Hatch size	~9 mm TL (possibly as small as 7 mm TL)
Incubation time/temp.	
Pigment	

Diagnostic characters

- Wild: Only egg with bright orange yolk laid in *Mytilus* beds^a

LARVAE

Preanal length	~50% SL
Length at flexion	~9-11 mm SL
Length at transformation	~26 mm SL (largest pelagic specimen caught = 16-17 mm SL)

Sequence of fin development

Pigment

- Finfold pigment
- Distinctive pigment over lateral surface of trunk
- Pectorals pigmented in band posteriorly

Diagnostic characters (see Table 3)

- Long pigmented pectorals (of sample size = 2, length of pectorals ranged from 42 to 59% SL)
- Bumps and parietal ridge

The following information will aid in identification of *N. pribilovius* (larvae unknown)^c and *N. robustus* (eggs and larvae unknown).

	<i>N. pribilovius</i>	<i>N. robustus</i>
Total vertebrae	36-37	35
Dorsal fin spines	7-10	7- 8
Dorsal fin rays	22-26	19-21
Anal fin rays	15-20	14-15
Pectoral fin rays	15-17	14-16
Pelvic fin rays	3	3
Range	SE Alaska - Chukchi Sea	Wash. - Bering Sea

^aJ. Marliave, Vancouver Public Aquarium, P.O. Box 3232, Vancouver, B.C., Canada V6B 3X8, pers. commun., 16 Oct. 1986.

^bFitch and Lavenberg 1975

^cAndriashev 1954; eggs 2.5-2.7 mm.

Ref: Blackburn 1973, Marliave 1975a, Richardson and Washington 1980, Washington et al. 1984b.




Figure A, NWAFC original (B. Vinter); B–C, Richardson and Washington 1980.

AGONIDAE

Poachers and alligatorfishes range from Baja California to the Bering Sea, with the center of abundance in the North Pacific. They are mostly elongate with large fused bony plates covering the body. Pectoral fins are fanlike. Adults are demersal and occur from moderate depths to 1250 m. Some have also been found in tidepools (Hart 1973, Garrison and Miller 1982). The family is represented by 25 species and 15 genera within the study area. Among the few species studied, demersal adhesive eggs are attached to holdfasts of laminarians (Breder and Rosen 1966, Garrison and Miller 1982). Larvae are pelagic and are occasionally taken in plankton nets near the surface. Bony plates, characteristic of agonids, first appear in the larvae as spines. Settlement in some species may occur at approximately 2 months after hatching (Marliave 1975a). Descriptions of larvae of three species are available for inclusion here (*Agonomalus mozinoi*, *Bothragonus swani*, and *Xeneretmus latifrons*). Illustrations of single specimens are provided for *Stellerina*, *Hypsagonus*, *Ocella*, *Aspidophoroides* (species outside study area), and an unidentified agonid. Two illustrations are provided for Agonidae A, another unidentified agonid from the study area.


Table 39
Meristic characters of family Agonidae. All have pelvic counts of 1,2 and 6 branchiostegal rays.

Taxon	Distribution	Vertebrae		Fins			Gill rakers	
		Precaudal	Caudal	Dorsal	Anal	Pectoral	Upper	Lower
		(Total)					(Total)	
<i>Agonomalus mozinoi</i>	Cent. Calif.-Brit. Col.			VIII-IX,6-8	11-12	11-12		
<i>Agonopsis vulsa</i>	SSC-Gulf of Alaska		(39-42)	VIII-X,7-9	10-12	13-15		
<i>Agonus acipenserinus</i> ^a	N. Calif.-Chukchi Sea		(39-41)	VII-X,6-9	6-9	16-19		
<i>Agonus decagonus</i> ^a	Bering Sea-Arctic		(47-48) ^b	V-VII,5-8	6-8	13-16		
<i>Anoplagonus inermis</i>	N. Calif.-Aleutian Is.		(41-45)	4-6	4-5	8-10		
<i>Aspidophoroides bartoni</i>	Gulf of Alaska-Arctic		(51-53)	4-6	4-6	9-10		
<i>Aspidophoroides oltriki</i>	Bering Sea-Arctic		(38-40)	5-7	5-7	13-16		
<i>Bathyagonus alascanus</i>	N. Calif.-Bering Sea		(39-41)	V-VIII,5-8	6-8	15		
<i>Bathyagonus infraspinus</i>	N. Calif.-Bering Sea		(38-39)	V-VIII,5-8	5-8	15-16		
<i>Bathyagonus nigripinnis</i>	N. Calif.-Bering Sea		(44-45)	VI-VIII,6-7	7-9	15-16		
<i>Bathyagonus pentacanthus</i>	S. Calif.-Bering Sea		(40-46)	V-VIII,5-8	6-9	14-16		
<i>Bothragonus swani</i>	Cent. Calif.-Gulf of Alaska		(29-31)	II-V,4-6	4-5	10-12		
<i>Hypsagonus quadricornus</i>	Wash.-Bering Sea		(36)	IX-XI,5-7	9-11	12-14		
<i>Ocella dodecaedron</i>	Aleutian Is.-Chukchi Sea		(38-39)	IX-XI,7-8	14-16	14-15		
<i>Ocella impi</i> ^c	Brit. Col.		(37) ^b	IX,6	9	18		
<i>Ocella verrucosa</i>	Cent. Calif.-Bering Sea	13-14	21-24	VII-IX,6-9	7-12	14-15	1-2	8-12
<i>Odontopyxis trispinosa</i>	SSC-SE Alaska	10-12	27-30	III-VI,5-7	5-7	13-15		7
<i>Pallasina barbata</i>	Cent. Calif.-Bering Sea		(45-47)	V-IX,6-7	9-14	10-13		
<i>Percis japonicus</i>	Bering Sea		(42)	V-VII,6	7-9	12		
<i>Sarritor frenatus</i>	Brit. Col.-Bering Sea		(46-48)	VI-VIII,6-8	6-7	15-17		
<i>Sarritor leptorhynchus</i>	Gulf of Alaska-Bering Sea		(42-44)	VI-IX,5-8	6-8	13-15		
<i>Stellerina xyosterna</i>	SSC-Brit. Col.		(34-37)	VI-VIII,5-7	8-9	17-19	1-2	8-12 (10-14)
<i>Xeneretmus latifrons</i>	SSC-Brit. Col.	11-13	28-30	VI-VIII,6-8	6-9	13-15		10-11
<i>Xeneretmus leiops</i>	S. Calif.-SE Alaska		(39-42)	VI-VII,6-8	5-8	13-15		
<i>Xeneretmus triacanthus</i>	SSC-Brit. Col.	12	29-30	V-VII,6-7	5-7	12-14	1	8-13

^a Placement in the genus *Agonus* is questionable (Lea and Dempster 1982).

^b W.A. Laroche, 24 Maple Park, Box 216, Enosburg Falls, VT 05450, pers. commun., 10 Dec. 1986.

^c *Ocella impi* may not be a valid species, rather it may be the juvenile of *Stellerina xyosterna* (A.E. Peden, Brit. Col. Prov. Mus., Victoria, B.C., Canada V8V 1X4, pers. commun., 22 Jan. 1987).



Figures A–B, NWAFC originals (B. Vinter); C–G, Washington et al. 1984b (Figure C was misidentified in Washington et al. (1984b) as *Bothragonus swani*; G, Atlantic specimen).

MERISTICS

Vertebrae	Total: X-X-X
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 2-2-2
Dorsal fin	S: 8-X-9 R: 6-X-8
Pectoral fin	R: 11-X-12
Anal fin	R: 11-X-12
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Cent. California, 34-38°N, to Brit. Col., 48°30'-55°N
Ecology	Nearshore shelf demersal, intertidal to 11 m
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Winter-spring ^a Area: Mode: Repeated deposition of eggs in separate masses ^b Migration:
Fecundity	Range/function: 6-25 eggs/mass ^b
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	~1 mm
No. of oil globules	
Oil globule diameter	
Yolk	Red
Envelope	
Hatch size	5.5 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE^c**

Preanal length	
Length at flexion	
Length at transformation	
Sequence of fin development	
Pigment	
• Dorsal and ventral finfolds	
• Heavily pigmented body, pigment increasing with development	

Diagnostic characters

- Based on one specimen (8.2 mm SL), they are superficially similar to cottid *Hemitripteris* group (p. 448)

^aW.A. Laroche, 24 Maple Park, Box 216, Enosburg Falls, VT 05450, pers. commun., 10 Dec. 1986.

^bJ. Marliave, Vancouver Public Aquarium, P.O. Box 3232, Vancouver, B.C., Canada V6B 3X8, pers. commun., 16 Oct. 1986.

^cIncomplete series.




Figure A, Marliave 1978; B, NWAFC original (B. Vinter).

MERISTICS

Vertebrae	Total: 29-30-31 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 2-2-2
Dorsal fin	S: 2-3-5 R: 4-5-6
Pectoral fin	R: 10-12-12
Anal fin	R: 4-4-5
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Cent. California, 34-38°N, to Gulf of Alaska, 54-60°N
Ecology	Nearshore shelf demersal, intertidal to 18 m
ELH pattern	Oviparous; demersal, attached eggs; pelagic larvae
Spawning	Season: Winter-spring ^a Area: On kelp holdfasts ^b Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	2 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	7.5 mm TL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	
Length at flexion	~10-12 mm TL
Length at transformation	>16 mm TL
Sequence of fin development	
Pigment	
	<ul style="list-style-type: none"> • Upper and lower jaw • Dorsal head • Anterior gut and body • Outer edge of pectoral fin • Flexion and postflexion larvae develop several vertical bars on body


Diagnostic characters

- Large pigmented pectoral fins
- Body spines develop in flexion larvae
- Body short and stocky compared with *Xeneretmus latifrons*

^aW.A. Laroche, 24 Maple Park, Box 216, Enosburg Falls, VT 05450, pers. commun., 10 Dec. 1986.

^bMarliave 1975a

Ref: Marliave 1975a, Washington et al. 1984b.



Figures A–D, Marliave 1975a.

MERISTICS

Vertebrae	Total: 39-41-42 Precaudal: 11-12-13 Caudal: 28-29-30
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic S: 1-1-1 R: 2-2-2
Dorsal fin	S: 6-7-8 R: 6-7-9
Pectoral fin	R: 13-14-15
Anal fin	R: 6-7-9
Gill rakers	U: X-X-X L: 10-X-11

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epi- and mesobenthal, 18-400 m
ELH pattern	Oviparous, eggs probably demersal, pelagic larvae
Spawning	Season: Spring ^{a,b} Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	~7 mm TL
Incubation time/temp.	
Pigment	


Diagnostic characters**LARVAE**

Preanal length	
Length at flexion	~10 mm TL
Length at transformation	
Sequence of fin development	
Pigment ^c	<ul style="list-style-type: none"> • Body pigment extending into finfold in four vertical bars: Over posterior gut, midbody, at 3/4 body length, and tail region • Preflexion larvae have pigment on jaws

Diagnostic characters

- Four vertical pigment bars extending into finfolds
- Body spines develop in postflexion larvae
- Long slender body compared with *Bothragonus swani*

^aMarliave 1975a^bFitch and Lavenberg 1968^cJ. Marliave, Vancouver Public Aquarium, P.O. Box 3232, Vancouver, B.C., Canada V6B 3X8, pers. commun., 16 Oct. 1986.



Figures A–B, D, Marliave 1975a (D, length in Marliave 1975a is given as 21.0 mm TL [footnote c]) ; C, Washington et al. 1984b.

The family Cyclopteridae is composed of two subfamilies: the Cyclopterinae (lumpsuckers) and the Liparidinae (snailfishes). Although both subfamilies possess a ventral sucking disc, many differences distinguish the two groups (e.g., lumpsuckers have 2 dorsal fins and about 23-29 vertebrae, and snailfishes have a single dorsal fin and about 38-86 vertebrae).

Cyclopterinae

Cyclopterines are found exclusively in the cooler waters of the Northern Hemisphere. The subfamily is represented by eleven species in five genera in the Northeast Pacific and Bering Sea. Mostly benthic, adults may be found clinging to rocks in tidal zones or in rocky habitat as deep as 225 m (Hart 1973). Eggs, which are adhesive, 1.9-5.0 mm in diameter, and may contain one or more oil globules, are laid in nest sites within rocky crevices or shells and are guarded during incubation (Able et al. 1984). Newly hatched larvae may be identified as cyclopterines by their stout body shape, heavy pigmentation, large sucking disc on the ventral surface, and advanced state of development. They are found attached to substrate and prefer areas with current (A.C. Matarese and S.F. Borton, unpubl.). Larvae of only two species can be identified in our area, *Aptocyclus ventricosus* and *Eumicrotremus orbis*.

Table 40
Meristic characters of subfamily Cyclopterinae. All have pelvic discs and six branchiostegal rays.

Taxon	Distribution	Vertebrae		Fins			Gill rakers	
		Precaudal	Caudal	Dorsal	Anal	Pectoral	Upper	Lower
<i>Aptocyclus ventricosus</i>	Brit. Col.-Bering Sea	14	13-15	V,8-11	6-9	19-22		5-7
<i>Cyclopteropsis phrynoides</i>	Gulf of Alaska-Bering Sea			VII,11		25		
<i>Eumicrotremus andriashevi</i>	Bering Sea-Chukchi Sea			VI-VII,10-12	10-11	23-27		
<i>Eumicrotremus barbatus</i>	Aleutian Is.	11	16	VII,11	10	23		
<i>Eumicrotremus birulai</i>	Gulf of Alaska-Bering Sea	11	17	VI-VII,9-12	9-11	25-29		6-9
<i>Eumicrotremus gyrinops</i>	Aleutian Is.-Bering Sea			VIII,9	9	24		
<i>Eumicrotremus orbis</i>	Wash.-Bering Sea	10-11	17-18	V-VII,9-11	9-11	19-27		5-6
<i>Eumicrotremus soldatovi</i>	Bering Sea	11	18	VI,11-12	10	24		
<i>Eumicrotremus taranetzi</i>	Bering Sea	11	16	V-VI,9-10	9-10	24-26		6-7
<i>Lethotremus muticus</i>	Aleutian Is.-Bering Sea			VII,11	10	23		
<i>Pelagocyclus vitiazii</i>	Bering Sea			IV-V,9-10	8-9	19-21		

MERISTICS

Vertebrae	Total: 27-X-29
	Precaudal: 14-14-14
	Caudal: 13-X-15
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Disc
	S: X-X-X R: X-X-X
Dorsal fin	S: 5-5-5 R: 8-10-11
Pectoral fin	R: 19-20-22
Anal fin	R: 6-8-9
Gill rakers	U: X-X-X L: 5-X-7

LIFE HISTORY

Range	Brit. Col., 48°30'-55°N, to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathypelagic, 0-1500 m
ELH pattern	Oviparous; demersal, adhesive eggs; demersal larvae
Spawning	Season: Area: Mode: Males guard eggs ^a Migration:
Fecundity	Range/function: 3800 ^a
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	2.3-2.4 mm
No. of oil globules	(Present)
Oil globule diameter	
Yolk	
Envelope	
Hatch size	6.5-7.0 mm TL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	33% SL
Length at flexion	6.5-7.0 mm TL
Length at transformation	
Sequence of fin development	Pectorals and pelvic disc form before hatching
Pigment	

- Initially, body is lightly pigmented but with development pigment increases to cover entire body


Diagnostic characters

- Smooth body (lack of spines)
- Distinguished from other cyclopterines by
- Distribution
- Meristic characters:

Total vertebrae	27-29
Dorsal fin spines	5
Dorsal fin rays	8-11
Anal fin rays	6-9

^a Able et al. 1984

Ref: Able et al. 1984, Kobayashi 1962.



Figures A–D (D, ventral view), Kobayashi 1962 (A, reared from specimens collected near Hokkaido, Japan).

MERISTICS

Vertebrae	Total: 26-X-29
	Precaudal: 10-X-11
	Caudal: 17-X-18
Branchiostegal rays	6-6-6
Caudal fin	X, 5+4, X
	Total rays=9-12
Pelvic fin	Disc
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 5-6-7 R: 9-10-11
Pectoral fin	R: 19-23-27
Anal fin	R: 9-10-11
Gill rakers	U: X-X-X L: 5-X-6

LIFE HISTORY

Range	Washington, 46-48°30'N, to Chukchi Sea, north of 66°N
Ecology	Epi-, meso-, and bathybenthal, 0-575 m
ELH pattern	Oviparous; demersal, attached, adhesive eggs; demersal larvae
Spawning	Season: Fall-winter (Brit. Col.) ^a Area: Mode: Males guard eggs laid in nests ^b Migration:
Fecundity	Range/function: 305-1590 (737 at 52.5 mm)/ $F=47.67 \times L - 1766.86$, $L=TL \text{ mm}^b$
Age at first maturity	1 yr (males) ^b
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.9-2.2 mm
No. of oil globules	One to multiple
Oil globule diameter	
Yolk	Clear, homogeneous
Envelope	Translucent
Hatch size	4.5-4.7 SL
Incubation time/temp.	26 d/9.7-12.0°C
Pigment	

Diagnostic characters

- Degree of body pigment
- Precocious fin development

LARVAE

Preanal length	50-75% SL
Length at flexion	Prior to hatch
Length at transformation	6.9-8.7 mm SL
Sequence of fin development	Pelvics and caudal (prior to hatch), dorsal spines, dorsal rays, pectorals, anal (although caudal development is initiated prior to hatch, it is the last to complete development)

Pigment


- Initially, heavily pigmented except caudal peduncle, area around anus, first three anal rays, and snout

Diagnostic characters

- Pigment: Small melanophores covering most of body and dorsal fin
- Morphology: Globular shape, spines
- Precocious fin development

^aArita 1969^bA.C. Matarese and S.F. Borton, unpubl.

Ref: Matarese and Borton, unpubl.



Figures A–E (D, ventral view), Matarese and Borton, unpubl.

CYCLOPTERIDAE


Liparidinae

This large subfamily is circumpolar about both poles in distribution. Within the study area there are 74 described species within 14 genera. Adults are demersal or pelagic and have been found from tidepools to depths of 7000 m (Hart 1973, Able et al. 1984). Spawning habits are diverse. Adhesive clumps of eggs (1-8 mm) have been seen attached to algae, mollusc shells, and tubeworms; eggs and larvae of *Careproctus* spp. have been found in gill cavities of lithodid crabs (Anderson and Cailliet 1974). Parental care of eggs (i.e., hiding, paternal guarding, or both) is exhibited in most taxa studied. Most species studied, especially deep-water forms, hatch at an advanced stage of development (Able et al. 1984). Larvae are planktonic or benthic. Only one complete larval series is available from our area, *Rhinoliparis barbulifer*. Illustrations of two unidentified larvae are provided since each of them is relatively common in our collection. The small larva (4.0 mm SL) is common in our Gulf of Alaska collections whereas the larger larva (5.8 mm SL) is routinely collected in our surveys along the Pacific Coast and represents one of the ten most abundant groups of ichthyoplankton collected off Oregon (Mundy, pers. commun.)¹. Representatives of specimens from other areas (Atlantic, Arctic) are presented for three genera: *Careproctus*, *Paraliparis*, and *Liparis*. Early juvenile specimens of *Paraliparis* sp. (probably *P. deani*) are occasionally collected in plankton nets. Illustrations of a tentatively identified partial series of *Nectoliparis pelagicus* are also provided.

¹B. Mundy, NMFS Southwest Fish. Cent., Honolulu Lab., 2570 Dole St., Honolulu, HI 96822-2396, pers. commun., 1 Oct. 1986.

Unidentified Liparidinae

A



B





Figure A, NWAFC original (B. Vinter); B, Able et al. 1984.

LIPARIS
L. fabricii




Figures A–C (B, ventral view), Able et al. 1984 (Arctic specimens).

CAREPROCTUS


C. reinhardti

A



12.6 mm SL

B




12.6 mm SL

PARALIPARIS

P. copei


C




24.0 mm SL

P. calidus

D



E



12.9 mm SL

Figures A–E (B, E, ventral views), Able et al. 1984 (Atlantic specimens).

CYCLOPTERIDAE

Table 41
Meristic characters of subfamily Liparidinae.^a

Taxon	Distribution	Vertebrae		Fins				Gill rakers		Branchiostegals
		Precaudal (Total)	Caudal	Dorsal	Anal	Pectoral	Pelvic	Upper (Total)	Lower	
<i>Acantholiparis</i> ^b	Bering Sea						Absent		6	
<i>Acantholiparis caecus</i>	Oregon	8-10	43-46	48-52	43-45	21	Absent	(10-11)	6-7	
<i>Acantholiparis opercularis</i>	Calif.-Bering Sea	9	41	45-52	38-47	20-24	Absent	(8-10)	6	
<i>Careproctus abbreviatus</i>	SE Alaska-Bering Sea			39	32	21	Disc		6	
<i>Careproctus attenuatus</i>	Bering Sea			48	40	34	Disc		6	
<i>Careproctus bowersianus</i>	Bering Sea	9-10	47-50	51-54	46-48	34-38	Disc		6	
<i>Careproctus cameliae</i>	Bering Sea			50	45	28	Disc		6	
<i>Careproctus canus</i>	Aleutian Is.	11-12 (55-58)	44-47	51-53	43-46	33-36	Disc		6	
<i>Careproctus colletti</i>	Aleutian Is.-Bering Sea			52-58	47-52	25-31	Disc		6	
<i>Careproctus cypselurus</i>	Oregon-Bering Sea	8 (64-70)	57-59	58-64	52-58	32-37	Disc		6	
<i>Careproctus ectenes</i>	Aleutian Is.-Bering Sea			48	44	30-32	Disc		6	
<i>Careproctus filamentosus</i>	Oregon	9-10 (63-68)	54-58	58-63	51-55	21-24	Disc		6	
<i>Careproctus furcellus</i>	Bering Sea	10-11 (66-71)	56-61	60-65	54-59	32-37	Disc		6	
<i>Careproctus gilberti</i>	Cent. Calif.-Bering Sea	8-9 (55-58)	47-49	45-55	41-48	30-33	Disc		6	
<i>Careproctus longifilis</i>	SSC-Oregon	7-10 (55-58)	46-50	50-54	44-48	17-23	Disc		6	
<i>Careproctus melanurus</i>	SSC-Bering Sea	8-11 (57-63)	49-53	53-58	47-51	27-33	Disc		6	
<i>Careproctus microstomus</i>	Oregon	9-10 (67-69)	58-59	61-67	54-60	22-27	Disc		6	
<i>Careproctus mollis</i>	Bering Sea			51	47	35	Disc		6	
<i>Careproctus opisthotremus</i>	Aleutian Is.-Bering Sea			46	36	32	Disc		6	
<i>Careproctus oregonensis</i>	Oregon	8-10 (65-69)	57-60	61-67	55-57	19-23	Disc		6	
<i>Careproctus ostentum</i>	Aleutian Is.-Bering Sea			54	47	32	Disc		6	
<i>Careproctus ovigerum</i>	Oregon-Brit. Col.	10-12 (47-49)	35-39	43-45	34-37	31-34	Disc		6	
<i>Careproctus pellucidus</i>	SE Alaska			52-55	45-48	33-35	Disc		6	
<i>Careproctus phasma</i>	Bering Sea			53	45	34	Disc		6	
<i>Careproctus pycnosoma</i>	Bering Sea	10 (46-49)	36-39	42-45	36-39	38-39	Disc		6	
<i>Careproctus rastrinus</i>	SE Alaska-Bering Sea	8-9 (59-64)	50-52	55-59	49-52	33-37	Disc		6	
<i>Careproctus scottae</i>	SE Alaska-Bering Sea			52-56	47-51	32-34	Disc	2	10	
<i>Careproctus simus</i>	Aleutian Is.-Bering Sea	10-11	48-52	54-58	47-51	31-37	Disc		6	
<i>Careproctus spectrum</i>	SE Alaska-Bering Sea			52	47	32	Disc		6	
<i>Careproctus zachirus</i>	Aleutian Is.	12-13 (56-58)	44-46	51-58	43-45	28-31	Disc		6	
<i>Crystallichthys cyclospilus</i>	Gulf of Alaska-Bering Sea	11 (52-53)	42	48-50	42-43	33-36	?		6	
<i>Crystallichthys mirabilis</i>	Bering Sea			53	44	30-33	?		6	
<i>Elassodiscus caudatus</i>	Cent. Calif.-SE Alaska	9-11	46-51	49-55	41-50	27-29	Disc		6	
<i>Elassodiscus tremebundus</i>	Bering Sea	9-12 (62-74)	52-63	55-67	49-60	25-33	Disc		6	
<i>Gyrinichthys minytremus</i>	Aleutian Is.-Bering Sea			25	14	25	Disc		6	
<i>Liparis Bristolensis</i>	Bering Sea-Chukchi Sea		(49)	38-40	30-35	33-37	Disc		6	
<i>Liparis callyodon</i>	Wash.-Bering Sea		(41-42)	33-35	25-27	28-31	Disc		6	

Table 41 (continued)

Taxon	Distribution	Vertebrae		Fins				Gill rakers		Branchiostegals
		Precaudal (Total)	Caudal	Dorsal	Anal	Pectoral	Pelvic	Upper (Total)	Lower	
<i>Liparis catharus</i>	SE Alaska	12	38	46	36	37	Disc			6
<i>Liparis cyclopus</i>	Oregon-Bering Sea	(42-44)		35-37	29-31	29-32	Disc			6
<i>Liparis dennyi</i>	Wash.-Aleutian Is.	(44-45)		37-40	30-34	36-39	Disc			6
<i>Liparis florae</i>	S. Calif.-Bering Sea	(39-40)		31-33	25-27	29-33	Disc	0-1	3-4	
<i>Liparis fucensis</i>	Cent. Calif.-SE Alaska	10-11	29-30	33-35	27-29	37-43	Disc	(8-9)		6
<i>Liparis gibbus</i>	SE Alaska-Arctic	10-12	34-38	38-46	32-37	37-45	Disc	(6-10)		6
<i>Liparis grebnitzki</i>	Bering Sea			32	27	29	Disc			6
<i>Liparis mednius</i>	Bering Sea			29 ^c	27	27	Disc			6
<i>Liparis megacephalus</i>	Bering Sea			43-44	36	36-38	Disc			6
<i>Liparis micraspidophorus</i>	Aleutian Is.-Bering Sea			31-32	25-27	30-32	Disc			6
<i>Liparis mucosus</i>	SSC-SE Alaska	(36-38)		28-32	22-25	27-32	Disc			6
<i>Liparis ochotensis</i>	Gulf of Alaska-Bering Sea			45	36-38	39-42	Disc			6
<i>Liparis pulchellus</i>	Cent. Calif.-Bering Sea	11-12	40-42	47-53	39-42	36-37	Disc		6-9	6
<i>Liparis rutteri</i>	N. Calif.-Bering Sea	(37-39)		30-32	23-27	30-33	Disc			6
<i>Liparis tunicatus</i>	Bering Sea-Arctic	10-12	35-38	39-44	33-37	32-38	Disc		5-11	6
<i>Lipariscus nanus</i>	Cent. Calif.-SE Alaska			40-52	37-49	13-15	Disc			5
<i>Nectoliparis pelagicus</i>	Cent. Calif.-Bering Sea	9-12	50-52	44-56	40-51	19-25	Absent			5
		(61-64)								
<i>Odontoliparis ferox</i>	Oregon	11	48	51	46	17	Absent			6
<i>Osteodiscus cascadae</i>	Oregon-Brit. Col.	8-9	43-47	47-52	40-44	20-25	Absent			6
		(51-56)								
<i>Paraliparis cephalus</i>	S. Calif.-Bering Sea	9-10	47-54	50-57	44-51		Absent			6
<i>Paraliparis dactylosus</i>	Cent. Calif.-Bering Sea	(59-61)		54-56	49-51	28-30	Absent			6
<i>Paraliparis deani</i>	N. Calif.-SE Alaska			56-58	44-48	18-22	Absent			6
<i>Paraliparis holomelas</i>	Bering Sea			58-61	54	23	Absent			6
<i>Paraliparis latifrons</i>	SSC-Oregon	9-10	51-52	54-57	48-50	21-24	Absent			6
		(61)								
<i>Paraliparis megalopsis</i>	Oregon	9	67	66-71	63-65	16-19	Absent			6
		(76)								
<i>Paraliparis melanobranchus</i>	Oregon			60	53	17	Absent			6
<i>Paraliparis mento</i>	Cent. Calif.-Wash.	9-10	51-52	55-59	49-51	16-18	Absent			6
<i>Paraliparis paucidens</i>	Oregon-Brit. Col.	10-12	55-56	58-60	53-54	19-24	Absent			6
		(66-67)								
<i>Paraliparis pectoralis</i>	Oregon-Bering Sea	10-11	51-54	55-58	49-52	28-32	Absent			6
		(61-64)								
<i>Paraliparis rosaceus</i>	SSC-Brit. Col.	11-13	56-61	57-69	53-60	18-22	Absent			6
		(67-74)								
<i>Paraliparis ulochir</i>	SSC-Bering Sea	9-10	62-65	65-69	60-64	21-24	Absent			6
		(72-74)								
<i>Polypera beringiana</i>	Wash.-Bering Sea			38-39	31-32	36-37	Disc			6
<i>Polypera greeni</i>	Brit. Col.-Bering Sea	(47)		37-40	31-32	33-37	Disc			6
<i>Rhinoliparis attenuatus</i>	Cent. Calif.-Bering Sea	(80-83)		74-78	70-73	21-25	Absent			6
<i>Rhinoliparis barbulifer</i>	S. Calif.-Bering Sea	(68)		63-68	57-59	18-21	Absent			6
<i>Tennocora candida</i>	Gulf of Alaska-Bering Sea	(68)		45-48	39	33-37	Disc			6

^aThe NWAFC meristic database was updated for the Liparidinae by the papers of Kido (1983, 1984, 1985), Kido and Kitagawa (1986), and unpublished data from D. Stein (Oregon St. Univ., Corvallis, OR 97331, pers. commun., 29 Oct. 1986).

^bAt least two undescribed species in addition to *A. opercularis* probably occur in the Bering Sea.

^cAccording to Soldatov and Lindberg (1930), *Liparis mednius* has six dorsal spines.

MERISTICS

Vertebrae	Total: 61-X-64 Precaudal: 9-X-12 Caudal: 50-X-52
Branchiostegal rays	5-5-5
Caudal fin	
Pelvic fin	Absent
Dorsal fin	R: 44-X-56
Pectoral fin	R: 19-20-25
Anal fin	R: 40-X-51
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Cent. California, 34-38°N, to Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal
ELH pattern	Oviparous, eggs probably demersal, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE^a**


Preanal length Changes with development
Length at flexion
Length at transformation
Sequence of fin development
Pigment
• Darkly pigmented peritoneum
• Ventral half of head pigmented, appearing bearded

Diagnostic characters


- Presence of two separate pectoral fin lobes
- General tadpole shape
- Position of anus: Apparently moves forward with development; in adults anus is directly under eye
- Absence of disc
- Number of branchiostegal rays (five)

^aIdentification of small larvae prior to pectoral fin development and movement of anus forward is tentative. A cleared and stained 7.9-mm SL specimen has five branchiostegal rays.

A



B



Figures A–B, NWAFC originals (B. Vinter).

MERISTICS

Vertebrae	Total: X-X-X Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Absent
Dorsal fin	R: 56-X-58
Pectoral fin	R: 18-X-22
Anal fin	R: 44-X-48
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	N. California, 38-42°N, to SE Alaska, 55-59°N
Ecology	Epi-, meso-, and bathybenthal, 55-1008 m
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Continuous and periodic spawning occurs with other <i>Paraliparis</i> spp. ^a Area: Probably under rocks or in/on invertebrates with hard exoskeletons ^a Mode: Parental care (egg guarding) likely for periodic spawners ^a Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	~2 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE^b**

Preanal length
Length at flexion
Length at transformation
Sequence of fin development
Pigment


Diagnostic characters

- Characteristic notch in pectoral fin
- General body shape
- Absence of disc

^aStein 1980b^bJuvenile specimens only, probably *P. deani*.

Ref: Able et al. 1984, Hart 1973.

A



~ 28.5 mm SL

Figure A, NWAFC original (B. Vinter).

MERISTICS

Vertebrae	Total: 68-68-68 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	6-6-6
Caudal fin	X, 3, X
Pelvic fin	Absent
Dorsal fin	S: 63-X-65 R: X-X-X
Pectoral fin	R: 18-X-20
Anal fin	S: 57-X-59 R: X-X-X
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Meso- and bathybenthal
ELH pattern	Parity unknown, eggs probably demersal, benthopelagic larvae ^a
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	2.5 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	≤9.1 mm SL
Incubation time/temp.	
Pigment	• Unpigmented

Diagnostic characters

- Yolk absorbed by 11.7 mm SL
- Flexion and fin ray development occurs prior to hatching

LARVAE (postflexion and juvenile)


Preanal length	<50% SL
Length at flexion	<9.1 mm SL
Length at transformation	
Sequence of fin development	Dorsal, anal, and caudal; pectoral
Pigment	• Larvae <11.0 mm SL have no pigment • Nape, peritoneum, and lateral body pigmented by 12.5 mm SL • Pigment on the head and body increases in larger specimens

Diagnostic characters


- Loose skin over body
- Morphology
 - Depressed head
 - Large snout
 - Slender tapering body
- Absence of disc

^aKido and Kitagawa 1986

Ref: Able et al. 1984, Kido and Kitagawa 1986.



Figures A–C (A, yolksac larva; B–C, juveniles), Kido and Kitagawa 1986 (specimens from Iwate Prefecture, Honshu, Japan).



Perciformes

The most diversified of all fish orders is the Perciformes. Although the order encompasses the entire range of fish forms and behavior, most species are adapted for life as predators in shallow or surface waters. More than a dozen characters (many secondarily lost) define this group of spiny-rayed fishes. Most obvious of these are fin spines, two dorsal fins, scales generally ctenoid, pelvic fin inserted forward of the abdomen, and vertical insertion of the pectoral fin. There are 22 suborders, 150 families, 1,367 genera, and about 7,800 species within the order (J. Nelson 1984). The most speciose groups are the percoids and zoarcoids. Within the study area, nine suborders are represented; four of them are composed of a single species. Most species are members of the families Stichaeidae and Zoarcidae.

Some early-life-history stages are known for most families within our area except for Zoarcidae and Scytalinidae. When describing the sequence of fin ray development for percoids, we used the patterns A-F summarized in G.D. Johnson (1984) (see also Introduction, p. 29). A summary of meristic characters of zoarcids is provided to assist in identification.

Families in study area:	Carangidae	Zaproridae
	Bramidae	Scytalinidae
	Caristiidae	Trichodontidae
	Sciaenidae	Clinidae
	Pentacerotidae	Icosteidae
	Bathymasteridae	Ammodytidae
	Zoarcidae	Gobiidae
	Stichaeidae	Trichiuridae
	Cryptacanthodidae	Scombridae
	Pholididae	Luvaridae
	Anarhichantidae	Stromateidae
	Ptilichthyidae	Centrolophidae
		Tetragonuridae

MERISTICS

Vertebrae	Total: 25-25-25
	Precaudal: 10-10-10
	Caudal: 15-15-15
Branchiostegal rays	7-X-8
Caudal fin	9-10, 9+8, 8-9
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 4-X-7 R: 24-X-29
Pectoral fin	R: X-X-X
Anal fin	S: 2-X-3 ^a R: 15-X-18
Gill rakers	U: 5-X-8 L: 12-X-19

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epipelagic
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	~1.3 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	Smooth
Hatch size	
Incubation time/temp.	24-48 hr/18-30°C
Pigment	

Diagnostic characters

LARVAE

Preanal length	~50-75% SL
Length at flexion	>4.1 mm SL
Length at transformation	
Sequence of fin development	“A” pattern: ^b 2nd dorsal (rays), anal, and caudal simultaneously followed by 1st dorsal (spines), pelvics, and pectorals (some carangids have precocious pelvics)

Pigment

- Uniform heavy pigmentation
- Body: Dorsolateral, lateral midline, and scattered on ventrolateral
- Vomer and branchiostegal membrane
- Antimedial rows on dorsal body margin present or absent^c

Diagnostic characters

- Relatively deep bodied
- Uniform heavy pigment except in caudal region
- Pigment on branchiostegal membrane
- 25 myomeres
- Flexion and postflexion
 - Large posttemporal and supracleithral, supraocular and preopercular spines (not serrate)
 - No supraoccipital crest

^aThe first two spines are physically separated from the third, shorter spine which is associated with the anal fin soft rays.

^bG.D. Johnson 1984

^cLaroche et al. 1984

Ref: Laroche et al. 1984, Sanzo 1931b.




Figure A, Laroche et al. 1984 (Gulf of Mexico specimen); B–C, Sanzo 1931b (eastern Atlantic specimens).

MERISTICS

Vertebrae	Total: 23-24-25	
	Precaudal: 10-10-10	
	Caudal: 14-14-14	
Branchiostegal rays	7-X-8	
Caudal fin	9-10, 9+8, 9-10	
Pelvic fin	Thoracic	
	S: 1-1-1	R: 5-5-5
Dorsal fin	S: 8-X-9	R: 28-X-38
Pectoral fin	R: X-X-X	
Anal fin	S: 2-X-3 ^a	R: 22-X-33
Gill rakers	U: 7-X-15	L: 25-X-42

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesopelagic, 0-403 m ^b
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Jan-Nov; ^c Mar-Aug (California) ^d Area: Epipelagic, offshore ^e Mode: Migration:
Fecundity	Range/function: 53,000 ^b
Age at first maturity	2-3 yr ^e
Longevity	>30 yr ^e

^aThe first two spines are physically separated from the third, shorter spine which is associated with the anal fin soft rays.

^bHart 1973

^cFrey 1971

^dMacCall and Stauffer 1983

^eFitch and Lavenberg 1971

^fRemains within this range from 2.2 to 50.0 mm SL.

^gG.D. Johnson 1984

Ref: Ahlstrom and Ball 1954, Laroche et al. 1984.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	0.90-1.08 mm
No. of oil globules	One
Oil globule diameter	0.25 mm
Yolk	Segmented
Envelope	Smooth, clear
Hatch size	1.91-2.38 mm SL
Incubation time/temp.	24-48 hr/18-30°C
Pigment	
	• Oil globule
	• Dorsal and ventral melanophores

Diagnostic characters

- Anterior position of oil globule in yolksac larvae
- Pigment
- Lack of yolk pigment distinguishes eggs from *Merluccius productus* (p. 186) and *Scomber japonicus* (p. 554)

LARVAE


Preanal length	52-61% SL ^f
Length at flexion	8-11 mm SL
Length at transformation	16 mm SL
Sequence of fin development	"A" pattern: ^g 2nd dorsal (rays), anal, and caudal simultaneously followed by 1st dorsal (spines), pelvics, and pectorals (some carangids have precocious pelvics)

Pigment

- Crown, dorsal and ventral midline
- Mediolateral streak
- Some superficial lateral pigment develops

Diagnostic characters

- Low myomere count (usually 24)
- Dorsal body margin pigment; antimedial rows absent with median rows only
- Flexion and postflexion
 - ~9 preopercular spines (not serrate)
 - Supraoccipital crest present



Figures A–F, Ahlstrom and Ball 1954.

MERISTICS

Vertebrae	Total: 39-40-41 Precaudal: 15-16-17 Caudal: 23-24-26
Branchiostegal rays	7-X-8
Caudal fin	8, 9+8, 7
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin ^a	S: 3-3-5 R: 30-X-35 D1+D2=33-36
Pectoral fin	R: 21-22-23
Anal fin ^a	S: 2-X-3 R: 25-X-29 A1+A2=27-30
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epipelagic, 0-200 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Spring (California) ^b Area: Pelagic ^b Mode: Migration: To the south to spawn ^b
Fecundity	Range/function:
Age at first maturity	
Longevity	>6 yr ^b

^aSince spines are weak or lacking in fins, there has been much confusion in the literature. According to data collected by G.D. Johnson (Natl. Mus. Nat. Hist., Wash., D.C., 20560, pers. commun., 7 Nov. 1986), all dorsal and anal elements are soft rays. Total counts from Mead (1972) are therefore more useful and are presented here.

^bFitch and Lavenberg 1971

^cG.D. Johnson 1984

^dG.D. Johnson, pers. commun., 7 Nov. 1986.

Ref: Mead 1972.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.56-1.60 mm
No. of oil globules	One
Oil globule diameter	0.40×0.32 mm
Yolk	
Envelope	
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	<50% SL
Length at flexion	By 7.4 mm SL
Length at transformation	
Sequence of fin development	Both "C" and "D" patterns ^c described for the family are not present in this species ^d

Pigment


- Gut and anterior body
- Crown
- With development along hypural margin
- Pectoral fin fringes

Diagnostic characters

- Pigment covers anterior body
- Body depth: Deep upper body
- Large pectoral fin
- Patterns of preopercular spines are useful diagnostic characters for bramid larvae

Taractes asper larvae are unknown. The following information may aid in their identification.

Total vertebrae	41-42
Caudal	17-18
Precaudal	23-24
Dorsal fin (D1+D2)	31-34
Anal fin (A1+A2)	23-26
Pectoral fin rays	18-20
Pelvic fin	I,5
Range	Cent. Calif.- Gulf of Alaska



Figures A–C, Mead 1972 (eyes redrawn).

MERISTICS

Vertebrae	Total: 37-38-40 Precaudal: 16-16-16 Caudal: 19-19-19
Branchiostegal rays	7-7-7
Caudal fin	6-7, 9+8, 6
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin ^a	D1+D2=32-34
Pectoral fin	R: 14-X-19
Anal fin ^a	A1+A2=20-22
Gill rakers	U: 6-6-6 L: 16-16-16

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Mesopelagic, 305-610 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length
Length at flexion Between 5.8 and 10.1
 mm SL
Length at transformation
Sequence of fin "A" pattern:^b 2nd dorsal
development (rays), anal, and caudal
 simultaneously followed
 by 1st dorsal (spines),
 pelvics, and pectorals

Pigment

- Postanal bar pattern
- Head and mediolateral gut

Diagnostic characters

- Myomeres (>35)
- Distinct pigment pattern with bars
- Head and body shape

^aThere is confusion in the literature whether spines are present in the dorsal (1-3) and anal (1) fins. According to data collected by G.D. Johnson, all dorsal and anal elements are soft rays (Natl. Mus. Nat. Hist., Wash., D.C. 20560, pers. commun., 7 Nov. 1986). Total counts are therefore more useful and are presented here.

^bG.D. Johnson 1984




Figure A, G.D. Johnson 1984.

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: 10-11-12 Caudal: 14-15-16
Branchiostegal rays	7-X-8
Caudal fin	X, 9+8, X (15-17 secondary)
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-X-16 R: 18-22-25
Pectoral fin	R: 16-17-19
Anal fin	S: 2-2-2 R: 10-11-12
Gill rakers	U: 9-X-12 L: 17-X-21

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Nearshore shelf pelagic
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Winter-spring ^a Area: Pelagic ^b Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	0.85 mm
No. of oil globules	1-3, coalesce early to 1
Oil globule diameter	0.23 mm
Yolk	Homogeneous, becoming pigmented
Envelope	Transparent, smooth
Hatch size	1.5-2.0 mm SL
Incubation time/temp.	52 hr/20°C
Pigment	<ul style="list-style-type: none"> • Yolk and oil globule • Embryonic pigment increases on head and trunk (dorsally and dorsolaterally)

Diagnostic characters**LARVAE**

Prenatal length	38-53% SL
Length at flexion	5.4-6.4 mm SL
Length at transformation	>12.7 mm SL
Sequence of fin development	2nd dorsal (rays), anal, 1st dorsal (spines), pelvics

Pigment


- Presence of nape melanophore
- Melanophore above hindgut absent or small
- 2-21 ventral melanophores; number increasing with development

Diagnostic characters

Among the fish larvae occurring in the study area, *G. lineatus* most closely resembles *Scomber japonicus* (p. 554). The myomere counts will usually separate them.

<i>G. lineatus</i>	26
<i>S. japonicus</i>	30-31

^aHart 1973^bWatson 1982



Figures A–F, Watson 1982.

MERISTICS^b

Vertebrae	Total: 24-X-25
	Precaudal: 12-X-13
	Caudal: 13-13-13
Branchiostegal rays	7-7-7
Caudal fin	7, 9+8, 5-6
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 13-X-15 R: 8-X-10
Pectoral fin	R: 16-X-19
Anal fin	S: 3-4-5 R: 6-X-9
Gill rakers	U: 5-X-8 L: 14-X-20

LIFE HISTORY

Range	N. California, 38-42°N, to Gulf of Alaska, 54-60°N
Ecology	Epi- and mesopelagic, 0-402 m
ELH pattern	Oviparous, pelagic eggs, larvae probably pelagic ^c
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

^a *Pentaceros* = *Pseudopentaceros* (Hardy 1983). J. Nelson (1984) does not recognize the genus *Pseudopentaceros* and cites Smith (1964) who included *Pseudopentaceros* in the synonymy of *Pentaceros*. Hardy (1983) removed *Pseudopentaceros* from synonymy but his work is not cited in J. Nelson (1984). The genus *Pseudopentaceros* may include up to three species:

Pseudopentaceros richardsoni—Restricted to Southern Hemisphere

P. wheeleri—North Pacific Ocean from Japan to Hawaii

P. pectoralis—North Pacific, Hawaii to Aleutian Is. (overlaps with *P. wheeleri* in central Pacific)

Hardy (1983) describes *P. wheeleri* as the most slender form. He suggests they are the "slender forms" referred to by Zama et al. (1977a) in their description of *Pentaceros richardsoni*. Hardy does not synonymize their "high-bodied" forms with *P. wheeleri*. The juvenile form described by Zama et al. (1977a) may include *P. wheeleri* in part. Hardy (1983) names this species as the one targeted on the Hawaiian ridge for intense commercial fishing. *Pseudopentaceros wheeleri* and *P. pectoralis* are probably the same species at different ontogenetic stages. Most likely two species occur worldwide (Humphreys et al. 1989).

^b Hardy 1983; includes meristic data for *Pseudopentaceros richardsoni*, *P. pectoralis*, and *P. wheeleri*.

^c Juveniles collected at surface to at least 260 mm FL; the shift from pelagic to bottom life may be ambivalent, occurring between 260 and 300 mm FL.

Ref: Hardy 1983, Humphreys et al. 1989, G.D. Johnson 1984, Zama et al. 1977b.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length
Length at flexion <9 mm SL
Length at transformation
Sequence of fin
development
Pigment (early juvenile *P. richardsoni*)

- Entire body heavily pigmented with intense patches along dorsal body midline and on spinous dorsal, anal, and pelvic fins

Diagnostic characters (early juvenile *P. richardsoni*)

- Pigment pattern
- Bony cranial projections, spines by 10-16 mm SL; most prominent are the following: median supraoccipital; posttemporal; lateral expansions which develop over orbit, becoming highly serrated along the edge with development; preopercular
- Anteriorly serrated pelvic spine
- With development there is a reduction in spination




Figure A, G.D. Johnson 1984 (southern hemisphere specimen).

BATHYMASTERIDAE

This endemic North Pacific family occurs throughout the study area where four species from two genera are found. Ronquils are elongate with long dorsal and anal fins composed almost entirely of soft rays. Little is known of the adult habits in the family except for some members of the genus *Rathbunella*, occurring from south of Point Conception to northern California, which inhabit rocky areas 10-92 m deep (and are occasionally taken with trawl nets). Courtship of *Rathbunella* is paired and may result in spawning occurring over a protracted period of time. Demersal eggs, which measure 0.9-1.1 mm and contain one oil globule, are deposited in a nonadhesive mass that is guarded by the male (Fitch and Lavenberg 1975, NWAFC unpubl.).

Pelagic larvae of *Ronquilus jordani* are collected in ichthyoplankton surveys conducted off Alaska and along the Pacific coast to northern California. *Bathymaster* spp. (6-40 mm SL) are routinely collected in the Gulf of Alaska and in the Bering Sea. *Ronquilus* larvae can be separated from *Bathymaster* larvae by their lack of pigment around the urostyle. *Bathymaster* spp. larvae are presently not identifiable to species. Meristic characters offer some potential for larger larvae (e.g., *B. signatus* usually has higher counts, especially vertebrae and total caudal fin rays). Before their fin rays develop, bathymasterid larvae may often be confused with stichaeid larvae. Most stichaeids (except members of the Stichaeini tribe) have a myomere count >55, whereas bathymasterids have a count <55. *Ronquilus jordani* larvae (myomeres 49-50) most closely resemble *Stichaeus punctatus* larvae (myomeres 51-55) due to the presence of distinctive lateral pigment along the hypaxial myomeres. *Bathymaster* spp. larvae (myomeres 49-55) most closely resemble those of *Bryozoichthys-Chirolphis* (myomeres about 60-75) due to the presence of pigment around the urostyle.



Table 42
Meristic characters of family Bathymasteridae.^a All have pelvic fin counts of I,5, 7+7 principal caudal fin rays, and 6 branchiostegal rays.

Taxon	Distribution	Vertebrae		Fins						
		Precaudal	Caudal (Total)	Dorsal ^b	Anal ^b	Pectoral	Caudal		Gill rakers	
							Upper	Lower	Upper	Lower
<i>Bathymaster caeruleofasciatus</i>	Brit. Col.-Bering Sea	14-15-16	35-38-39 (50-53-53)	44-47-48	33-35-36	16-18-19			5	12-14
<i>Bathymaster leurolepis</i>	Gulf of Alaska-Bering Sea	14-15-15	35-36-37 (49-51-52)	45-46-47	32-34-34	17-18-19	10	10	4-5	12-14
<i>Bathymaster signatus</i>	Wash.-Arctic	15-16-16	34-37-39 (54-54-55)	46-48-49	33-34-36	19-20-21	10-12	9-11		15-18
<i>Ronquilus jordani</i>	N. Calif.-Bering Sea	13-14-15	34-36-37 (49-50-50)	44-45-46	33-35-35	17-17-19	5-7	5-7		



^aFor some characters modal values are listed between range of values.

^bTotal counts include weak anterior spines; *Bathymaster* spp. usually possess two dorsal spines and one anal spine, and *Ronquilus* usually has one dorsal and one anal spine.

Bathymaster A



Ronquilus jordani



Figures A–D, NWAFC originals (B. Vinter).

ZOARCIDAE

Eelpouts are found chiefly in colder marine waters of the Northern Hemisphere. They have elongate, tapered bodies with long dorsal and anal fins confluent with the caudal fin. The pelvic fins are small and jugular when present. The head is large and the mouth is often big with thick lips. Within the study area there are 48 species within 14 genera. Adults are found from the intertidal zone to depths of more than 1900 m (Hart 1973). Although members of the genus *Zoarces* are viviparous, all species in our area, where reproductive mode is known, are oviparous. Oviparous species have been observed guarding their eggs by wrapping themselves around the egg mass. Demersal eggs are adhesive, spherical, and possess one oil globule. Diameters range from 1.7 to 9.0 mm. Newly hatched larvae are quite advanced and strongly resemble adult zoarcids (Anderson 1984b). Larvae of only five taxa have been illustrated; four are from outside the study area. Larvae probably become demersal or semidemersal soon after hatching because they are virtually never collected in plankton nets.

Table 43
Meristic characters of family Zoarcidae (taxonomy after Anderson 1984a,b).





Taxon	Distribution	Vertebrae		Fins				Gill rakers		Branchiostegals
		Precaudal	Caudal (Total)	Dorsal	Anal	Pectoral	Pelvic ^a	Upper (Total)	Lower	
<i>Bothrocara brunneum</i>	SSC-Bering Sea	22	94	107-112	92-96	14-17	ab	3-5	14-15	
<i>Bothrocara hollandi</i>	Bering Sea			114-117	94-99	15-17	ab	4	11	6
<i>Bothrocara molle</i>	SSC-Bering Sea		(120)	100-112	89-101	13-14	ab		(22)	
<i>Bothrocara pusillum</i>	SE Alaska-Brit. Col.	18-20	95-101	113-121	100-107	14-17	ab			
<i>Bothrocara remigerum</i>	Cent. Calif.-Wash.	23	96	107-117	93-94	13-16				
			(116)							
<i>Derepodichthys alepidotus</i>	SSC-Brit. Col.	22-26	92-98	110-116	94-101	10-11	3	0-1	11-12	6
<i>Gymnelis hemifasciatus</i>	Gulf of Alaska-Arctic	18-21	65-77	80-92 ^b						
			(85-95)							
<i>Gymnelis popovi</i>	Gulf of Alaska-Bering Sea			101	89		ab	3	12	
<i>Gymnelis viridis</i>	Aleutian Is.-Arctic		(87-99)	92-93	74	10-12	ab			
<i>Krusensterniella pavlovskii</i>	Bering Sea	20-21	88-89	70-75		12	ab			
<i>Lycenchelys altus</i>	Aleutian Is.	21	67	83	68	18		1	7	
			(88)						(8)	
<i>Lycenchelys camchaticus</i>	SSC-Bering Sea	21-24	97-103	112-117	98-105	13-17		2	13-16	6
									(14-18)	
<i>Lycenchelys crotalinus</i>	S. Calif.-Bering Sea	22-24	98-107	113-123	99-109	14-17		1	14-18	6
<i>Lycenchelys hippopotamus</i>	Bering Sea	23-24	109-113			13-17		3	12-14	6
<i>Lycenchelys jordani</i>	Calif.-SE Alaska	22-24	100-109	116	93	15-17		1	15-18	6
<i>Lycenchelys longirostris</i>	Bering Sea	21-22	92-93	108-109	93-95	15-16			(9-11)	
			(113-115)							
<i>Lycenchelys microporus</i>	Bering Sea	29	94			18		2	10	6
<i>Lycenchelys pliciferus</i>	Bering Sea	28-29	96-97			15				6
<i>Lycenchelys rassi</i>	Bering Sea	23-25	98-103	119	104	15-16		1-2	8-11	6
<i>Lycenchelys ratmanovi</i>	Bering Sea	22-23	88-89			16-19		1-2	7-9	6
<i>Lycenchelys roseus</i>	Aleutian Is.	28	118-119	130-133	114-115	14-15		1-2	9-10	6
<i>Lycenchelys volki</i>	Bering Sea	30	96			17		2	14	6
<i>Lycodapus derjugini</i>	Bering Sea	14-15	56-59	70-79	57-68	6-7	ab			6
<i>Lycodapus dermatinus</i>	SSC-SE Alaska	13-15	62-68	70-75	62-66	6-7	ab			6
<i>Lycodapus endemoscotus</i>	SSC-Brit. Col.	14-17	72-79	84-91	74-81	6-8	ab			6
<i>Lycodapus fierasfer</i>	SSC-Bering Sea	13-15	69-77	78-85	68-74	6-8	ab			6
<i>Lycodapus leptus</i>	Bering Sea	16-19	78-82	91-94		6-8	ab			6
<i>Lycodapus mandibularis</i>	S. Calif.-Bering Sea	14-17	67-80	76-90	65-79	6-9	ab			6
<i>Lycodapus pachysoma</i>	Oregon-Brit. Col.	14-16	60-63	70-74	58-64	7-8	ab			6
<i>Lycodapus parviceps</i>	Wash.-Bering Sea	18-20	81-85	94-98		8-9	ab			6
<i>Lycodapus poecilus</i>	Bering Sea	15-17	65-72	75-83		5-7	ab			6
<i>Lycodapus psarosomatus</i>	Bering Sea	17-19	77-82	89-93		8	ab			6
<i>Lycodes brevipes</i>	Oregon-Bering Sea	20-22	80-82	85-102	74-89	19-21	3	2-3	11	
<i>Lycodes concolor</i>	Bering Sea	22	92-93	47-118	98-99	21		0-2	12	
<i>Lycodes cortezianus</i>	S. Calif.-SE Alaska	22-24	83-90	112-114	95-97	18-21	3	2	11	6
<i>Lycodes diapterus</i>	S. Calif.-Bering Sea	21-23	100	90-124	94-107	18-25	3	1	13	
<i>Lycodes mucosus</i>	Bering Sea-Arctic		(90-92)	88-93	69-73	17-18				
<i>Lycodes pacifica</i>	SSC-Gulf of Alaska	21-23	79-85	90-107	70-90	16-19	3	0-2	8-12	
<i>Lycodes palearis</i>	Oregon-Chukchi Sea		(96-105)	94-106	83-90	17	3	2-3	10	
<i>Lycodes raridens</i>	Bering Sea-Arctic		(97)	83-93	72-76	18-19		3	10-12	
<i>Lycodes turneri</i>	Bering Sea-Arctic		(90-100)	89-97	69-78	15-18	3			
<i>Lycinema barbatum</i> ^c	SSC-Bering Sea	20-21	86-93	101-105	90-94	15		2	8-9	
<i>Melanostigma pammelas</i>	SSC-Brit. Col.	19-20	69-72	73-88	64-75	6-8	ab		(11-13)	6-7
<i>Nalbantichthys elongatus</i> ^c	Bering Sea	25	119-125	143-152	121-127	6	ab			7
<i>Opaeophacus acrogeineus</i> ^{c,d}	Bering Sea	25-26		141-148	121-124	4-5	ab	3	11	6
			(144-149)							
<i>Pachycara bulbiceps</i>	SSC-Brit. Col.			109	89	16	ab			6
<i>Puzanovia rubra</i> ^c	Bering Sea	22-24	110-125	137-147	115-128	10-12	ab	3-5	9-13	6
<i>Taranetzella lycoderma</i> ^c	Oregon-Bering Sea	19-20	69-78	84-91	71-79	15	3	3	13	
			(90-97)							

^aab = absent.

^bDorsal fin count does not include one spine.

^cTotal principal caudal fin ray counts available for only the following species: *Lycinema barbatum*, 12; *Nalbantichthys elongatus*, 7-10; *Opaeophacus acrogeineus*, 8-9; *Puzanovia rubra*, 9-12; *Taranetzella lycoderma*, 8.

^dBond and Stein 1984

A Unidentified zoarcid**B** *Zoarces viviparous***C** *Bothrocara hollandi***D** *Gymnelus viridis***E** *Macrozoarces americanus*

Figures A–E, Kendall et al. 1983 (B, collected from Gulf of Riga, Baltic Sea; C, Japan Sea specimen; D, Barents Sea specimen; E, collected near New Brunswick).

STICHAEIDAE

Pricklebacks are found mostly in the North Pacific in inshore areas. Adults are long and somewhat eel-like with a long dorsal fin composed of all spines in most species. Seventeen genera and 26 species are found in the study area. This family is separated into eight tribes grouped within four subfamilies. Six tribes within three subfamilies occur in our area.¹ Larvae are abundant in the area but very few larval series have been described. Before fin rays develop, small larvae are elongate and resemble other elongate forms, especially bathymasterids and pholidids (see Table 4). In general, bathymasterids have fewer myomeres and pholidids have a longer gut and more myomeres. Since so few complete larval series are available, general characters based on only a few species are presented for each tribe.

Stichaeinae

Adult characters include: Pelvic fins I,3-4
(*Gymnoclinus* I,2)
Large pectoral fins
Vertebrae 49-76

Stichaeini

Genera found within the study area: *Eumesogrammus* (one species) and *Stichaeus* (one species). Myomere counts are low (<55). *Stichaeus* larvae are identified by the distinctive lateral pigment along the hypaxial myomeres.


Chirolophini

Genera found within the study area: *Bryzoichthys* (two species), *Chirolophis* (four species), and *Gymnoclinus* (one species). Myomere counts are >55. Larvae are generally more pigmented than Stichaeini. Melanophores occur along the dorsal and ventral body midline, over the notochord internally, and, in some taxa, around the urostyle.


¹Makushok (1958) places the genera *Eulophias* and *Azygopterus* in separate tribes under Xiphisterinae. Anderson (1984b) places them together in the Eulophiini. Other nomenclatural changes not affecting taxa in the study area are presented by Yatsu (1986).

STICHAEINI
Stichaeus punctatus

A




B




CHIROLOPHINI
Bryozoichthys–Chirolophis

C



D



Figures A–B (B, ventral view), Fahay 1983 (after Faber 1976, North Atlantic specimen); C–D, NWAFC originals (B. Vinter).

STICHAEIDAE

Lumpeninae

Adult characters include: Pelvic fins I,3 or absent
Large pectoral fins
Vertebrae 60-81


Lumpenini

Genera found within the study area: *Acantholumpenus* (one species), *Anisarchus* (one species), *Lumpenella* (one species), *Lumpenus* (four species), and *Poroclinus* (one species). Lumpenini larvae generally lack pigment along the dorsal midline. Diagnostic pigment usually occurs over the dorsal surface of the gut and anus, along the ventral midline, and in the hypural area.



Lumpenella larvae generally have >70 myomeres and pigment around the urostyle, and, with development, can be identified by their distinctive snout and the presence of up to five anal spines. *Lumpenus* spp. larvae can be identified by meristics, number of postanal ventral melanophores, and number of melanophores on the anus. Hypural pigment is usually restricted to several spots along the posterior edge of the hypural area. *Poroclinus* larvae have distinctive pigment occurring above and below the notochord in the caudal peduncle area. They can also be distinguished from other Lumpenini by the presence of three anal spines.

LUMPENINI



Lumpenella longirostris




Lumpenus maculatus



Lumpenus sagitta



Poroclinus rothrocki



Figures A, C, F, Garrison, unpubl.; B, Fahay 1983 (after Faber 1976, Atlantic specimen); D-E, NWAFC originals (B. Vinter).

STICHAEIDAE

Opisthocentrini

Genera found within the study area: *Allolumpenus* (one species), *Opisthocentrus* (one species), and *Plectobranchnus* (one species). Larvae are known only for *Opisthocentrus* and *Plectobranchnus*. *Plectobranchnus* larvae have a distinctive pigment pattern quite dissimilar to *Opisthocentrus* (see species pages). *Opisthocentrus* larvae appear to resemble Aletrini larvae (e.g., similar pigment pattern and lack of pelvic fins).

Xiphisterinae

Adult characters include: Pelvic fins absent
Pectoral fins small
Vertebrae 57-84


Aletrini

Genera found within the study area: *Aletridium* (one species) and *Anoplarchus* (two species). Only larvae of *Anoplarchus purpurescens* are known from our area. They have a row of postanal ventral midline melanophores and a few spots along the hypural margin. No single unique character distinguishes them from Xiphisterini larvae but usually a combination of characters allows identification. These characters include fewer myomeres (58-68 vs. generally >70), smaller relative eye diameter, fewer melanophores dorsally on gut, and presence of hypural spots throughout the larval period (see species pages).


Xiphisterini

Genera found within the study area: *Cebidichthys* (one species), *Phytichthys* (one species), and *Xiphister* (two species). Although series of *Phytichthys* and *Xiphister* have been described, larvae are difficult to distinguish from one another and from *Anoplarchus*. Xiphisterini larvae usually have a row of postanal ventral melanophores and a row of internal pigment above the notochord.


OPISTHOCENTRINI
Opisthocentrus ocellatus



Plectobranthus evides



ALECTRINI
Anoplarchus purpurescens



XIPHISTERINI
Xiphister atropurpureus




Figure A, Shiogaki 1982 (reared from Japanese specimens); B–D, NWAFC originals (B. Vinter).

Table 44
Meristic characters of family Stichaeidae.

Taxon	Distribution	Vertebrae		Dorsal	Fins			Gill rakers		Branchiostegals
		Precaudal	Caudal (Total)		Anal	Pectoral	Pelvic ^a	Upper	Lower	
STICHAEINAE										
Stichaeini										
<i>Eumesogrammus praecisus</i>	Bering Sea-Arctic	15-16	34-36 (50-52)	XLVII-XLIX	II,31-33	18	I,3			6
<i>Stichaeus punctatus</i>	Brit. Col.-Arctic	14-16	36-40 (51-55)	XLVI-XLIX	I-II,32-35	15-16	I,4			6
Chirolophini										
<i>Bryzoichthys lysimus</i>	Aleutian Is.-Bering Sea	16-17	53-54	LXIII-LXVI	I,49-50	14-15	I,3	4-5	9-11	6
<i>Bryzoichthys marjorius</i>	Brit. Col.-Aleutian Is.	15-17	56-59 (72-75)	LXII-LXXI	I,51-59	14-16	I,3	5-6	9-10	6
<i>Chirolophus decoratus</i>	N. Calif.-Bering Sea			LXI-LXIII	I,44-51	14-15	I,4			6
<i>Chirolophus nugator</i>	S. Calif.-Aleutian Is.			LIII-LV	I,37-42	13-14	I,4			6
<i>Chirolophus snyderi</i>	Bering Sea	16-17	46-49 (63-65)	LVIII-LXI	I,43-45	15	I,4			6
<i>Chirolophus tarsodes</i>	Brit. Col.-Bering Sea			LVIII-LX	I,43-45	14-15	I,3-4			6
<i>Gymnoclinus cristulatus</i>	Bering Sea			LXI	40-43	14	I,1-2			6
LUMPENINAE										
Lumpenini										
<i>Acantholumpenus mackayi</i>	Bering Sea	27-29	49-52 (76-80)	LXVIII-LXXV	II,41-47	14-15	I,3			6
<i>Anisarchus medius</i>	SE Alaska-Arctic			LXI-LXXI	II-V,36-42	13-14	I,2-3			6
<i>Lumpenella longirostris</i> ^b	Brit. Col.-Gulf of Alaska	24-25	47-49	LXI-LXXI	I,40-43	13-16	I,3			6
<i>Lumpenus fabricii</i> ^b	SE Alaska-Arctic	26-28	44-48 (70-75)	LXI-LXV	I,40-43	13-16	I,3			6
<i>Lumpenus maculatus</i> ^b	Wash.-Arctic	26-30	39-43 (66-72)	LVII-LXIV	I-II,34-40	14-16	I,3-4			6
<i>Lumpenus medius</i>	SE Alaska-Arctic	23-25	43-50 (65-70)	LVIII-LXIII	I,37-42		I,3			6
<i>Lumpenus sagitta</i> ^b	N. Calif.-Bering Sea	26-28	46-54	LXIV-LXXII	I,45-50	15-17	I,3-4			6
<i>Poroclinus rothrocki</i>	S. Calif.-Bering Sea		(65)	LVII-LXXVII	III,40-44	13-15	I,3 ^c			6
Opisthocentrini										
<i>Allolumpenus hypochromus</i>	S. Calif.-Brit. Col.			XLIV-XLIX	I,31	12	I,3			6
<i>Opisthocentrus ocellatus</i>	Bering Sea	22-23	40-44 (63-67)	LVIII-LXII	II,37-39	20-21	ab			5
<i>Plectobranchnus evides</i> ^b	S. Calif.-Brit. Col.		(60)	LIV-LVII	II-III,34-36	15	I,3			5-6
XIPHISTERINAE										
Alectrini										
<i>Alectridium aurantiacum</i>	Bering Sea	19-21	46-48 (65-68)	LIX-LXIII	I,41-44	11	ab			5
<i>Anoplarchus insignis</i>	N. Calif.-Aleutian Is.	17-19	44-49	LVII-LXIV	40-46	9-10	ab			5
<i>Anoplarchus purpureus</i>	S. Calif.-Bering Sea	17-19	40-46 (58-64)	LIV-LX	I,36-41	9-10	ab	3-5	5-10	5
Xiphisterini										
<i>Cebidichthys violaceus</i>	SSC-Oregon	23-25	40-47 (65-71)	XXII-XXV, 40-43	I-II,39-42	10-11	ab	3-4	6-10	6
<i>Phytichthys chirus</i> ^b	S. Calif.-Bering Sea	24-25	50-52 (75-76)	LXIX-LXXXVIII	II-III,40-50	15	ab			6
<i>Xiphister atropurpureus</i> ^b	SSC-Bering Sea	22-24	51-56 (75-80)	LXV-LXXII	I,49-55	11-12	ab	2-3	6-10	6
<i>Xiphister mucosus</i> ^b	Cent. Calif.-SE Alaska	29-31	44-53 (73-83)	LXXI-LXXXVIII	I,46-50	12	ab			6

^aab = absent.

^bPrincipal caudal fin ray counts available for only the following species: *Lumpenella longirostris*, 6+6-7; *Lumpenus fabricii*, 6+7; *Lumpenus maculatus*, 7+6; *Lumpenus sagitta*, 6+7; *Plectobranchnus evides*, 6+5; *Phytichthys chirus*, 6+7; *Xiphister atropurpureus*, 6+7; *Xiphister mucosus*, 6+7.

^cRudimentary.

STICHAEIDAE
(Opisthocentrini)

***Opisthocentrus ocellatus* (Tilesius 1811)**

MERISTICS

Vertebrae	Total: 63-X-67 Precaudal: 22-X-23 Caudal: 40-X-44
Branchiostegal rays	5-5-5
Caudal fin	
Pelvic fin	Absent
Dorsal fin	S: 58-X-62
Pectoral fin	R: 20-X-21
Anal fin	S: 2-2-2 R: 37-X-39
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Dec-Jan (Mutsu Bay, Japan) ^a Area: Narrow cavities under stones on muddy sand bottoms ^a Mode: Eggs spawned in masses and guarded by female ^a Migration:
Fecundity	Range/function: 700-3300 ^a
Age at first maturity	1 yr ^a
Longevity	2-3 yr ^a

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.89-2.01 mm
No. of oil globules	One large, many small (yellow)
Oil globule diameter	
Yolk	Colorless
Envelope	
Hatch size	9-10 mm SL
Incubation time/temp.	48 d/5-10°C
Pigment	

Diagnostic characters

LARVAE


Preanal length	<50% SL
Length at flexion	
Length at transformation	33-40 mm SL ^b
Sequence of fin development	
Pigment	
	• Dorsally and ventrally on gut
	• Postanal ventral melanophores
	• Hypural spot

Diagnostic characters

^aShiogaki 1982

^bCollected with a small trawl net.

Ref: Shiogaki 1982.



Figures A–D, Shiogaki 1982 (A–C, reared from specimens collected from Mutsu Bay, Japan; D, collected from Mutsu Bay, Japan-probably newly settled).

STICHAEIDAE
(Opisthocentrini)

Plectobranthus evides Gilbert 1890

MERISTICS

Vertebrae	Total: 60-60-60 Precaudal: X-X-X Caudal: X-X-X
Branchiostegal rays	5-X-6
Caudal fin	X, 6+5, X
Pelvic fin	Thoracic S: 1-1-1 R: 3-3-3
Dorsal fin	S: 54-X-57
Pectoral fin	R: 15-15-15
Anal fin	S: 2-X-3 R: 34-X-36
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	S. California, 32-34°N, to Brit. Col., 48°30'-55°N
Ecology	Epi- and mesobenthal, 84-274 m
ELH pattern	Probably oviparous, eggs probably demersal, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters

LARVAE^a


Preanal length ~50% SL
Length at flexion Between 10 and 16 mm SL
Length at transformation
Sequence of fin
development
Pigment

- Blotch medially over gut near cleithrum
- Blotches on ventral body at midgut, hindgut over anus, and at myomeres 30 and 45
- Pigment becomes more internal in larger specimens
- Above and below notochord, increasing with development anteriorly in area where hypurals form

Diagnostic characters

- Pigment pattern: Five blotches along body
- Meristics: anal spines 2-3
pelvic fin 1,3

^aLarvae obtained from Bruce Mundy, formerly of Oregon State University (present address, NMFS Southwest Fish. Cent., Honolulu Lab., 2570 Dole St., Honolulu, HI 96822-2396); originally identified by S.L. Richardson (deceased).



Figures A-C, NWAFC originals (B. Vinter).

MERISTICS

Vertebrae	Total: 58-60-64 Precaudal: 17-18-19 Caudal: 40-43-46
Branchiostegal rays	5-5-5
Caudal fin	
Pelvic fin	Absent
Dorsal fin	S: 54-X-60
Pectoral fin	R: 9-X-10
Anal fin	S: 1-1-1 R: 36-39-41
Gill rakers	U: 3-X-5 L: 5-X-10

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal, intertidal to 30 m
ELH pattern	Oviparous; demersal, adhesive, attached eggs; pelagic larvae
Spawning	Season: Jan ^a -Mar ^b Area: Demersal, under rocks or on shells ^a Mode: Pairs; eggs laid in masses; ^c females guard nests ^d Migration:
Fecundity	Range/function: 2000-3000 ^d
Age at first maturity	2-3 yr ^d
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.27-1.45 mm
No. of oil globules	1-3, with development 1 large and 1 small
Oil globule diameter	
Yolk	
Envelope	White
Hatch size	~7.5 mm TL
Incubation time/temp.	
Pigment	

Diagnostic characters

- Eggs laid in pedestal-shaped mass (flattened with lateral constriction)^c

LARVAE

Preanal length	40-45% SL
Length at flexion	~10 mm SL
Length at transformation	~12 mm SL
Sequence of fin development	
Pigment	

- Postanal ventral midline: A melanophore on nearly every myoseptum, at base of each anal fin ray
- A few melanophores at caudal fin base along hypural margin
- Dorsal and ventral gut pigment: Dorsally about five spots on posterior half, ventrally a continuous line on anterior 2/3 of gut
- Heavy uniform superficial body pigment on larvae >12 mm SL (early juveniles)

Diagnostic characters

- Distinguished from *Xiphister* spp. (p. 516, 518) and *Phytichthys chirus* (p. 514) by
- Smaller relative eye diameter
 - Fewer dorsal gut melanophores (<8)
 - Hypural spots throughout larval period


^aMarliave 1975a

^bSchultz and DeLacy 1932

^cJ. Marliave, Vancouver Public Aquarium, P.O. Box 3232, Vancouver, B.C., Canada V6B 3X8, pers. commun., 16 Oct. 1986.


^dPeppar 1965

A




6.1 mm SL

B




9.0 mm SL

C



12.0 mm SL


D



12.0 mm SL

Facial pattern develops in larvae approaching settlement

E



12.0 mm SL

Figures A–E, NWAFC originals (B. Vinter, reared).

STICHAEIDAE
(Xiphisterini)

Phytichthys chirus (Jordan and Gilbert 1880)

MERISTICS

Vertebrae	Total: 75-X-76 Precaudal: 24-X-25 Caudal: 50-X-52
Branchiostegal rays	6-6-6
Caudal fin	X, 6+7, X
Pelvic fin	Absent
Dorsal fin	S: 69-X-78
Pectoral fin	R: 15-15-15
Anal fin	S: 2-X-3 R: 40-X-50
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Intertidal, nearshore shelf demersal
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Mid-winter (Brit. Col.) ^a Area: Exposed intertidal, under rocks ^a Mode: Pairs; single parent guards ^a Migration:
Fecundity	Range/function: 1100 (one mass) ^a
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION^a

EGGS

Diameter	2.25 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	White
Hatch size	11.2 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

- Eggs laid in conical-shaped mass

LARVAE

Preanal length	45% SL
Length at flexion	13 mm SL
Length at transformation	18-21 mm SL
Sequence of fin development	Caudal, dorsal and anal, pectorals
Pigment	

- Postanal ventral midline with about 30 spots
- Dorsally about eight spots on gut; ventral pigment present (not shown on figure)
- Row of internal pigment over notochord

Diagnostic characters

Distinguished from *Xiphister* spp. (p. 516, 518) by

- Relatively larger at comparable stages of development (e.g., hatching occurs >10.0 mm SL)
- Larger relative eye diameter


- Dorsal body midline unpigmented in preflexion and flexion larvae

Distinguished from *Anoplarchus purpurescens* (p. 512) by

- Lack of hypural spots in preflexion and flexion larvae

^aJ. Marliave, Vancouver Public Aquarium, P.O. Box 3232, Vancouver, B.C., Canada V6B 3X8, pers. commun., 16 Oct. 1986. Drawings by C. Hui.

Ref: Marliave, unpubl.



Figures A–C, Marliave, unpubl.

STICHAEIDAE
(Xiphisterini)

***Xiphister atropurpureus* (Kittlitz 1858)**

MERISTICS

Vertebrae	Total: 75-75-80
	Precaudal: 22-23-24
	Caudal: 51-52-56
Branchiostegal rays	6-6-6
Caudal fin	X, 6+7, X
Pelvic fin	Absent
Dorsal fin	S: 65-X-72
Pectoral fin	R: 11-X-12
Anal fin	S: 1-1-1 R: 49-X-55
Gill rakers	U: 2-X-3 L: 6-X-10

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal, intertidal to 8 m
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Late winter-early spring (British Columbia) ^a Area: Intertidal, under rocks ^a Progression: Early in protected waters, later on exposed shores ^a Mode: Pairs; males guard nests ^b Migration:
Fecundity	Range/function: 900-1700 ^b
Age at first maturity	2-3 yr ^c
Longevity	

EARLY LIFE HISTORY DESCRIPTION^d

EGGS

Diameter	2.25 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	White
Hatch size	8.5 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	43-45% SL
Length at flexion	11 mm SL
Length at transformation	18 mm SL
Sequence of fin development	Caudal, anal and dorsal, pectorals
Pigment	

- Postanal ventral midline with about 36 spots (range 28-46)
- Dorsally about eight spots on gut; ventral pigment present (not shown on figure)
- Row of internal pigment over notochord

Diagnostic characters

- See *Anoplarchus purpureus* (p. 512) and *Phytichthys chirus* (p. 514)
- Distinguished from *X. mucosus* by
- Relatively smaller at comparable stages of development
- Internal pigment over notochord generally more prominent
- Relatively short snout-anus length (preanal myomeres 23-25 vs. 28-30 in *X. mucosus*)


^aMarliave 1975b

^bMarliave and DeMartini 1977

^cWingert 1974

^dJ. Marliave, Vancouver Public Aquarium, P.O. Box 3232, Vancouver, B.C., Canada V6B 3X8, pers. commun., 16 Oct. 1986. Drawings by C. Hui.

Ref: Marliave, unpubl.



Figures A–C, Marliave, unpubl.

STICHAEIDAE
(Xiphisterini)

Xiphister mucosus (Girard 1858)

MERISTICS

Vertebrae	Total: 73-81-83
	Precaudal: 29-30-31
	Caudal: 44-50-53
Branchiostegal rays	6-6-6
Caudal fin	X, 6+7, X
Pelvic fin	Absent
Dorsal fin	S: 71-X-78
Pectoral fin	R: 12-12-12
Anal fin	S: 1-1-1 R: 46-X-50
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Cent. California, 34-38°N, to SE Alaska, 55-59°N
Ecology	Nearshore shelf demersal, intertidal to 18 m
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Late winter-spring (British Columbia) ^a Area: Exposed intertidal, under rocks ^a Mode: Pairs; males guard nests ^b Migration:
Fecundity	Range/function: 5500-9500 ^a
Age at first maturity	5 yr ^c
Longevity	

EARLY LIFE HISTORY DESCRIPTION^d

EGGS

Diameter	2.5 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	White
Hatch size	9.5 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	47-49% SL
Length at flexion	12 mm SL
Length at transformation	18 mm SL
Sequence of fin development	Caudal, dorsal and anal, pectorals
Pigment	

- Postanal ventral midline about 26 spots (range 18-35)
- Dorsally about eight spots on gut; ventral pigment present (not shown on figure)
- Row of internal pigment over notochord

Diagnostic characters




- See *Anoplarchus purpureus* (p. 512) and *Phytichthys chirus* (p. 514)
- Distinguished from *X. atropurpureus* by
- Generally larger at comparable stages of development
 - Internal pigment over notochord generally lighter

^aMarliave 1975a

^bMarliave and DeMartini 1977

^cWingert 1974

^dJ. Marliave, Vancouver Public Aquarium, P.O. Box 3232, Vancouver, B.C., Canada V6B 3X8, pers. commun., 16 Oct. 1986. Drawings by C. Hui.



Figures A–C, Marliave, unpubl.

MERISTICS *Delolepis gigantea* (Kittlitz 1858)

Vertebrae	Total: 81-X-85 Precaudal: X-X-X Caudal: 49-X-51
Branchiostegal rays	6-6-6
Caudal fin	X, 7+8, X
Pelvic fin	Absent
Dorsal fin	S: 73-X-77
Pectoral fin	R: 13-13-13
Anal fin	S: 2-2-2 R: 43-X-49
Gill rakers	U: X-X-X L: X-X-X

MERISTICS *Lyconectes aleutensis* Gilbert 1896

Vertebrae	Total: 71-X-77 Precaudal: X-X-X Caudal: 47-X-51
Branchiostegal rays	6-6-6
Caudal fin	X, 7+8, X
Pelvic fin	Absent
Dorsal fin	S: 60-X-69
Pectoral fin	R: 12-X-13
Anal fin	S: 2-X-3 R: 45-X-49
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	N. California, 38-42°N, to Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal (<i>D. gigantea</i> , 6-128 m); epi- and mesobenthic (<i>L. aleutensis</i> , 46-350 m)
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Spring-summer (<i>L. aleutensis</i>) ^b Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	4.41-4.85 mm (<i>D. gigantea</i>); 1.8 mm (<i>L. aleutensis</i>)
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	Smooth (<i>D. gigantea</i>)
Hatch size	Larvae with yolk at 16-17 mm SL (<i>D. gigantea</i>)
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**


Preanal length	~50% SL (<i>D. gigantea</i>); <50% SL (<i>L. aleutensis</i>)
Length at flexion	
Length at transformation	
Sequence of fin development	Caudal, pectorals, dorsal and anal; fin development may begin prior to hatching in <i>D. gigantea</i>
Pigment	

Diagnostic characters (see Table 3)


- Number of myomeres: 81-85 in *D. gigantea*, 71-77 in *L. aleutensis*
- Pigment on isthmus and over gut (absent in *D. gigantea*)
- Preanal length: ~50% SL in *D. gigantea*, <50% SL in *L. aleutensis*
- Size at stage of development: *D. gigantea* larvae hatch at a larger size and more advanced stage of development

^aNawojchik (1986) includes *Lyconectes* and *Delolepis* in the genus *Cryptacanthodes*.^bHart 1973

A *Delolepis gigantea*



B *Lyconectes aleutensis*



Figures A–B, NWAFC originals (B. Vinter).

PHOLIDIDAE

Although some species of gunnels are found in the North Atlantic, most members of this family are located in the eastern Pacific. Pholidids are eel-like with long compressed bodies, a dorsal fin extending from head to caudal fin and made up entirely of spines, and small pelvic fins (I,1) when present. There are nine species in three genera within the study area.¹ Adults are demersal, occurring from as deep as 75 m to tidepools (Hart 1973). Pholidids are noted for paired spawning and subsequent guarding of eggs which may be 1.4-3.0 mm and have one oil globule (A.C. Matarese, unpubl.). One or both partners may guard the eggs during incubation by coiling around them (Breder and Rosen 1966). Larvae are pelagic and may settle after 50 days (Garrison and Miller 1982).

Presently, larvae of *Pholis* spp. cannot be identified to species in our study area. The following characters may be helpful in separating *Pholis* spp. larvae from those of *Apodichthys flavidus*.

- Morphology** Head and eye generally smaller in *Pholis* at comparable stages of development
- Meristics** Presence of pelvic fins and two anal spines in *Pholis*
- Pigment** Internal pigment above notochord is less pronounced and disappears sometime before larvae undergo flexion; presence of a continuous series of melanophores along ventral surface of gut; pigment spots along dorsal surface of gut are smaller and more numerous (postflexion)

Meristic characters may be useful in separating larger *Pholis* postflexion larvae and juveniles. Off Washington, Oregon, and California, myomere counts enable separation of *P. clemensi* (94-98) and *P. schultzi* (89-93) from other species (usually <90).

Larvae of *Xererpes fucorum* are inadequately known. *X. fucorum* can be separated from *Pholis* spp. by their lack of pelvic fins and from *Apodichthys flavidus* by the presence of one or two anal spines and a lower dorsal spine count (83-87 vs. 90-94).


¹Yatsu (1985) proposes alternative generic placements for several species; according to his arrangement, nine species within four genera occur in the area.

Table 45
Meristic characters of family Pholididae.*



Taxon	Distribution	Vertebrae		Fins				Gill rakers		Branchiostegals
		Precaudal	Caudal	Dorsal	Anal	Pectoral	Pelvic	Upper	Lower	
<i>Apodichthys flavidus</i>	S. Calif.-Gulf of Alaska	50-51	48	XC-XCIV	I,36-42	14		3	10-13	5
<i>Pholis clemensi</i>	N. Calif.-SE Alaska	37-39	57-59	LXXXVII-XCI	II,48-53	11-14	I,1		11	5
<i>Pholis dolichogaster</i>	Aleutian Is.-Bering Sea	41-45	56-57	XCIII-XCVI	II,48-51	13-15	I,1			5
<i>Pholis fasciata</i>	Bering Sea	40-44	50	LXXXIII-LXXXVIII	II,41-44	11-13	I,1			5
<i>Pholis gilli</i>	Bering Sea			LXXXIV	II,43		I,1			5
<i>Pholis laeta</i>	N. Calif.-Bering Sea	40-42	43-44	LXXIV-LXXXI	II,32-38	11-12	I,1			5
		(81-89)								
<i>Pholis ornata</i>	Cent. Calif.-Brit. Col.		(80-87)	LXXIV-LXXX	II,34-38	11-12	I,1			5
<i>Pholis schultzi</i>	Cent. Calif.-Brit. Col.		(89-93)	LXXX-LXXXIX	II,40-44	10-12	I,1-2	1-2	7-10	
<i>Xererpes fucorum</i>	SSC-Brit. Col.	52	40	LXXXII-LXXXVII	I,29-38	12		1-2	6-9	5

*Yatsu (1985) places *Pholis clemensi*, *P. laeta*, and *P. schultzi* in the genus *Allopholis*. *P. dolichogaster* is placed in the genus *Rhodymenichthys*. *Xererpes fucorum* is placed in the genus *Apodichthys*.

Apodichthys flavidus



Pholis sp.



Figures A–C, NWAFC originals (B. Vinter).

MERISTICS

Vertebrae	Total: 88-X-89
	Precaudal: 29-X-31
	Caudal: 57-X-59
Branchiostegal rays	6-X-7
Caudal fin	
Pelvic fin	Absent
Dorsal fin	S: 81-X-88
Pectoral fin	R: 20-X-22
Anal fin	S: X-X-X R: 50-X-55
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Area: Mode: Pairs; eggs guarded by one or both parents ^a
Fecundity	Migration:
Age at first maturity	Range/function:
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	4-8 mm
No. of oil globules	One
Oil globule diameter	
Yolk	
Envelope	
Hatch size	~17-18 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	50% SL
Length at flexion	<20 mm SL
Length at transformation	~40 mm SL
Sequence of fin development	

Pigment (see Table 3)

- Heavily pigmented over body except ventral surface of gut, pectoral fin base, and posteriormost opercular area

Diagnostic characters

Distinguished from *Anarrhichthys ocellatus* by

- Morphology: Body not elongate
- Lower vertebral count (88-89 myomeres)
- Eye diameter (large)

^aBreder and Rosen 1966

Ref: Andriashev 1954, Barsukov 1959.




Figure A, NWAFC original (B. Vinter); B-C, Kobayashi 1961a.

MERISTICS

Vertebrae	Total: 221-247-251 Precaudal: 36-37-39 Caudal: 183-204-214
Branchiostegal rays	6-X-7
Caudal fin	
Pelvic fin	Absent
Dorsal fin	S: 218-X-250
Pectoral fin	R: 19-X-20
Anal fin	S: 0-0-1 R: 180-X-233
Gill rakers	U: 3-X-5 L: 11-X-15

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthal, 0-226 m
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Oct-Feb ^a Area: In caves or rocky crevices ^a Mode: Pairs; eggs guarded by both parents ^a Migration: Mated pairs sedentary, den site permanent ^a Range/function: ~10,000 ^a
Fecundity	
Age at first maturity	
Longevity	4 yr ^a

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	5.2-5.4 mm
No. of oil globules	
Oil globule diameter	
Yolk	Reddish
Envelope	Smooth; white, becoming brownish
Hatch size	33.5 mm SL
Incubation time/temp.	3-4.5 mo
Pigment	

Diagnostic characters

- Embryo coils around 3-3.5 times within envelope

LARVAE

Preanal length	25% SL at hatching, decreasing with develop- ment
Length at flexion	Tail fully formed at hatching
Length at transformation	Hatch as juveniles
Sequence of fin development	Fins fully formed at hatching with yolk present
Pigment	
	• Dorsal head
	• Upper body over gut
	• Small concentrated melanophores along dorsal and ventral body midlines that extend onto the body laterally and into the finfolds with development

Diagnostic characters

- Elongate body shape
- High number of myomeres (221-251)

^aMarliave 1987

Ref: Marliave 1975a, 1987.




Figure A, Marliave 1975a.

MERISTICS^a

Vertebrae	Total: 227-X-240	
	Precaudal: 53-X-59	
	Caudal: 174-X-181	
Branchiostegal rays	5-5-5	
Caudal fin		
Pelvic fin	Absent	
Dorsal fin	S: 83-X-90	R: 137-X-145
Pectoral fin	R: 13-13-13	
Anal fin	R: 180-X-196	
Gill rakers	U: X-X-X	L: X-X-X

LIFE HISTORY

Range	Oregon, 42-46°N, to Bering Sea, 54-66°N
Ecology	Nearshore shelf demersal, 0-80 m
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Spring ^b Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter
No. of oil globules
Oil globule diameter
Yolk
Envelope
Hatch size
Incubation time/temp.
Pigment

Diagnostic characters**LARVAE**

Preanal length Gut length 35-40% SL
(decreases with
development)

Length at flexion
Length at transformation ~114 mm SL
Sequence of fin Dorsal and anal first at
development 40 mm SL
Pigment (larvae 20.3-36.0 mm SL)

- Head
 - Lower jaw
 - Isthmus
 - Internally at base of hindbrain
- Gut: Dorsal and ventral surface
- Body: Concentrated dorsally and ventrally
- Caudal: "Fleshy caudal extension" is distinctly pigmented; pigment is scattered evenly dorsally and ventrally on body and into finfolds

Diagnostic characters

- Morphology: Elongate form, gut length (40% SL)
- Number of myomeres (>225)
- Pigment pattern: Concentrated ventrolateral spots and pigment on fleshy caudal extension

^aKobayashi (1961b) reports slightly different meristics in specimens from Japanese waters:

Dorsal fin spines	79-83
Dorsal fin rays	141-157
Anal fin rays	166-193
Pectoral fin rays	11-13
Branchiostegal rays	6

Richardson and Dehart (1975) note that in four larval specimens collected off Oregon the total vertebral counts of 227 were considerably lower than counts in Bering Sea specimens (236-240).

^bLarvae 20.3-36.0 mm SL collected March-May 18 km from coast of Oregon.

Ref: Richardson and Dehart 1975.





Figure A, Richardson and Dehart 1975.



Figures A–D, Haryu and Nishiyama 1981.

MERISTICS

Vertebrae	Total: 44-X-47 Precaudal: 12-12-14 ^a Caudal: 32-34-40 ^a
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 14-X-16 R: 18-X-20
Pectoral fin	R: 21-X-22
Anal fin	S: 1-1-1 R: 28-X-29
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	N. California, 38-42°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthic, 20-375 m
ELH pattern	Oviparous; demersal, adhesive, attached eggs; pelagic larvae
Spawning	Season: Winter-spring (British Columbia) ^a Area: Rocky intertidal ^a Mode: Migration: Along shore to rocky areas ^a
Fecundity	Range/function: ~1000 ^a
Age at first maturity	
Longevity	

^aMarliave 1981c^bM. Okiyama, Univ. Tokyo, Ocean Res. Inst., 1-15-1 Minamidai, Nakano-Ku, Tokyo 164, Japan, pers. commun., 8 Nov. 1985.

Ref: Marliave 1981c.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	3.5 mm
No. of oil globules	One
Oil globule diameter	
Yolk	
Envelope	
Hatch size	13 mm SL (preserved)
Incubation time/temp.	~1 yr
Pigment	

Diagnostic characters

- Precocious caudal development

LARVAE


Preanal length	35-40% SL, increases with development
Length at flexion	Occurs prior to hatch
Length at transformation	
Sequence of fin development	Caudal, dorsal and anal, pectorals and pelvics
Pigment	<ul style="list-style-type: none"> • On about every other myomere along ventral midline • Distinct spots on hypural margin, increase in number with development to become a line • Large melanophores cover entire surface of gut • Several large melanophores on crown • Anterior dorsal midline starts on nape, increases posteriorly

Diagnostic characters

- Early caudal development
- Hypural margin pigment: First forms line, then an
anteriorly open bracket; similar smaller bracket
develops on the caudal peduncle
- Slightly separate dorsal fins
- Anal fin longer than second dorsal fin

To distinguish from *Arctoscopus japonicus* (present in the
Bering Sea)^b

- *A. japonicus* larvae are less advanced at similar
stages of development than those of *T. trichodon*,
e.g., newly hatched *A. japonicus* larvae possess no
developing fin rays and an unflexed notochord
- Dorsal margin pigment not present until juvenile
stage in *A. japonicus*
- *A. japonicus*: Dorsal X-XI,13; anal 30-31



Figures A–D, Marliave 1981b.

Clinids are found in both Atlantic and Pacific tropical and temperate waters. They are small (many <7.5 cm) and elongate, and somewhat deep-bodied. The dorsal fin extends from behind the head almost to the caudal fin. In the northeastern Pacific the family is represented by only three species in two genera. Adults inhabit nearshore rocky areas from intertidal zones to depths of 50 m. Spawning behavior consists of nest building by a male in a rocky crevice, on seaweed, or in the lumen of a living sponge (Breder and Rosen 1966). Eggs, which are laid by one or more females, form clumps. The adhesive eggs (0.85-1.7 mm) have one or more oil globules which may be uncolored, pale yellow, or orange; sticky filaments are present on eggs of some species. Males of some tribes possess a modified anal fin or intromittent organ (Ophiclini, Clinini), and reproduction within those tribes is either viviparous or ovoviviparous. Larvae are pelagic for a brief period before settling to the bottom (Breder and Rosen 1966). The early-life-history stages of *Heterostichus rostratus* are presented here. Other clinids in the study area include two species of the genus *Gibbonsia*. In comparison to *H. rostratus*, members of *Gibbonsia* have fewer myomeres (48-54), anal fin rays (23-29), pectoral fin rays (11-13), and upper gill rakers (3-4). They undergo flexion and transformation earlier than *H. rostratus* (5.0-8.1 mm SL and ~19 mm SL, respectively).¹ Pigment is generally limited to the ventral midline and dorsally over the swimbladder.

¹W. Watson, Marine Ecological Consultants, 531 Encinitas Blvd., Suite 110, Encinitas, CA 92024, pers. commun., 1983.

Table 46
Meristic characters of family Clinidae.

Taxon	Distribution	Vertebrae		Fins				Gill rakers		
		Precaudal	Caudal	Dorsal	Anal	Pectoral	Pelvic	Upper	Lower	Branchiostegals
<i>Gibbonsia metzi</i>	SSC-Brit. Col.	18-20	32-34	XXXIV-XXXVII, 7-10	II,24-29	11-13	I,3	3-4	7-8	5-6
<i>Gibbonsia montereyensis</i>	SSC-Brit. Col.	16-17	32-35	XXXIV-XXXVI, 5-8	II,23-28	11-13	I,3	3-4	7-12	6
<i>Heterostichus rostratus</i>	SSC-Brit. Col.	21-22	34-36	XXXV-XXXVIII, 11-13	II,31-35	12-14	I,3	5-7	12-13	6

MERISTICS

Vertebrae	Total: 55-57-58
	Precaudal: 21-22-22
	Caudal: 34-35-36
Branchiostegal rays	6-6-6
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 3-3-3
Dorsal fin	S: 35-37-38 R: 11-X-13
Pectoral fin	R: 12-13-14
Anal fin	S: 2-2-2 R: 31-X-35
Gill rakers	U: 5-X-7 L: 12-X-13

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Nearshore shelf demersal, intertidal to 40 m
ELH pattern	Oviparous; adhesive, attached eggs; pelagic ^a larvae
Spawning	Season: Mar; ^b Feb-Apr; ^c spring ^a Area: Egg mass found on floating kelp filaments ^a Mode: Pairs; males guard nests ^a Migration:
Fecundity	Range/function: 400-1200 (may spawn several times/yr) ^a
Age at first maturity	1-1.5 yr ^a (females 14 cm TL)
Longevity	5 yr (females) ^a 3 yr (males) ^a

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.35 mm; ^d 1.4 mm ^a
No. of oil globules	One
Oil globule diameter	
Yolk	Red or brown
Envelope	16 filaments in a cluster
Hatch size	5.5-6.2 mm TL
Incubation time/temp.	12-17 d/18°C
Pigment	
	• Yolk
	• Ventral midline melanophores on embryo

Diagnostic characters

- Filaments

LARVAE

Preanal length	50% SL
Length at flexion	~7-9 mm SL
Length at transformation	By 25 mm SL, 30-50 mm TL
Sequence of fin development	Caudal, dorsal and anal, pectorals and pelvics
Pigment	

- Postanal ventral midline series: Denser posteriorly, with development becoming about one melanophore at base of each anal fin ray
- Anteriorly and dorsally on gut

Diagnostic characters

- Pigmented swimbladder (not shown on figures)
- Ventral midline melanophores
- More myomeres (56-60) than *Gibbonsia* spp. (47-53)
- Dorsal and anal fin development begins posteriorly

^aStepien 1986^bWang 1981^cW. Watson, Marine Ecological Consultants, 531 Encinitas Blvd., Suite 110, Encinitas, CA 92024, pers. commun., 3 Nov. 1986.^dBarnhart 1932




Figure A, Barnhart 1932; B, D, Stepien 1986 (reared); C, E, Matarese et al. 1984a.

MERISTICS

Vertebrae	Total: 66-X-68 Precaudal: 23-23-23 Caudal: 45-45-45
Branchiostegal rays	6-X-7
Caudal fin	6-9, 9+8, 6-9
Pelvic fin	Abdominal in larvae, absent in adults S: 1-1-1 R: 4-4-4
Dorsal fin	R: 52-X-56
Pectoral fin	R: 20-X-21
Anal fin	R: 34-X-44
Gill rakers	U: 1-1-1 L: 6-6-6

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesopelagic
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Spring-early summer, early fall; ^a winter ^b Area: Pelagic ^a Mode: Migration: To coastal areas ^b
Fecundity	Range/function: 230,000-430,000 ^b
Age at first maturity	3-4 yr ^b
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	2.8-3.1 mm
No. of oil globules	One
Oil globule diameter	Initially 0.42-0.60 mm
Yolk	Homogeneous, opaque becoming clear
Envelope	Smooth
Hatch size	6.5 mm SL
Incubation time/temp.	
Pigment	<ul style="list-style-type: none"> • Yolk • Oil globule • Embryo: Finfolds, above and below tail, body

Diagnostic characters

- Large size
- Large oil globule, decreasing in size with embryonic development

LARVAE

Preanal length	~40% SL
Length at flexion	11-17 mm SL
Length at transformation	
Sequence of fin development	Pectorals, pelvics, dorsal, anal; caudal not complete in 28.5 SL specimen

Pigment

- Head and gut covered with discrete spots
- Dorsal body margin
- Caudal finfold pigment becoming less prominent with development
- Pectoral and pelvic fin bases (postflexion)
- Three opposing blotches on dorsal and ventral finfolds in preflexion larvae


Diagnostic characters

- Preflexion pigment: Three opposing blotches in median finfold, one in caudal finfold
- Pelvic fin present in larvae, lost in adults
- Dorsal and anal fin begin to develop in finfold, have deep bases
- Morphological changes with development, from elongate to deep-bodied
- Blunt head
- Small preopercular spines

^aNWAFIC, unpubl.

^bFitch and Lavenberg 1971

Ref: Matarese et al. 1984b; Matarese, unpubl.



Figures A–D, Matarese et al. 1984b.

MERISTICS

Vertebrae	Total: 65-67-74 Precaudal: 40-44-47 Caudal: 23-24-25
Branchiostegal rays	6-X-8
Caudal fin	X, 8+7, X
Pelvic fin	Absent
Dorsal fin	R: 54-X-63
Pectoral fin	R: 13-14-15
Anal fin	R: 24-X-32
Gill rakers	U: 3-X-6 L: 16-X-22

LIFE HISTORY

Range	S. California, 32-34°N, to Arctic, not specific
Ecology	Epi- and mesobenthal, intertidal to 275 m
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Nov-Feb ^a Areas: In areas of strong current ^b Mode: Migration:
Fecundity	Range/function: 1000 ^c (<i>A. personatus</i> , western Pacific)-22,100 ^d (southwestern Barents Sea)
Age at first maturity	1 yr ^e (western Pacific)
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	0.67-0.91 mm (0.80 mm)
No. of oil globules	One
Oil globule diameter	~0.26 mm
Yolk	
Envelope	
Hatch size	6-7 mm SL (as small as 4 mm)
Incubation time/temp.	2-12 wk
Pigment	• Embryo: Eyes, dorsal and ventral body

Diagnostic characters

LARVAE


Preanal length	~60% SL
Length at flexion	11-13 mm SL
Length at transformation	16-31 mm SL
Sequence of fin development	Caudal, pectorals, dorsal and anal
Pigment	• Gut • Double row postanal ventral pigment

Diagnostic characters (see Table 4)

- Elongate body
- Gut length (~60% SL)
- Lightly pigmented
- Postanal ventral pigment (double row)
- Dorsal and anal fins begin development opposed to each other
- Elongate head shape in larger specimens

^aTrumble 1973
^bAndriashev 1954
^cInoue et al. 1967
^dMacy et al. 1978
^eHamada 1966

Ref: Kobayashi 1961c, Stevens et al. 1984.



Figures A–B, D, (B, ventral view), NWAFC originals (B. Vinter); C, Stevens et al. 1984.

Gobiidae is the most speciose family of marine fishes, although some species occur in brackish or freshwater environments. Gobies are generally small bottom-dwelling fishes with pelvic fins united to form a sucking disc. Found mainly in subtropical and tropical areas, adults inhabit shallow to moderate depths in salt and brackish water (some in freshwater). The study area includes three species: *Clevelandia ios*, *Coryphopterus nicholsi*, and *Lepidogobius lepidus*. Larvae are easily recognized by their conspicuously pigmented swimbladder and pigment patterns. Larvae are commonly collected inshore and in bays and estuaries. They are rare in coastal ichthyoplankton collections in the study area.

Small larvae of *Clevelandia* and *Lepidogobius* have been confused for some time in the literature. Since we have incomplete developmental series of the two species, we are presenting a consensus of opinion from researchers who have had more experience studying or collecting gobies. Gobiid larvae with three dorsal melanophores previously assigned to *Clevelandia* are now considered *Lepidogobius* (Wang 1986; W. Watson and G. McGowen, pers. commun.¹).

¹W. Watson, Marine Ecological Consultants, 531 Encinitas Blvd., Suite 110, Encinitas, CA 92024, pers. commun., 3 Nov. 1986; G. McGowen, Los Ang. Cty. Mus. Nat. Hist., 900 Exposition Blvd., Los Angeles, CA 90036, pers. commun., 31 Oct. 1986.

Table 47
Early-life-history characters of gobiid larvae from the Northeast Pacific (Wang 1981, in part).

	<i>Clevelandia ios</i>	<i>Coryphopterus nicholsi</i>	<i>Lepidogobius lepidus</i>
Spawning Site	Burrow	Rocky reef	Burrow
Egg Shape	Elliptical with narrow, blunt distal end	Spindle-shaped, narrow, elongate	Elliptical
Larvae			
Total myomeres	34-36	25-26	36-38
Preanal myomeres	16-18	9-10	14-17
Postanal myomeres	16-19	14-17	19-23
Distinguishing pigmentation	Single large melanophore along dorsal midline of body which forms a band at about myomere 26	A series of 10-15 melanophores along ventral body midline and a shorter series along dorsal midline near caudal	Three dorsal midline melanophores, posteriormost forming a band at about myomeres 19-26
Juveniles			
Dorsal fin	IV-VI; 0-I, 14-17	V-VI; I-II, 9-15	VI-IX; 0-I, 14-18
Anal fin	0-I, 14-17	0-I, 11-14	0-I, 13-16
Pectoral fin	18-21	21-24	20-22
Vertebrae	35-37	26	37-38
Distribution	Seawater-polyhaline-oligohaline	Seawater-polyhaline	Seawater-polyhaline-oligohaline
Distinguishing characteristics	~12 dark bands on dorsum; ~12 close melanophores on lateral line	Large black eye; black margin on spinous dorsal fin; side of body without dark vertical bands	Black margin on spinous dorsal fin; broad dark band at base of caudal region; body pigmentation light

MERISTICS

Vertebrae	Total: 35-36-37 Precaudal: 15-15-15 Caudal: 21-21-22
Branchiostegal rays	3-X-5
Caudal fin	
Pelvic fin	Thoracic R: 5-5-5
Dorsal fin	1st S: 4-X-6 2nd S: 0-X-1 R: 14-X-17
Pectoral fin	R: 20-20-20
Anal fin	S: 0-X-1 R: 14-X-17
Gill rakers	U: 1-X-3 L: 5-X-7

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Intertidal
ELH pattern	Oviparous; demersal, attached eggs; pelagic larvae
Spawning	Season: Nov-June or later ^a Area: Mode: Migration:
Fecundity	Range/function: 750-1000 (may be multiple spawners) ^{b/} $F=0.0306 \times L^{2.04}$ ^c
Age at first maturity	1 yr ^c
Longevity	2-3 yr ^c

^aWang 1981^bPrasad 1958^cBrothers 1975

Ref: Prasad 1958, Ruple 1984, Wang 1981.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	0.70-0.85 mm
No. of oil globules	Many coalescing to one
Oil globule diameter	
Yolk	
Envelope	Transparent; adhesive threads at one pole
Hatch size	2.7-3.8 mm SL
Incubation time/temp.	10-12 d/15-15.5°C
Pigment	

Diagnostic characters

- Ellipsoidal

LARVAE

Preanal length	45-52% SL
Length at flexion	
Length at transformation	
Sequence of fin development	Caudal, 2nd dorsal and anal, 1st dorsal and pectorals, pelvics

Pigment



- Single large melanophore along dorsal midline of body which forms a band at about myomere 26
- Dorsal surface of swimbladder

Diagnostic characters (see Table 47)

- Conspicuously pigmented swimbladder (family)
- Distinguished from *Coryphopterus nicholsi* by
- Total myomeres (fewer in *C. nicholsi*, 25-26 vs. 35-37)
- Distinguished from *Lepidogobius lepidus* by
- Pigment pattern: No anterior melanophores along dorsal midline; present in *L. lepidus*

Postflexion

- Dorsal spine count (4-6), anal count (14-17), mouth size



Figures A–B, NWAFC originals (B. Vinter; specimens loaned by Bruce Mundy, formerly of Oregon State University. The 15.0 mm SL specimen was slightly damaged, so the swimbladder could not be accurately illustrated).

MERISTICS

Vertebrae	Total: 26-26-26 Precaudal: 11-11-11 Caudal: 15-15-15
Branchiostegal rays	3-X-5
Caudal fin	
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	1st S: 4-X-8 2nd S: 1-X-2 R: 9-X-15
Pectoral fin	R: 16-X-23
Anal fin	S: 0-X-1 R: 11-X-13
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Nearshore shelf demersal, intertidal to 106 m
ELH pattern	Oviparous; demersal, attached eggs; pelagic larvae
Spawning	Season: Feb ^a -Oct ^b (California) Area: Under rocks ^a Mode: Pairs; eggs guarded by males ^a Migration:
Fecundity	Range/function: 3274-4788 (may be multiple spawners) ^a
Age at first maturity	2-5 yr (females) ^a 3-5 yr (males) ^a
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	2.1×0.48 mm
No. of oil globules	Multiple
Oil globule diameter	
Yolk	
Envelope	Transparent, smooth
Hatch size	2.94 mm TL
Incubation time/temp.	
Pigment	

Diagnostic characters

- Ellipsoidal (pointed at each end)
- Pigmented dorsally over yolk and along ventral body midline

LARVAE

Preanal length	50% SL
Length at flexion	
Length at transformation	
Sequence of fin development	Caudal, 2nd dorsal and anal, 1st dorsal and pectorals, pelvics

Pigment

- Dorsal surface of swimbladder
- Dorsal surface of posterior gut
- Series of dorsal midline melanophores along posterior half of body
- 10-15 postanal ventral melanophores
- With development, pigment in hypural area

Diagnostic characters (see Table 47 and *Clevelandia ios*, p. 544)

- Myomeres (26)

^aWiley 1973^bEbert and Turner 1962

Ref: Ruple 1984, Wang 1981.




Figure A, NWAFC original (B. Vinter; specimen loaned by Bruce Frost, University of Washington, for illustration).

MERISTICS

Vertebrae	Total: 37-37-38 Precaudal: 15-15-15 Caudal: 22-22-23
Branchiostegal rays	3-X-4
Caudal fin	
Pelvic fin	Thoracic R: 5-5-5
Dorsal fin	1st S: 6-X-9 2nd S: 0-X-1 R: 14-X-18
Pectoral fin	R: 20-20-20
Anal fin	S: 0-X-1 R: 13-X-16
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Nearshore shelf demersal, intertidal to 201 m
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Nov-June (California) ^a Area: Intertidal mudflats ^a Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.3-1.8 × 0.8-1.0 mm (unfertilized)
No. of oil globules	Many (unfertilized)
Oil globule diameter	
Yolk	Granular, yellowish
Envelope	Transparent, smooth
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	42-45% TL
Length at flexion	
Length at transformation	
Sequence of fin development	Caudal, 2nd dorsal and anal, 1st dorsal and pectorals, pelvics

Pigment

- Dorsal surface of swimbladder
- Three dorsal midline melanophores, posteriormost forming a band at about myomeres 19-26

Diagnostic characters (see Table 47 and *Clevelandia ios*, (p. 544)

Distinguished from *C. ios* by



- Anterior melanophores along dorsal midline; not present in *C. ios*

Distinguished from *Coryphopterus nicholsi* by

- Total myomeres (fewer in *C. nicholsi*, 25-26 vs. 37-38)

^aWang 1981

Ref: Ruple 1984, Wang 1981.



Figures A–B, NWAFC originals (B. Vinter; specimens loaned by Bruce Mundy, formerly of Oregon State University, for illustration).

There are four species of cutlassfishes in three genera of this family in the northeastern Pacific. General body shape is elongate and ribbonlike, tapering to a small caudal fin. Voracious predators, most have a large, well-toothed mouth and pointed snout. Adult cutlassfishes are benthopelagic but have been found at the surface at night (Fritzsche 1978). Spawning occurs offshore, resulting in pelagic eggs which can be 1.6-2.5 mm and have a single reddish-yellow oil globule (Breder and Rosen 1966, Fritzsche 1978). Pelagic larvae have a high myomere count (generally >100) and develop three dorsal spines, the first of which may be elongate (Schmidt and Strubberg 1918). Larvae have a small head with a long, tapering body (Ozawa 1986e). No early-life-history stages have been collected in our area.

Table 48
Meristic characters of family Trichiuridae.

Taxon	Distribution	Vertebrae		Fins				Gill rakers		Branchiostegals
		Precaudal	Caudal	Dorsal	Anal	Pectoral	Pelvic	Upper	Lower	
<i>Aphanopus carbo</i>	N. Calif.-Brit. Col.	42-46	55-59	XXXVIII-XLIII, 53-56	I-II, 44-49	12	I,1			7-8
<i>Benthodesmus elongatus</i>	Cent. Calif.-Brit. Col.	(148-153)		XLIV-XLVII, 98-102	II, 91-98	12	I,1	5	9	7
<i>Benthodesmus tenuis</i>	Brit. Col.	(121-131)		XXXIX-XLII, 79-88	II, 69-75					
<i>Lepidopus fitchi</i>	SSC-Oregon	35 (84-92)	50	IX, 78-86	II, 41-49	12	I,1*	7	10	7

*The soft ray is rudimentary (Rosenblatt and Wilson 1987).

MERISTICS *B. elongatus*
(Steindachner 1891)

Vertebrae	Total: 148-X-153
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	
Pelvic fin	Thoracic
	S: 1-1-1 R: 1-1-1
Dorsal fin	S: 44-44-47 R: 98-X-102
Pectoral fin	R: 12-12-12
Anal fin	S: 2-2-2 R: 91-93-98
Gill rakers	U: 5-5-5 L: 9-9-9

MERISTICS *B. tenuis* (Günther)

Vertebrae	Total: 121-X-131
	Precaudal: X-X-X
	Caudal: X-X-X
Branchiostegal rays	7-7-7
Caudal fin	
Pelvic fin	Thoracic
	S: X-X-X R: X-X-X
Dorsal fin	S: 39-X-42 R: 79-X-88
Pectoral fin	R: X-X-X
Anal fin	S: 2-2-2 R: 69-X-75
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Cent. California, 34-38°N, ^a to Brit. Col., 48°30'-55°N
Ecology	Epi- and mesopelagic
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

^a*B. tenuis* range restricted to British Columbia.

^bDescription based on *B. elongatus simonyi*, a subspecies from the North Atlantic, and *B. elongatus pacificus* from Japan. Ozawa (1986e) describes and figures one late postflexion larva of *B. tenuis* (27.1 mm SL). Larvae of *B. elongatus* and *B. tenuis* from our area may differ.

^cOzawa 1986e

Ref: Collette et al. 1984b, Evseenko 1982, Gorbunova 1982b, Ozawa 1986e, Peden and Hughes 1986.

EARLY LIFE HISTORY DESCRIPTION

EGGS - Family

Diameter	1.7-2.0 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	4.5-6.5 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE^b

Preanal length	
Length at flexion	
Length at transformation	
Sequence of fin development	Dorsal first fin to develop
Pigment	
	<ul style="list-style-type: none"> • Blotch on upper preopercle • Crown • Dorsal midline beginning just anterior to dorsal fin origin and running along developing fin, increasing with development • Dorsolateral gut • Ventrolateral blotch about mid-postanal body; with development becoming a series along ventral midline about 3/4 length of anal fin • <i>B. elongatus pacificus</i> has a blotch at the origin of the anal fin (not shown on figure)^c

Diagnostic characters

- Family
 - Serrate spines in dorsal, anal, and pelvic fins
 - In some genera (e.g., *Benthodesmus*), the anterior-most rays of the dorsal fin are extremely elongate in the smallest larvae
- Genus
 - Caudal fin development distinct
 - Pelvic fin located approximately below pectoral fin
 - Number of dorsal fin spines (>30)


BENTHODESMUS

Figure A, Collette et al. 1984b (Gulf of Mexico specimen); B, Gorbunova 1982b; C, Evseenko 1982 (B–C, redrawn; subspecies from North Atlantic).

MERISTICS

Vertebrae	Total: 30-31-31 Precaudal: 14-14-15 Caudal: 16-17-17
Branchiostegal rays	7-7-7
Caudal fin	X, 9+8, X
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 9-10-12 R: 11-12-12
Pectoral fin	R: 20-20-20
Anal fin	S: 1-1-1 R: 11-X-12
Gill rakers	U: 11-13-14 L: 27-28-30

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59°N
Ecology	Epi- and mesopelagic, 0-300 m ^a
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Apr-July (California) ^b Area: 0-72 m, 3-32 km from shore ^c Mode: Migration:
Fecundity	Range/function: >1 million (may spawn more than once each year) ^b
Age at first maturity	2 yr ^b -6 yr ^d
Longevity	10 yr ^b ; 12 yr ^e

^a Collette and Nauen 1983^b Hart 1973^c Fritzsche 1978^d Schaefer 1980^e Fitch and Lavenberg 1971

Ref: Berrien 1978, Collette et al. 1984b, G.D. Johnson 1984, Kramer 1960.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	0.9-1.3 mm
No. of oil globules	One, ventral
Oil globule diameter	0.26 mm
Yolk	Homogeneous
Envelope	Smooth, clear
Hatch size	3.3 mm SL
Incubation time/temp.	
Pigment	
• Oil globule	
• Yolk	

Diagnostic characters

- Posterior position of oil globule

LARVAE


Preanal length	52%, increasing with development to 64% SL
Length at flexion	6 mm SL
Length at transformation	15 mm SL
Sequence of fin development	2nd dorsal (rays) and anal, 1st dorsal (spines), pectorals and pelvics

Pigment

- Pigment on crown extends anteriorly with development
- Dorsolaterally on gut
- Ventral midline
- Postflexion
 - Short dorsal midline series develops under dorsal fin, spreading anteriorly
 - Ventral midline becomes restricted to posterior half of body
 - Mediolateral pigment

Diagnostic characters

- Large head with teeth
- Myomeres (30-31)
- Pigment pattern



Figures A–E, Kramer 1960 (redrawn).

MERISTICS

Vertebrae	Total: 22-22-23	
	Precaudal: 10-10-10	
	Caudal: 12-12-13	
Branchiostegal rays	5-X-6	
Caudal fin	X, 16, X	
Pelvic fin ^a	Thoracic	
	S: 1-1-1	R: 4-4-4
Dorsal fin ^a	S: 2-2-2	R: 24-24-24
Pectoral fin	R: 17-X-20	
Anal fin ^a	S: 14-X-17	
Gill rakers	U: 4-X-6	L: 11-X-14

LIFE HISTORY

Range	South of southern California to Washington, 46-48°30'N
Ecology	Epi- and mesopelagic
ELH pattern	Parity and eggs unknown, pelagic larvae
Spawning	Season: Spring-early summer ^b Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

^aMeristics for larvae and adults are different:

	Larvae	Adults
Dorsal fin rays	24	11-14
Anal fin rays	18	13-15
Pelvic fin rays	4	absent

^bFitch and Lavenberg 1971^cBased on illustrations only, specimens were not available.^dG.D. Johnson, Natl. Mus. Nat. Hist., Wash., D.C. 20560, pers. commun., 7 Nov. 1986.

Ref: Leis and Richards 1984, Nishikawa 1987.

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	<3.5 mm SL
Incubation time/temp.	
Pigment	





Diagnostic characters

LARVAE

Preanal length	~60% SL, decreasing with development
Length at flexion	
Length at transformation	
Sequence of fin development	Pelvics and anterior dorsal fin spines form early
Pigment ^c	<ul style="list-style-type: none"> • Few spots on upper jaw • With development above urostyle • Lightly on caudal fin, hypural region with several spots • Gut • With development on pectoral fin rays

Diagnostic characters

- Morphology
 - Deep-bodied but not as kite-shaped as acanthurids
 - Large square-shaped head with small terminal mouth
 - Extensive head spination
 - With development, minute spines on soft rays and along body surface
 - Dorsal and pelvic spines elongate, finely serrated
 - Loss of meristic elements with growth
- Not shown on figure^d
 - Spines on ascending process of premaxillary bone (important feature uniquely shared with *Zanclus canescens*, a closely related acanthuroid, and acanthurids)
 - Small dorsal spine anterior to first dorsal spine appears later in development



Figures A–D, Nishikawa 1987 (western Pacific specimen).

MERISTICS

Vertebrae	Total: 56-59-62
	Precaudal: 22-24-25 (adults) ^a
	Caudal: 33-35-38
Branchiostegal rays	7-7-7
Caudal fin	11-14, 9+8, 10-13
Pelvic fin	Thoracic
	S: 1-1-1 R: 5-5-5
Dorsal fin	S: 3-3-3 R: 34-39-43
Pectoral fin	R: 18-X-21
Anal fin	S: 3-3-3 R: 20-X-29
Gill rakers	U: 4-X-6 L: 12-X-14

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epipelagic, 0-91 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.52-1.80 mm
No. of oil globules	One
Oil globule diameter	0.30-0.44 mm (large)
Yolk	Homogeneous
Envelope	Clear, smooth
Hatch size	
Incubation time/temp.	
Pigment	<ul style="list-style-type: none"> • Dorsal pigment along body • Ventral pigment from head to tip of tail • Underside of oil globule becomes pigmented with development

Diagnostic characters**LARVAE**

Preanal length	50-58% SL
Length at flexion	~9.3-11.0 mm SL
Length at transformation	~20 mm SL
Sequence of fin development	Caudal, 2nd dorsal (rays), anal, pectorals, 1st dorsal (spines), pelvics


Pigment

- Dorsal and ventral body midline pigment
- Lateral line dashes develop during flexion
- Head and tail pigment increases with development
- Become heavily, uniformly pigmented including fin membranes

Diagnostic characters

- Distinguished from *Tetragonurus cuvieri* (p. 560) by
- High myomere counts (56-62)
 - Preflexion: Dorsal body pigment over posterior half of body; only on tail in *T. cuvieri*
 - Flexion: No preopercular spines
 - Postflexion
 - Dorsal and anal spines weak, few (three) dorsal spines
 - More dorsal and anal fin rays than in *T. cuvieri*

^aNumber of preanal myomeres appears to decrease with development (B. Sumida, NMFS Southwest Fish. Cent., P.O. Box 271, La Jolla, CA 92038, pers. commun., 16 Dec. 1986).



Figures A–F, Ahlstrom et al. 1976.

MERISTICS

Vertebrae	Total: 52-53-57 Precaudal: 25-28-29 (adults) ^a Caudal: 24-26-28
Branchiostegal rays	5-6-6
Caudal fin	9-13, 9+8, 9-12
Pelvic fin	Thoracic S: 1-1-1 R: 5-5-5
Dorsal fin	S: 15-17-21 R: 10-12-17
Pectoral fin	R: 14-X-17
Anal fin	S: 2-2-2 R: 9-X-15
Gill rakers	U: 6-6-6 L: 7-X-14

LIFE HISTORY

Range	South of southern California to Aleutian Is., 51-55°N
Ecology	Epipelagic
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.10-1.28 mm
No. of oil globules	One
Oil globule diameter	0.25-0.30 mm (amorphous)
Yolk	Homogeneous
Envelope	Smooth, golden
Hatch size	4.0-4.1 mm SL

Incubation time/temp.**Pigment**

- Oil globule
- Distinctive pigment on embryo; double dorsal line separating prior to head, outlining brain, and extending forward to snout
- Ventral pigment above digestive tract, continuing along tail

Diagnostic characters

- Pigment

LARVAE

Preanal length	60-70% SL
Length at flexion	7.6-10.1 mm SL
Length at transformation	~21.4 mm SL
Sequence of fin development	Caudal; 2nd dorsal (rays), anal, and pectorals; 1st dorsal (spines); pelvics


Pigment

- Tail
- Lateral line notable during flexion
- Eye bar
- Dorsal midline on caudal peduncle extending anteriorly with development

Diagnostic characters

- Distinguished from *Icichthys lockingtoni* (p. 558) by
- Myomere count (52-57)
 - Pigment
 - Notochord tip and base of caudal
 - Ventral row over gut to tail
 - Early formation of opposing 2nd dorsal and anal fins
 - Slender body with long caudal peduncle

^aNumber of preanal myomeres appears to decrease with development (B. Sumida, NMFS Southwest Fish. Cent., P.O. Box 271, La Jolla, CA 92038, pers. commun., 16 Dec. 1986).



Figures A–E, Ahlstrom et al. 1976.

MERISTICS

Vertebrae	Total: 28-30-31 Precaudal: 11-13-14 (adults) ^a Caudal: 17-17-19
Branchiostegal rays	6-6-6
Caudal fin	7-9, 9+8, 6-8
Pelvic fin	Absent
Dorsal fin	S: 2-3-4 R: 41-45-48
Pectoral fin	R: 19-21-23
Anal fin	S: 2-3-3 R: 35-39-44
Gill rakers	U: 3-3-3 L: 11-11-11

LIFE HISTORY

Range	South of southern California to Brit. Col., 48°30'-55°N
Ecology	Epipelagic, 9-91 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Spring-summer (California) ^b Area: Mode: Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	
No. of oil globules	One
Oil globule diameter	0.2 mm (in yolksac larva)
Yolk	
Envelope	
Hatch size	1.8-2.0 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	55%, with development decreasing to 44% SL
Length at flexion	5-7 mm SL
Length at transformation	~20 mm SL
Sequence of fin development	

Pigment

- Anterior body
- Median ventral series of melanophores from isthmus to anus
- Large melanophores on head and anterior half of body, becoming uniformly pigmented on anterior 3/4 of body, later over whole body


Diagnostic characters

- Pigment
- Increase in relative body depth during ontogeny
- Absence of pelvic fins


^aNumber of preanal myomeres appears to decrease with development (B. Sumida, NMFS Southwest Fish. Cent., P.O. Box 271, La Jolla, CA 92038, pers. commun., 16 Dec. 1986).

^bFitch and Lavenberg 1971

Ref: D'Vincent et al. 1980.



Figures A–F (B, dorsal view), D’Vincent et al. 1980.



Pleuronectiformes

Pleuronectiforms are benthic fishes that are asymmetrical with both eyes on one side of the head. Asymmetry is also reflected in dentition, cranial osteology, pelvic fin placement and morphology, pigment, and squamation. Worldwide in distribution, there are 7 families, with well over 120 genera and 500 species (Ahlstrom et al. 1984a). Most families are defined as dextral (eyes on right side) or sinistral (eyes on left side), although some species are indiscriminate. A total of 3 families with 31 species occur in the study area. Much is known about the early life history of many species. Eggs of most are pelagic, spherical, and have homogeneous yolks. Chorions may be smooth, striated, or ornamented with raised polygonal patterns (Ahlstrom et al. 1984a). Early larvae are bilaterally symmetrical and swim upright. Of help in larval identification are body shape (short gut, elongate body), pigment patterns (postanal bands, pigment on the urostyle and finfolds), and meristic characters. Size at transformation and the development of spines and elongate dorsal or pelvic fin rays can be diagnostic. During transformation, the eyes migrate to their adult position and the larvae assume the distinctive flat profile unique to the order. At this point, juveniles settle and become benthic inhabitants.

Families in study area: **Paralichthyidae**
Pleuronectidae
Cynoglossidae

Table 49
Characters useful in identifying pleuronectiform larvae. Size data are after Ahlstrom et al. (1984a), in part.
All measurements are in millimeters NL or SL. Taxa listed alphabetically within families.

Species ^a	Size at hatching	Size at notochord flexion	Size at transformation	Total myomeres
Paralichthyidae				
<i>Citharichthys sordidus</i>	2.0	10.0-11.4	20.0-39.0	39-40
<i>C. stigmaeus</i>	2.0	9.2-10.2	24.0-36.0	36-39
Pleuronectidae				
<i>Acanthopsetta nadeshnyi</i>	<3.0	8.4-9.9	20.0-24.0	39-40 ^b
<i>Atheresthes evermanni</i>	<8.4	11.5-15.0	—	—
<i>A. stomias</i>	<10.0	10.0-12.0	33.0- ?	47-50
<i>Clidoderma asperrimum</i>	—	—	—	42-44 ^b
<i>Embassichthys bathybius</i>	9.0	15.4-16.2	16.2- ?	57-65
<i>Eopsetta jordani</i>	2.8	—	—	41-45
<i>Glyptocephalus stelleri</i>	4.1-5.2	15.0-17.0	19.0-48.0	52-60
<i>G. zachirus</i>	5.0-6.0	15.3-24.0	49.0-72.0	63-66
<i>Hippoglossoides elassodon</i>	5.3-6.9	9.0-10.2	18.0- ?	43-47
<i>H. robustus</i>	4.0	11.0- ?	>28.6	44
<i>Hippoglossus stenolepis</i>	7.8-8.5	13.6-17.8	14.7-24.1	49-51
<i>Inopsetta ischyra</i>	—	—	—	41
<i>Isopsetta isolepis</i>	2.7-2.9	9.1-14.0	15.0->21.9	41-42
<i>Lepidopsetta bilineata</i>	3.4-4.0	8.4-9.9	>17.7	39-42
<i>Limanda aspera</i>	2.2-2.8	7.5-9.5	10.0- ?	40-41
<i>L. proboscidea</i>	<4.8	—	—	—
<i>Liopsetta glacialis</i>	3.7	—	—	37-41
<i>Lyopsetta exilis</i>	5.6	9.0-10.9	15.7-24.7	43-47
<i>Microstomus pacificus</i>	6.0	10.0-15.0	20.0->45.0	50-55
<i>Parophrys vetulus</i>	2.3-2.8	8.8-10.5	17.5- ?	42-47
<i>Platichthys stellatus</i>	1.9-2.1	5.5-6.0	8.0- ?	35-38
<i>Pleuronectes quadrituberculatus</i>	5.6	8.0-10.0	10.0- ?	41-42
<i>Pleuronichthys coenosus</i>	3.9	6.2-8.5	8.2->11.4	37-39
<i>P. decurrens</i>	4.9-5.5	7.8-11.0	10.5->21.0	38-41
<i>Psettichthys melanostictus</i>	<3.0	8.0-10.0	>22.6	38-41
<i>Reinhardtius hippoglossoides</i>	10.0-16.0	25.0-27.0	45.0-65.0	61-64
Cynoglossidae				
<i>Symphurus atricauda</i>	1.9	9.4-10.8	19.0-24.2	50-52

^aThe following nomenclatural changes have been recommended by Sakamoto (1984):

<i>Glyptocephalus</i> (= <i>Errex</i>) <i>zachirus</i>	<i>Liopsetta</i> (= <i>Pleuronectes</i>)
<i>Isopsetta</i> (= <i>Pleuronectes</i>)	<i>Parophrys</i> (= <i>Pleuronectes</i>)
<i>Lepidopsetta</i> (= <i>Pleuronectes</i>)	<i>Lyopsetta</i> (= <i>Eopsetta</i>)
<i>Limanda</i> (= <i>Pleuronectes</i>)	

^bM. Okiyama, Ocean Res. Inst., Univ. Tokyo, 1-15-1 Minamidai, Nakano-Ku, Tokyo 164, Japan, pers. commun., 17 Oct. 1986.

MERISTICS

Vertebrae	Total: 39-X-40 Precaudal: 11-11-12 Caudal: 27-28-29
Branchiostegal rays	6-X-7
Caudal fin	X, 7+6, X Total rays=17 ^a
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 86-91-102
Pectoral fin	R: 12-12-12
Anal fin	R: 67-72-81
Gill rakers	U: 6-X-9 L: 12-13-16

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathybenthal, 0-549 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Mar-May (Puget Sound) ^b ; July-Sept (Calif.); may spawn twice each season ^c Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	2-3 yr ^b
Longevity	

^a 11 branched rays.^b Smith 1936^c Arora 1951^d Ahlstrom et al. 1984a

^e Early preflexion larvae (<6-7 mm SL) of *C. sordidus* and *C. stigmaeus* are not identifiable to species until the elongate dorsal fin rays form on *C. sordidus* (~7 mm SL). Preflexion larvae collected in our study area do not resemble the published illustrations of either species (Ahlstrom and Moser 1975); they have ventral gut pigment extending more anteriorly, a series of postanal ventral midline melanophores, no finfold pigment, and less intense caudal pigment (Fig. A). Specimens >6.9 mm SL that have elongated dorsal fin rays are *C. sordidus* (Fig. B), but do not exhibit the intense caudal pigment shown by Ahlstrom and Moser (1975). Our specimens indicate a decrease in intensity of caudal pigment with development for *C. sordidus* and an increase for *C. stigmaeus*. The preflexion larva of *C. sordidus* illustrated by Ahlstrom and Moser (1975) may belong to another species probably south of our study area, most likely *C. xanthostigma* (B. Sumida, NMFS Southwest Fish. Cent., Box 271, La Jolla, CA 92038, pers. commun., 16 Dec. 1986).

Ref: Ahlstrom 1965, Ahlstrom and Moser 1975, Ahlstrom et al. 1984a.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	0.55-0.77 mm; ^c 0.51-0.79 mm ^d
No. of oil globules	One
Oil globule diameter	0.09-0.12 mm
Yolk	Homogeneous
Envelope	Smooth
Hatch size	~2.0 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

We cannot identify *Citharichthys* spp. eggs to species. We collect two types of paralichthyid eggs: the first type has an average diameter of ~0.64-0.68 mm, and the other type averages ~0.78-0.88 mm.

LARVAE^e


Preanal length	<50% SL
Length at flexion	<10.0-11.4 mm SL
Length at transformation	20->39 mm SL
Sequence of fin development	Dorsal, anal, caudal, pelvics, pectorals <u>or</u> caudal, dorsal and anal, pelvics, pectorals

Pigment

- Ventral body midline pigment posterior to anus, coalesces later into two melanophores
- Band at 3/4 BL
- Posteriorly on gut
- Tips of elongate dorsal and pelvic fin rays

Diagnostic characters

- Elongate dorsal and pelvic fin rays



Figures A–B, NWAFC originals (B. Vinter); C–D, Ahlstrom and Moser 1975.

MERISTICS

Vertebrae	Total: 36-37-39
	Precaudal: 9-10-10
	Caudal: 27-28-29
Branchiostegal rays	6-X-7
Caudal fin	2, 7+6, 2
	Total rays = 17 ^a
Pelvic fin	Thoracic
	R: 6-6-6
Dorsal fin	R: 75-84-97
Pectoral fin	R: 12-12-12
Anal fin	R: 58-64-77
Gill rakers	U: 3-X-5 L: 7-8-10

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59°N
Ecology	Epi- and mesobenthal, 0-366 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Spring-summer (Calif.) ^b Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	0.75-0.83 mm
No. of oil globules	One
Oil globule diameter	0.09-0.10 mm
Yolk	Homogeneous
Envelope	Smooth
Hatch size	2 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

See notes on *C. sordidus* (p. 568)

LARVAE

Preanal length	
Length at flexion	9.2-10.2 mm SL
Length at transformation	24.0-35.5 mm SL
Sequence of fin development	Dorsal, anal, caudal, pelvics, pectorals <u>or</u> caudal, dorsal and anal, pelvics, pectorals

Pigment (see *C. sordidus*)

- Urostyle pigment increases in intensity to flexion stage, then fades
- Small melanophores along ventral body midline posterior to anus; these migrate onto anal fin pterygiophores in postflexion larvae
- Band at 3/4 BL, lost after flexion
- Caudal fin


Diagnostic characters

- Lack of elongate fin rays

^a 11 branched rays.

^b Ahlstrom and Moser 1975

Ref: Ahlstrom et al. 1984a.



Figures A–B, Ahlstrom and Moser 1975.

The majority of species in the order (27 out of a total of 31) belong to the family Pleuronectidae. Ahlstrom et al. (1984a) summarize the following ontogenetic characters for the family.

Egg size ranges from 0.66 to 4.50 mm, yolk is homogeneous, oil globules are absent in most species, and the chorion is smooth (exceptions include *Pleuronichthys ritteri* and *P. cornutus*). Some light sculpturing of the chorion is seen in several species.

Larvae hatch from 1.7 to 16.0 mm SL. They possess no elongate dorsal or pelvic fin rays, gut is normal, and spines are limited to the preopercular, otic, and frontal regions, although absent in most.

The caudal fin ray count is not stabilized within pleuronectids, in contrast to most other taxa. Most pleuronectids have 17-19 total caudal fin rays (18 is most common). Within species there is variability in the total fin ray count and in the number of rays that are branched. Since variability exists, total counts, number of branched rays, and principal and secondary rays are given when available.

Table 50
Pigment characters and myomere counts that may be useful in the identification of pleuronectid larvae during the preflexion and early flexion stages.*

Genus	Pigment characters					Total myomeres
	Number of bands	Slashlike hypaxial pigment	Urostyle	Finfold		
				Dorsal	Anal	
Postanal pigment bands present						
<i>Embassichthys</i>	3	No	Yes	Yes	Yes	57-65
<i>Eopsetta</i>	2	No	Yes	No	No	41-45
<i>Glyptocephalus</i>	3-4	No	Yes	Yes/no	Yes/no	52-66
<i>Hippoglossoides</i>	3-4	Yes	Yes	Yes	Yes	44-51
<i>Isopsetta</i>	3	No	Yes	No	Yes	41-42
<i>Lepidopsetta</i>	1	No	No, few spots	Yes	Yes	39-42
<i>Microstomus</i>	4	No	Yes	Yes	Yes	50-55
Postanal pigment bands absent						
<i>Acanthopsetta</i>	—	Yes	Yes	No	No	39-40
<i>Atheresthes</i>	—	No	Yes/no	No	No	47-50
<i>Hippoglossus</i>	—	No	Yes	Yes	Yes	49-51
<i>Limanda</i>	—	Yes	No	No	Yes	40-41
<i>Liopsetta</i>	—	No	Yes	No	No	37-41
<i>Lyopsetta</i>	—	Yes	Yes	Yes	Yes	43-47
<i>Parophrys</i>	—	Yes	Yes	No	Yes	42-47
<i>Platichthys</i>	—	No	Yes	Yes	Yes	35-38
<i>Pleuronectes</i>	—	No	No	No	Yes	41-42
<i>Pleuronichthys</i>	—	No	Yes	Yes	Yes	38-41
<i>Psettichthys</i>	—	No	Yes	Yes	Yes	38-41
<i>Reinhardtius</i>	—	No	Yes	Yes	Yes	61-64

*Characters are discussed only for taxa where at least some early-life-history stages are known. Only general trends are presented since pigment may vary from specimen to specimen. In cases where actual specimens were not available, subjective decisions were made based on previously published illustrations.

MERISTICS

Vertebrae	Total: 39-X-42 Precaudal: 9-X-10 Caudal: 30-X-31
Branchiostegal rays	7-X-8
Caudal fin	Total rays=18
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 67-X-78
Pectoral fin	R: 9-X-11
Anal fin	R: 54-X-62
Gill rakers	U: X-X-X L: 10-X-12

LIFE HISTORY

Range	Bering Sea, 54-66°N
Ecology	Epi- and mesobenthal
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Summer ^a Area: Deep water ^a Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	0.86-1.03 mm (0.92 mm)
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	Smooth
Hatch size	<3 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE

Preanal length	<50% SL
Length at flexion	8.4-9.9 mm SL
Length at transformation	~20-24 mm SL
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals
Pigment	

- Series of ventral midline melanophores extending around urostyle
- Several spots above notochord, number increases with development
- During flexion, pigment develops along hypaxial myomeres
- Initially, gut pigment restricted to dorsal and anteroventral surface; with development it extends posterodorsally and along ventral surface


Diagnostic characters

- Pigment
 - Pigment pattern without bands
 - Presence of hypaxial pigment
 - Presence of pigment around urostyle


^a Pertseva-Ostroumova 1961

Ref: Ahlstrom et al. 1984a, Pertseva-Ostroumova 1961.

A




B Urostyle pigment



4.5 mm


C



8.4 mm

Lateral pigment along hypaxial myomeres


D



9.9 mm


Median finfolds pigmented

E



19.5 mm

F



22.5 mm

Figures A–F, Pertseva-Ostroumova 1961.

MERISTICS

Vertebrae	Total: 49-X-52 Precaudal: 10-X-11 Caudal: 39-X-41
Branchiostegal rays	7-X-8
Caudal fin	
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 95-107-112
Pectoral fin	R: 13-X-14
Anal fin	R: 75-X-92
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	Gulf of Alaska, 54-60°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthal
ELH pattern	Oviparous, bathypelagic eggs, pelagic larvae
Spawning	Season: Fall-winter ^a Area: Demersal, deep water ^a Mode: Migration:
Fecundity	Range/function: 130,000- 500,000 ^a
Age at first maturity	9-10 yr (females) ^a 6-7 yr (males) ^a
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	2.05-2.20 mm; 2.5-3.5 mm ^a
No. of oil globules	None
Oil globule diameter	
Yolk	Greenish yellow
Envelope	Thin, smooth, clear
Hatch size	<8.4 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**


Preanal length	
Length at flexion	~11.5-15.0 mm SL
Length at transformation	
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals
Pigment	<ul style="list-style-type: none"> Two dorsolateral patches which, with development, meet and extend over the caudal peduncle

Diagnostic characters

According to figures in Pertseva-Ostroumova (1961), pigment pattern and morphology are similar to *A. stomias*. No larvae of *A. evermanni* have been identified from the study area.

^aPertseva-Ostroumova 1961

Ref: Ahlstrom et al. 1984a, Pertseva-Ostroumova 1961.



Figures A–B, Pertseva-Ostroumova 1961 (sizes not given).

MERISTICS

Vertebrae	Total: 47-49-50 Precaudal: 12-12-12 Caudal: 35-37-38
Branchiostegal rays	7-X-8
Caudal fin	Total rays = 17 ^a
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 92-108-115
Pectoral fin	R: 14-X-15
Anal fin	R: 72-87-99
Gill rakers	U: 4-4-4 L: 11-X-13

LIFE HISTORY

Range	Cent. California, 34-38°N, to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathybenthal, 18-900 m
ELH pattern	Oviparous, mesopelagic eggs, mesopelagic larvae
Spawning	Season: Dec-Mar (Bering Sea); ^b Aug (Gulf of Alaska) ^c Area: 108-360 m ^c Mode: Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	20 yr ^d

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	~3 mm
No. of oil globules	None
Oil globule diameter	
Yolk	
Envelope	Smooth
Hatch size	
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**


Preanal length	<50% SL
Length at flexion	~10-12 mm SL
Length at transformation	25.6-33.0 mm SL
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals
Pigment	

- Pigment on midbrain, spreading with development to dorsal surface of head, lower jaw, opercle
- Dorsally on gut, spreading laterally with development
- Two dorsolateral patches which, with development, meet and increase in size to cover caudal peduncle posteriorly

Diagnostic characters

- Distinguished from other pleuronectids by
- Head spination: Early development of preopercular spines
 - Pigmented supraocular crest
 - Pigment pattern without bands; two dorsal patches with a lack of ventral pigment
 - Head and snout blunt
- Distinguished from *A. evermanni* by
- See *A. evermanni* (p. 576)

^a 13 branched rays.^b Pertseva-Ostroumova 1961^c Hirschberger and Smith 1983^d Kabata and Forrester 1974



Figures A–C, NWAFC originals (B. Vinter).

MERISTICS

Vertebrae	Total: 57-62-65 Precaudal: 13-14-14 Caudal: 44-48-51
Branchiostegal rays	7-X-8
Caudal fin	X, 10+9, X Total rays = 19-21 ^a
Pelvic fin	Thoracic R: 5-X-6
Dorsal fin	R: 109-X-117
Pectoral fin	R: 11-11-11
Anal fin	R: 94-X-98
Gill rakers	U: 6-X-9 L: 14-X-16

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Meso- and bathybenthal, 320-1433 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Spring (possibly winter-spring) ^b Area: Mode: Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

^a17 branched rays.

^bRichardson 1981b

^cAccording to Figures C and E from Richardson (1981b), flexion occurs rapidly between 15.4 and 16.2 mm SL. However, the 18.5-mm NL specimen (Fig. D) illustrated by Ahlstrom et al. (1984a) does not appear fully flexed. Differences in size may be a result of preservation.

Ref: Ahlstrom et al. 1984a, Richardson 1981b.

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	2.8-3.1 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Smooth
Hatch size	~9 mm SL
Incubation time/temp.	
Pigment	
	• Eye
	• Hindgut extending out into yolk
	• Three postanal bands

Diagnostic characters

- Size
- Pigment pattern


LARVAE

Preanal length	40% SL
Length at flexion	15.4-16.2 mm SL ^c
Length at transformation	>16.2 mm SL when eye migration commences
Sequence of fin development	Caudal, dorsal, and anal; pelvics; pectorals
Pigment	

- Initially, three postanal bands with pigment on the finfold in region of bands; number of dorsal and ventral spots change with size and stage of development
- Pigmentation increases with development along finfold margin (~7 spots along dorsal and 5 spots along anal in flexion larvae)

Diagnostic characters

- Postanal band pattern in preflexion larvae (three bands)
 - High myomere count (57-65)
- Distinguished from similar larvae with high myomere count by
- Presence of three postanal pigment bands and median finfold pigment
 - Reinhardtius hippoglossoides* (p. 622): Lightly pigmented and without bands
 - Glyptocephalus* spp. (p. 584, 586): Three bands (more laterally intense) in *G. stelleri*; in *G. zachirus*, caudal band, no finfold pigment
 - Unidentified ophidiid larvae (p. 213): More myomeres (>75), less finfold pigment, and less pronounced loop in gut



Figures A–C, E, Richardson 1981b; D, Ahlstrom et al. 1984a.

MERISTICS

Vertebrae	Total: 41-X-45
	Precaudal: 11-11-11
	Caudal: 30-32-34
Branchiostegal rays	7-X-8
Caudal fin	X, 9+8, X
	Total rays=19 ^a
Pelvic fin	Thoracic
	R: 6-6-6
Dorsal fin	R: 82-97-103
Pectoral fin	R: 13-13-13
Anal fin	R: 67-73-79
Gill rakers	U: X-X-X L: 15-X-17

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi-, meso-, and bathybenthal, 0-550 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Dec-Apr ^b Area: Off continental slope, 274-366 m ^c Mode: Migration: To deep water for spawning ^c
Fecundity	Range/function: 400,000-1,200,000 ^d
Age at first maturity	3-8 yr (females) ^c 4-9 yr (males) ^c
Longevity	25 yr (females) ^c 21 yr (males) ^c

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.21-1.25 mm
No. of oil globules	None
Oil globule diameter	
Yolk	
Envelope	
Hatch size	2.8 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE^e

Preanal length	<50% SL
Length at flexion	
Length at transformation	
Sequence of fin development	
Pigment	<ul style="list-style-type: none"> • Unpigmented at hatching • Preflexion <ul style="list-style-type: none"> —Midbody patch and urostyle pigment —Melanophores on body over anus —Posteriorly on gut and on head • Flexion and early postflexion larvae are undescribed

Diagnostic characters

- Distinguished from other pleuronectids by
- Presence of two postanal pigment bands including pigment on urostyle
 - Presence of preopercular spines

^a 15 branched rays.


^b Alderdice and Forrester 1971

^c Frey 1971

^d Forrester 1969

^e Larvae are rare. Since most routine ichthyoplankton surveys sample only to 200 m, the absence of small larvae in field collections indicates larvae may occur below 200 m.

Ref: Ahlstrom et al. 1984a, Alderdice and Forrester 1971.



Figures A–B, Alderdice and Forrester 1971; C, Ahlstrom et al. 1984a.

MERISTICS

Vertebrae	Total: 52-X-60	
	Precaudal: 9-9-9	
	Caudal: 43-43-43	
Branchiostegal rays	7-7-7	
Caudal fin		
Pelvic fin	Thoracic	
	R: 6-6-6	
Dorsal fin	R: 83-X-97	
Pectoral fin	R: 10-X-12	
Anal fin	R: 72-X-80	
Gill rakers	U: X-X-X	L: 7-X-10

LIFE HISTORY

Range	Bering Sea, 54-66°N
Ecology	Epi- and mesobenthal
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.20-1.61 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Thick, sometimes striated
Hatch size	4.1-5.2 mm SL
Incubation time/temp.	
Pigment	
	• Yolksac
	• Late-stage embryo

Diagnostic characters**LARVAE**


Preanal length	<50% SL
Length at flexion	15-17 mm SL
Length at transformation	50-60 mm SL; ^a eye migration begins at 35 mm SL
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals
Pigment	
	• Finfold (dorsal and anal)
	• Three postanal bands

Diagnostic characters

- Distinguished from other pleuronectids by
- Presence of three postanal pigment bands
 - High myomere count (52-60)
 - Presence of preopercular spines
- Distinguished from *G. zachirus* by
- Fewer myomeres (52-60)
 - Pigment pattern
 - Presence of finfold pigment
 - Difference in band pattern (fewer bands)

^a Ahlstrom et al. 1984a: 19-48 mm SL.

Ref: Ahlstrom et al. 1984a, Okiyama 1963, Okiyama and Takahashi 1976, Pertseva-Ostroumova 1961.



Figures A–D, Pertseva-Ostroumova 1961 (B–C, reversed); E, Okiyama and Takahashi 1976.

MERISTICS

Vertebrae	Total: 63-64-66 Precaudal: 12-13-14 Caudal: 50-51-52
Branchiostegal rays	7-7-7
Caudal fin	4, 8+7, 4 Total rays=22-23 ^a
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 87-102-110
Pectoral fin	R: 11-X-13
Anal fin	R: 78-85-93
Gill rakers	U: 4-4-4 L: 5-X-8

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathybenthal, 0-850 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Jan-June ^b Area: Mode: Migration:
Fecundity	Range/function: 3900-238,000/ $F=0.00000053797 \times L^{4.22667}$, L=TL mm ^b
Age at first maturity	5-9 yr (females) ^b 3-5 yr (males) ^b
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.80-2.2 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Surface may be irregular or appear bumpy
Hatch size	~5 mm SL
Incubation time/temp.	
Pigment	<ul style="list-style-type: none"> • Pigment on yolksac • Late stage (three bands + caudal)

Diagnostic characters

Distinguished from *Lyopsetta exilis* (p. 606) and *Microstomus pacificus* (p. 608) by

- Size
- Pigment on yolksac, late-stage embryo
- Wide perivitelline space
- Size of embryo (coiling in *G. zachirus*)

LARVAE

Preanal length	<50% SL
Length at flexion	15.3-24.0 mm SL
Length at transformation	49-59 mm SL ^c
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals
Pigment	<ul style="list-style-type: none"> • Postanal body <ul style="list-style-type: none"> —Three bands —Caudal band (above, below, and in finfold) —Melanophores between bands • Head, lower jaw, gut • With development, caudal pigment becomes restricted to hypural area

Diagnostic characters

- See *G. stelleri* (p. 584)
- Distinguished from other pleuronectids by
- Presence of four postanal pigment bands
 - High myomere count (63-66)
 - Presence of preopercular spines

See *Embassichthys bathybius* (p. 580) and *Microstomus pacificus* (p. 608) for similar larvae

^a13 branched rays.

^bHosie and Horton 1977

^cAccording to Ahlstrom et al. (1984a), transformation may occur as late as 72 mm SL.




Figure A, Ahlstrom and Moser 1980; B, NWAFC original (B. Vinter); C–D, Ahlstrom et al. 1984a (after Ahlstrom and Moser 1975).

MERISTICS

Vertebrae	Total: 43-44-47 Precaudal: 12-13-13 Caudal: 32-X-35
Branchiostegal rays	7-7-7
Caudal fin	X, 7+7, X Total rays=18 ^a
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 72-79-90
Pectoral fin	R: 10-X-12
Anal fin	R: 55-62-71
Gill rakers	U: X-X-X L: 14-20-26

LIFE HISTORY

Range	Cent. California, 34-38°N, to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathybenthal, 0-875 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Feb-July ^b Area: 50-305 m ^b Mode: Migration:
Fecundity	Range/function: 70,000- 600,000 ^c
Age at first maturity	1-2 yr ^c
Longevity	21 yr ^b

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	2.75-3.75 mm (2.9-3.5 mm)
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous, colorless
Envelope	Smooth
Hatch size	5.3-6.9 mm SL ^d
Incubation time/temp.	
Pigment	
• Yolksac	
• Late-stage embryo:	On body and in dorsal and ventral finfold

Diagnostic characters

- Wide perivitelline space
- Body and finfold pigment

LARVAE

Preanal length	<50% SL
Length at flexion	9-10 mm SL
Length at transformation	Probably gradual, beginning between 18 and 21 mm SL
Sequence of fin development	Caudal, dorsal, and anal; pelvics; pectorals
Pigment	
• Three postanal bands and along ventral midline	
• Around urostyle	
• Finfold	
• With development becoming less pigmented	

Diagnostic characters

- Distinguished from other pleuronectids with three postanal pigment bands by
- Presence of hypaxial, fin, and urostyle pigment
 - Low myomere count (43-47)

^a 12 branched rays.^b Salveson 1976^c Miller 1969^d Pertseva-Ostroumova 1961




Figure A, Pertseva-Ostroumova 1961; B–E, NWAFC originals (B. Vinter).

MERISTICS

Vertebrae	Total: 44-44-44 Precaudal: 12-12-12 Caudal: 32-32-32
Branchiostegal rays	7-7-7
Caudal fin	
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 66-74-80
Pectoral fin	R: 10-10-10
Anal fin	R: 53-59-64
Gill rakers	U: 1-1-1 L: 10-12-17

LIFE HISTORY

Range	Aleutian Is., 51-55°N, to Chukchi Sea, north of 66°N
Ecology	Epi- and mesobenthal, 18-425 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Apr-June ^a Area: Shallow gulfs and bays (50-150 m) ^a Mode: Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	2.04-2.69 mm; occasionally up to 2.90 mm
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	Smooth, thin
Hatch size	4 mm SL
Incubation time/temp.	
Pigment	
	• Yolksac
	• Late-stage embryo: On body

Diagnostic characters

- Wide perivitelline space
- Smaller in diameter than *H. elassodon*

LARVAE

Preanal length	<50% SL
Length at flexion	~11 mm SL
Length at transformation	>28.6 mm SL ^b
Sequence of fin development	
Pigment	
	• Three postanal bands extending into finfold


Diagnostic characters

- See *H. elassodon* (p. 588)

^aPertseva-Ostroumova 1961^bAhlstrom et al. 1984a

Ref: Ahlstrom et al. 1984a, Pertseva-Ostroumova 1961.


A




2.04–2.69 mm

B


Three postanal bands



10.8 mm



C



31.0 mm SL

Figures A–C (B insert, ventral view of head), Pertseva-Ostroumova 1961.

MERISTICS

Vertebrae	Total: 49-50-51 Precaudal: 16-16-16 Caudal: 35-35-35
Branchiostegal rays	7-7-7
Caudal fin	Total rays=19
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 89-99-109
Pectoral fin	R: 19-19-19
Anal fin	R: 64-75-81
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Chukchi Sea, north of 66°N
Ecology	Epi-, meso-, and bathybenthal, 6-1110 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Nov-Mar ^a Area: 180-550 m ^a Mode: Migration: To deepwater spawning banks in Gulf of Alaska ^b
Fecundity	Range/function: 200,000-4 million ^c
Age at first maturity	7-20 yr (females) ^a 5-20 yr (males) ^a
Longevity	42 yr (females) ^d 27 yr (males) ^d

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	2.9-3.8 mm
No. of oil globules	
Oil globule diameter	
Yolk	Homogeneous, dense, yellow (after preservation)
Envelope	Shallow honeycomb pattern (not always visible on preserved specimens but slight surface irregularities are easily discernible)
Hatch size	7.8-8.5 mm SL

Incubation time/temp.

Pigment

- Embryo unpigmented at hatching

Diagnostic characters

- Large size (>3.0 mm)
- Lack of pigment

LARVAE

Preanal length	<50% SL
Length at flexion	13.6-17.8 mm SL
Length at transformation	14.7-24.1 mm SL ^e
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals

Pigment

- Lack of pigment in yolk sac larvae
- Series of melanophores along dorsal surface of notochord and along ventral midline
- Median finfolds pigmented along edge

Diagnostic characters

- Distinguished from other pleuronectids by
- Myomeres (49-51)
 - Pigment: Lack of bands, and presence of pigment on ventral midline and along edges of median finfolds

Distinguished from *Reinhardtius hippoglossoides* (both have large yolk sac larvae) by

- See *R. hippoglossoides* (p. 622)

^aSt. Pierre 1984


^bThompson and Van Cleve 1936

^cSchmitt and Skud 1978

^dWebber and Alton 1976

^eBecome juveniles at 28 mm SL.

Ref: Ahlstrom et al. 1984a, Pertseva-Ostroumova 1961, Thompson and Van Cleve 1936.



Figures A–B, Pertseva-Ostroumova 1961 (after Thompson and Van Cleve 1939); C–D, NWAFC originals (B. Vinter); E, Ahlstrom et al. 1984a.

MERISTICS

Vertebrae	Total: 41-42-42 Precaudal: 10-10-11 Caudal: X-X-X
Branchiostegal rays	7-X-8
Caudal fin	X, 7+6, X Total rays=18 ^a
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 78-86-90
Pectoral fin	R: 11-X-13
Anal fin	R: 58-66-69
Gill rakers	U: X-X-X L: 7-X-8

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthal, 20-425 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Feb-Apr ^b Area: Coastal waters ^b Mode: Schools ^b Migration:
Fecundity	Range/function: 350,000- 650,000 ^c
Age at first maturity	3 yr (females) ^b 2 yr (males) ^b
Longevity	11 yr (females) ^d 10 yr (males) ^d

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	0.84-1.00 mm; occasionally up to 1.10 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Striated
Hatch size	2.7-2.9 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

Very difficult to distinguish from the three other 1.0-mm pleuronectid eggs in the area:
Parophrys vetulus (p. 610)
Platichthys stellatus (p. 612)
Psettichthys melanostictus (p. 620)

LARVAE

Preanal length	<50% SL
Length at flexion	9-10 to 14 mm SL
Length at transformation	15->21 mm SL
Sequence of fin development	Caudal, dorsal, and anal; pelvics; pectorals
Pigment	<ul style="list-style-type: none"> • Three postanal bands at 50, 67, 90% SL • Melanophores extend ventrally on gut and along posterior portion of abdominal cavity

Diagnostic characters


- Three pigment bands
- Low myomere count (41-42)

^a 11 branched rays.

^b Smith 1936

^c Forrester 1969

^d Hart 1973



Figures A–F, Richardson et al. 1980 (B, reared).

MERISTICS

Vertebrae	Total: 39-40-42 Precaudal: 10-11-12 Caudal: 28-30-31
Branchiostegal rays	7-7-7
Caudal fin	Total rays=18-19 ^a
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 65-76-84
Pectoral fin	R: 8-11-13
Anal fin	R: 50-58-65
Gill rakers	U: 3-3-3 L: 5-X-8

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathybenthal, 0-579 m
ELH pattern	Oviparous; demersal, adhesive eggs; pelagic larvae
Spawning	Season: Feb-Apr ^b Area: Mode: Migration:
Fecundity	Range/function: 400,000-1,300,000 (Brit. Col.); ^c 150,000-400,000 (Bering Sea); ^d 80,000-920,000 (western Pacific)/ $F=0.0004891 \times L^{3.720}$, $L=BL \text{ cm}^e$
Age at first maturity	3-4 yr (females) ^b 2 yr (males) ^b
Longevity	15 yr (females) ^f 22 yr (males) ^f

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	0.87-1.00 mm; 1.02-1.09 mm ^g
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Thick, elastic, bright yellow or orange
Hatch size	5 mm SL; 3.6-4.0 mm SL; ^g 3.4-3.8 mm SL ^h
Incubation time/temp.	9-12 d/6.5-8.0°C
Pigment	

Diagnostic characters

- Demersal egg

LARVAE

Preanal length	<50% SL
Length at flexion	8.4-9.9 mm SL
Length at transformation	~20 mm SL; >17.7 mm SL ^h
Sequence of fin development	Caudal, dorsal, and anal; pelvics; pectorals
Pigment	

- Dorsally on peritoneum, ventrolateral gut pigment becoming posterolateral with development
- Postanal ventral melanophores ~3/4 length of body
- Two dorsolateral patches, the posterior one becoming a band meeting the ventral surface
- Anal finfold lightly pigmented on posterior half

Diagnostic characters

- Distinguished from other pleuronectids by
- Presence of one band posteriorly on body, not as conspicuous as most
 - Size at transformation (20 mm SL)
 - Advanced stage of development at hatching

See *Lepidopsetta* 2 (p. 599) and *Psettichthys melanostictus* (p. 620)

^a 12 branched rays.

^b Smith 1936

^c Forrester 1969

^d Fadeev 1965

^e Shvetsov 1979

^f Hart 1973

^g Pertseva-Ostroumova 1961

^h Ahlstrom et al. 1984a




Figure A, Pertseva-Ostroumova 1961; B–E, NWAFC originals (B. Vinter).

Lepidopsetta bilineata/*Lepidopsetta 2*/*Psettichthys melanostictus*

Two readily distinguishable types of *Lepidopsetta* larvae are collected in our study area which we designate *Lepidopsetta bilineata* and *Lepidopsetta 2*. Wilimovsky et al. (1967) had previously indicated that there are two subspecies of *Lepidopsetta* in the Northeast Pacific based on adult specimens (*L. b. bilineata* and *L. b. peracuata*). *Lepidopsetta 2* larvae are very similar to larvae of *Psettichthys melanostictus* at certain stages of development. We present *Lepidopsetta 2* as a separate series, since these larvae are distinct from those of *L. bilineata* and *P. melanostictus*. The following pigment patterns and morphological characters separate the three kinds of larvae.

Psettichthys melanostictus (see also p. 620)

Range from California to Alaska, but the center of distribution appears to be off California, Oregon, and Washington.

Pigment

- Postanal ventral midline melanophores: Usually restricted to 3-4 large spots
- Tail pigment less prominent than on *Lepidopsetta 2*
- Pigment on upper and lower jaw
- Isthmus pigment heavier than on others
- First spot along dorsal midline more anterior (over anus)

Morphology

- Gut coiled, shape of posterior gut
- Deeper body (wider finfold)

Lepidopsetta 2 (see p. 600, 601)

Range from Puget Sound to Alaska, and center of distribution appears to be off Southeast Alaska and in the Gulf of Alaska.

Pigment

- Postanal ventral midline melanophores: A series of small melanophores extending from the gut to the last myomere; sometimes a larger spot occurs about midbody (with larvae 6.3-9.7 mm SL, there may be more spots than at other sizes)
- First spot along dorsal midline behind anus
- Other: Tail pigment more intense, but mouth and isthmus pigment less intense than on *Psettichthys*

Morphology

- Eye relatively larger than on *Psettichthys* at similar ontogenetic stages
- Gut shape simple

Lepidopsetta bilineata (see also p. 596)

Range from southern California to the Bering Sea.

Pigment

- Postanal ventral midline melanophores: A series of small melanophores extending from the gut to about 2/3 body
- Only two spots along dorsal midline; anterior spot at midbody, posterior spot forming a band at about myomere 30
- No heavy pigment along edges of median finfold
- Tail pigment less prominent than on others
- Pigment along hypural edge

Morphology

- Gut shape simple

Finfold pigment may or may not be an additional criterion for separating *Lepidopsetta 2* and *Psettichthys*. Many *Psettichthys* specimens in our collection appear to have more prominent finfold pigment that is retained longer during development; however, many of these are laboratory-reared specimens. More wild-caught *Psettichthys* specimens are needed to verify whether they all have intense finfold pigment.



Yolksac larvae are more difficult to separate. *Psettichthys* larvae appear to possess upper and lower jaw pigment, smaller, more concentrated melanophores in the finfold and on the urostyle, and the last postanal band has a ventral stripe only. *Lepidopsetta 2* yolksac larvae appear to have only lower jaw pigment, larger, fewer, and evenly spaced melanophores in the finfold and urostyle, and the last postanal band has a dorsal and ventral stripe.

Literature: Specimens illustrated by Ahlstrom et al. (1984a) are both *Psettichthys* based primarily on the presence of the coiled gut and the 2-3 large ventral melanophores (Fig. 351C, p. 662, and Fig. 352E, p. 663). Hickman (1959) illustrated six *Psettichthys* larvae from specimens he reared from eggs (his Figs. 1 and 2) and from specimens he collected in Puget Sound (his Figs. 3-6). Hickman's Figures 3 and 4 appear to resemble *Lepidopsetta 2* larvae, while his other figures appear to resemble *Psettichthys*.



PLEURONECTIDAE

PSETTICHTHYS / LEPIDOPSETTA



Psettichthys melanostictus



Lepidopsetta 2




Lepidopsetta bilineata



Figures A–I, NWAFC originals (B. Vinter; A, D, G, include a detailed enlargement of tail).


PSETTICHTHYS / LEPIDOPSETTA

C




13.9 mm SL

F



16.4 mm SL

I



16.3 mm SL

MERISTICS

Vertebrae	Total: 40-40-41 Precaudal: 10-X-12 Caudal: 28-X-30
Branchiostegal rays	7-7-7
Caudal fin	Total rays = 18 ^a
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 61-69-77
Pectoral fin	R: 10-X-12
Anal fin	R: 49-52-58
Gill rakers	U: X-X-X L: 8-X-10

LIFE HISTORY

Range	Brit. Col., 48°30'-55°N, to Chukchi Sea, north of 66°N
Ecology	Epi-, meso-, and bathy-benthal, 10-600 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Summer ^b Area: Inner shelf region ^b Mode: Mass ^b Migration: To shallower water of shelf ^b
Fecundity	Range/function: 1,300,000-3,300,000/ $F=0.0747565 \times L^{2.86517}$ (F in 1000 eggs), L=TL cm ^b
Age at first maturity	6-7 yr (females) ^b 4-5 yr (males) ^b
Longevity	19 yr (females) ^b 17 yr (males) ^b

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	0.76-0.85 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous, yellowish
Envelope	Clear, smooth
Hatch size	2.25-2.80 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

- Size

LARVAE

Preanal length	<50% SL
Length at flexion	~7 mm SL; 7.5-9.5 mm SL ^c
Length at transformation	15-17 mm SL; may begin at 10 mm SL ^c
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals
Pigment	

- Mediolateral, along notochord
- Ventrolateral, along hypaxial myomeres
- Anal finfold

Diagnostic characters

- Distinguished from other pleuronectids without postanal pigment bands by
- Ventrolateral pigment along hypaxial myomeres
 - Size at transformation (15-17 mm SL)
 - Finfold pigment restricted to anal fin
 - Urostyle unpigmented

L. proboscidea and *L. sakhalinensis*^d eggs and larvae are incompletely known. The following information may aid in identification.

	<i>L. proboscidea</i>	<i>L. sakhalinensis</i>
Egg diameter	0.72-0.87 mm	
Total vertebrae	38-40	
Precaudal	11	
Caudal	27-29	
Dorsal fin rays	62-69	68-76
Pectoral fin rays	12	
Anal fin rays	46-50	53-59
Range	Bering Sea-Chukchi Sea	
Spawning season	Spring-summer ^e	

^a 12 branched rays.


^b Salvesson and Alton 1976b

^c Ahlstrom et al. 1984a

^d According to Allen and Smith (1988), this species occurs in the Bering Sea.

^e Pertseva-Ostroumova 1961

Ref: Ahlstrom et al. 1984a, Pertseva-Ostroumova 1961, Schmidt 1950.



Figures A–F (C, reversed), Pertseva-Ostroumova 1961.

MERISTICS

Vertebrae	Total: 37-X-41 Precaudal: 11-X-13 Caudal: 26-26-26
Branchiostegal rays	7-X-8
Caudal fin	Total rays=18
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 48-X-64
Pectoral fin	R: 8-X-12
Anal fin	R: 33-X-46
Gill rakers	U: 2-X-4 L: 7-X-8

LIFE HISTORY

Range	Bering Sea, 54-66°N, to Arctic, not specific
Ecology	Nearshore shelf demersal
ELH pattern	Oviparous, demersal eggs, pelagic larvae
Spawning	Season: Jan-Mar ^a Area: Shallow water, 5-10 m ^a Mode: Migration:
Fecundity	Range/function: 31,000-230,000 ^a
Age at first maturity	2+ yr (usually 4-5) ^b
Longevity	>9 yr ^b

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.2-1.7 mm (1.54-1.70 mm)
No. of oil globules	None
Oil globule diameter	
Yolk	Clear, homogeneous
Envelope	Thin, smooth
Hatch size	3.7 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

LARVAE^c

Preanal length	<50% SL
Length at flexion	
Length at transformation	
Sequence of fin development	
Pigment	<ul style="list-style-type: none"> • Newly hatched larvae heavily pigmented, but with development pigment appears to be restricted to postanal body above and below notochord

Diagnostic characters


- Distinguished from other pleuronectids without pigment bands by
- No pigment in finfolds or along hypaxial myomeres
 - Myomere count (37-41)

^aPertseva-Ostroumova 1961

^bAndriashev 1954

^cPreflexion larvae only.

Ref: Pertseva-Ostroumova 1961.



Figures A–C, Pertseva-Ostroumova 1961 (figures reversed).

MERISTICS

Vertebrae	Total: 43-45-47 Precaudal: 11-12-13 Caudal: 32-33-35
Branchiostegal rays	7-7-7
Caudal fin	X, 8+7, X Total rays = 19 ^a
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 72-77-88
Pectoral fin	R: 10-10-10
Anal fin	R: 57-60-66
Gill rakers	U: 2-X-3 L: 9-X-11

LIFE HISTORY

Range	South of southern California to Gulf of Alaska, 54-60°N
Ecology	Epi-, meso-, and bathybenthal, 25-800 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Feb; ^b Apr ^c Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	3-5 yr (females) ^c 2-3 yr (males) ^c
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.47-1.71 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Surface may be irregular and appear bumpy
Hatch size	~5.6 mm SL
Incubation time/temp.	
Pigment	
	• Yolksac
	• Late-stage embryo: Distinct caudal pigment visible

Diagnostic characters

- Caudal pigment
- Smallest size range of three similar-looking pleuronectid eggs in the 1.5-2.5 mm size range; see *Glyptocephalus zachirus* (p. 586) and *Microstomus pacificus* (p. 608)

LARVAE

Preanal length	<50% SL
Length at flexion	9.0-10.9 mm SL
Length at transformation	15.7-24.7 mm SL
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals

Pigment

- Early and midflexion larvae
 - Dorsal and ventral midline melanophores around tail and in finfold; with development, these melanophores move laterally on body along myoseptal lines
 - Finfold pigment increases to cover anal finfold and posterior half of dorsal finfold
- Gut pigment increases ventrally with development

Diagnostic characters


- Distinguished from other pleuronectids without pigment bands by
- Pigment pattern: Continuous dorsal and ventral midline
 - In preflexion larvae, the anterior melanophores of the dorsal midline series extend laterally
 - Finfold pigment

^a 12-13 branched rays.

^b Hart 1973

^c Smith 1936

Ref: Ahlstrom and Moser 1975, Ahlstrom et al. 1984a.



Figures A–E, Ahlstrom and Moser 1975.

MERISTICS

Vertebrae	Total: 50-52-55 Precaudal: 11-12-13 Caudal: 38-40-41
Branchiostegal rays	7-7-7
Caudal fin	X, 9+8, X Total rays=21 ^a
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 94-105-116
Pectoral fin	R: 8-X-12
Anal fin	R: 80-87-96
Gill rakers	U: 5-X-8 L: 8-X-11

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathybenthal, 9-1189 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Nov-Aug ^b Area: 80-550 m ^b Mode: Migration: Offshore for spawning ^c
Fecundity	Range/function: 37,188 ^d -260,000 ^e
Age at first maturity	5 yr ^f
Longevity	45 yr ^g

^a13-16 branched rays.
^bHirschberger and Smith 1983
^cWestrheim and Morgan 1962
^dHagerman 1952
^eHarry 1959
^fFrey 1971
^gChilton and Beamish 1982

Ref: Ahlstrom and Moser 1975, Ahlstrom et al. 1984a, Richardson 1981b.

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	2.05-2.68 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Smooth, slightly thick
Hatch size	~6 mm SL
Incubation time/temp.	27 d/10°C
Pigment	
• Yolksac	
• Late-stage embryo: Caudal pigment not as pronounced as <i>Lyopsetta exilis</i> (p. 606)	

Diagnostic characters

- Largest egg of three similar-looking pleuronectid eggs; see *L. exilis* and *Glyptocephalus zachirus* (p. 586)

LARVAE

Preanal length	<50% SL
Length at flexion	10-15 mm SL
Length at transformation	Eye migration begins at 20 mm SL; larvae remain pelagic >45 mm SL
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals
Pigment	

- Lower jaw; dorsally and ventrally on gut, with development becoming restricted to ventral surface (isthmus to anus)
- Preflexion
 - Three pigment patches in addition to tail and finfold
 - Anterior patch restricted to ventral area; two posterior patches have dorsal and ventral components extending into finfold
 - Around urostyle and in finfold
- Flexion: Pigment patches occur above notochord and along distal ends of finfold
- Postflexion: Pigment patches along dorsal and anal pterygiophores

Diagnostic characters

- Distinguished from other pleuronectids with 3-4 pigment bands by
- Pigment bands extending into finfold
 - Morphology: Early larvae long and slender, becoming deep-bodied during development
 - Otic spines
 - Myomeres (50-55)

See also *G. zachirus* (p. 586) and *Hippoglossoides elassodon* (p. 588)




Figure A, Ahlstrom and Moser 1975; B, Ahlstrom et al. 1984a (after Ahlstrom and Moser 1975); C–D, NWAFC originals (B. Vinter).

MERISTICS

Vertebrae	Total: 42-44-47 Precaudal: 10-11-12 Caudal: 31-33-34
Branchiostegal rays	7-X-8
Caudal fin	X, 7+7, X Total rays=18 ^a
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 72-80-82
Pectoral fin	R: 10-X-12
Anal fin	R: 54-60-70
Gill rakers	U: 4-X-6 L: 10-X-13

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathybenthal, intertidal to 550 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Oct-May (California); ^b Jan-Apr (Puget Sound) ^c Area: Demersal ^d Mode: Migration: Southern migration from feeding grounds to sheltered water in channels or bights ^d
Fecundity	Range/function: 150,000-1,950,000 ^e
Age at first maturity	3-4 yr (females) ^c 2 yr (males) ^c
Longevity	17 yr ^f

^a 12 branched rays.

^b Jow 1969

^c Smith 1936

^d Ketchen 1956

^e Ketchen 1947

^f Frey 1971

^g Ahlstrom et al. 1984a

Ref: Budd 1940, Ahlstrom and Moser 1975, Ahlstrom et al. 1984a.

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	0.80-1.05 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Thin, smooth, transparent
Hatch size	2.3-2.8 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

- See *Isopsetta isolepis* (p. 594)

LARVAE


Preanal length	43-45% SL
Length at flexion	~10 mm SL; 8.8-10.5 mm SL ^g
Length at transformation	Eye migration prior to 17.5 mm SL
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals
Pigment	

- Initially, anteroventrally on gut; with development, extending posteriorly; in postflexion, a line of pigment forms along the posterior hindgut and anus
- Dorsal and ventral midline: Ventral beginning prior to anus (~ myomere 5) and dorsal at ~ myomere 15 (pigment is variable); dorsal midline becoming less prominent with development as internal pigment develops above notochord through flexion stage
- Myoseptal pigment develops along hypaxial myomeres in postflexion larvae
- Finfold pigment restricted to anal finfold

Diagnostic characters

- Distinguished from other pleuronectids without pigment bands by
- Dorsal and ventral midline pigment, hypaxial pigment
 - Finfold pigment restricted to anal finfold

Note: The number of dorsal midline melanophores in preflexion larvae is highly variable.



Figures A–B, D, NWAFC originals (B. Vinter); C, Ahlstrom et al. 1984a (after Ahlstrom and Moser 1975).

MERISTICS

Vertebrae	Total: 35-36-38 Precaudal: 10-11-12 Caudal: 24-25-26
Branchiostegal rays	7-7-7
Caudal fin	Total rays=18
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 52-59-66
Pectoral fin	R: 9-10-10
Anal fin	R: 38-42-47
Gill rakers	U: 3-3-3 L: 6-X-8

LIFE HISTORY

Range	S. California, 32-34°N, to Arctic, not specific
Ecology	Epi- and mesobenthic, freshwater (upstream) to 375 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Nov-Feb (California); ^a Feb-Apr (Puget Sound); ^b May-June (Bering Sea) ^c Area: Shallow water ^a Migration:
Fecundity	Range/function: 900,000 ^c -11 million ^a
Age at first maturity	3-4 yr (females) ^a
Longevity	21 yr ^d

^aOrcutt 1950^bHart 1973^cFadeev 1965^dWolotira et al. 1977^eAhlstrom et al. 1984a

Ref: Ahlstrom et al. 1984a, Orcutt 1950, Pertseva-Ostroumova 1961, Yusa 1957.

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	0.88-1.28 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Striated, yellow
Hatch size	1.9-2.1 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

- See *Isopsetta isolepis* (p. 594)

LARVAE


Preanal length	<50% SL
Length at flexion	~7 mm SL; 5.5-6.0 mm SL ^e
Length at transformation	10.5 mm SL (may be slightly smaller, 8-9 mm SL)
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals

Pigment

- Lightly scattered over head and posteroventrally along gut
- Initially, in preflexion larvae, postanal pigment scattered along anal finfold and along posterior 1/3 of body
- Around urostyle
- Internally above notochord, becoming less concentrated with development
- Pigment in dorsal fin disappears with development
- Postflexion larvae lightly pigmented with lateral patches

Diagnostic characters

- Distinguished from other pleuronectids without pigment bands by
- Lack of dorsal midline pigment
 - Small size at transformation (<10 mm SL)



Figures A–B, Orcutt 1950 (B, reversed); C–F, NWAFC originals (B. Vinter).

MERISTICS

Vertebrae	Total: 41-X-42 Precaudal: 18-18-19 Caudal: X-X-X
Branchiostegal rays	7-X-8
Caudal fin	
Pelvic fin	Thoracic R: 6-6-6
Dorsal fin	R: 62-69-71
Pectoral fin	R: 11-11-11
Anal fin	R: 51-51-56
Gill rakers	U: X-X-X L: 7-7-7

LIFE HISTORY

Range	Gulf of Alaska, 54-60°N, to Chukchi Sea, north of 66°N
Ecology	Epi- and mesobenthal, 6-475 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Spring ^a Area: Mode: Migration: Inshore ^a
Fecundity	Range/function:
Age at first maturity	4 yr ^a
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.67-2.21 mm (1.7-1.9 mm)
No. of oil globules	None
Oil globule diameter	
Yolk	
Envelope	Thick wavy surface (color: bronze hue)
Hatch size	~5.85 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	<50% SL
Length at flexion	~8-10 mm SL
Length at transformation	Beginning by 10.7 mm SL
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals

Pigment


- Isthmus, posteroventrally on gut
- Initially, in preflexion larvae, dorsal spots over posterior half of body that become less prominent with development
- Ventral midline melanophores extend onto finfold and on caudal region
- Internal row above notochord throughout development

Diagnostic characters

- Distinguished from other pleuronectids without pigment bands by
- Pigment pattern
 - Urostyle unpigmented
 - No slash-like pigment along hypaxial myomeres
 - Finfold pigment mainly restricted to anal finfold
 - Size at transformation ~10 mm SL

^aPertseva-Ostroumova 1961

Ref: Ahlstrom et al. 1984a.



Figures A–C, F, Pertseva-Ostroumova 1961 (B–C, reversed); D–E, NWAFC originals (B. Vinter).

MERISTICS

Vertebrae	Total: 37-38-39
	Precaudal: 12-13-13
	Caudal: 24-25-26
Branchiostegal rays	7-X-8
Caudal fin	Total rays=19
Pelvic fin	Thoracic
	R: 6-6-6
Dorsal fin	R: 65-72-78
Pectoral fin	R: 9-11-12
Anal fin	R: 46-50-56
Gill rakers	U: 3-X-4 L: 8-X-11

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59°N
Ecology	Epi- and mesobenthal, 0-350 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Mar-Aug (California) ^a Area: Mode: Migration: Range/function:
Fecundity	
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION

EGGS

Diameter	1.20-1.56 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Sculptured with polygonal pattern
Hatch size	3.9 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

- Egg diameter (1.20-1.56 mm)
- Polygonal sculpturing on envelope surface

LARVAE

Preanal length	<50% SL
Length at flexion	6.2-8.5 mm NL
Length at transformation	8.2->11.4 mm SL
Sequence of fin development	Caudal slightly before dorsal and anal, pelvics, pectorals

Pigment

- Preflexion larvae: Opposing pigment clusters on dorsal and ventral finfolds, increasing with development
- Small melanophores covering all but posterior 1/4 of body

Diagnostic characters (see Table 3)

- Distinguished from other pleuronectids without pigment bands by
- Heavy pigment pattern
- Distinguished from *P. decurrens* by
- Pigment pattern: *P. coenosus* has less finfold pigment than *P. decurrens*
 - Precaudal vertebrae 12-13, whereas *P. decurrens* usually has 14-15
 - Lack of pterotic spines

Note: Preflexion larvae occasionally have more snout and lower jaw pigment than shown on figure.

^aBudd 1940

Ref: Ahlstrom et al. 1984a, Sumida et al. 1979.




Figure A, Budd 1940; B–E, Sumida et al. 1979.

MERISTICS

Vertebrae	Total: 38-39-41 Precaudal: 13-14-15 Caudal: 24-25-26
Branchiostegal rays	7-X-8
Caudal fin	2, 7+7, 2 Total rays = 19 ^a
Pelvic fin	Thoracic R: 4-6-7
Dorsal fin	R: 67-75-81
Pectoral fin	R: 9-12-14
Anal fin	R: 45-50-55
Gill rakers	U: 3-X-4 L: 6-X-9

LIFE HISTORY

Range	South of southern California to Bering Sea, 54-66°N
Ecology	Epi-, meso-, and bathybenthal, 8-532 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	1.84-2.08 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Sculptured with polygonal pattern
Hatch size	4.9-5.5 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

- Egg diameter (1.84-2.08 mm)
- Polygonal sculpturing on envelope surface

LARVAE

Preanal length	~50% SL
Length at flexion	7.8-11.0 mm NL
Length at transformation	10.5->21.0 mm SL ^b
Sequence of fin development	Caudal slightly before dorsal and anal, pelvics, pectorals

Pigment

- Nearly entire body and finfolds pigmented except posteriorly in preflexion larvae; banding pattern develops in postflexion larvae

Diagnostic characters (see Table 3)

- Distinguished from other pleuronectids without pigment bands by
- Heavy pigment pattern
 - Pterotic spines
- Distinguished from *P. coenosus* by
- See *P. coenosus* (p. 616)
 - In addition to having pterotic spines, more pigment, and more precaudal myomeres, *P. decurrens* is larger at various stages of development (see figures).

^a12 branched rays.^bEye migration by 10 mm SL, but not complete in some until >21 mm SL.




Figure A, Budd 1940; B-F, Sumida et al. 1979.

MERISTICS

Vertebrae	Total: 38-39-41
	Precaudal: 11-11-12
	Caudal: 28-28-30
Branchiostegal rays	7-7-7
Caudal fin	2, 7+7, 2
	Total rays = 18 ^a
Pelvic fin	Thoracic
	R: 6-6-6
Dorsal fin	R: 73-85-88
Pectoral fin	R: 10-X-12
Anal fin	R: 53-58-62
Gill rakers	U: 5-X-7 L: 14-X-18

LIFE HISTORY

Range	S. California, 32-34°N, to Bering Sea, 54-66°N
Ecology	Epi- and mesobenthic, 1-325 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Jan-Mar (Puget Sound); ^b July (British Columbia) ^c
	Area:
	Mode:
	Migration:
Fecundity	Range/function:
Age at first maturity	2-3 yr (females) ^d 2 yr (males) ^e
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	0.83-1.04 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Clear
Envelope	
Hatch size	<3 mm SL
Incubation time/temp.	
Pigment	
	• Yolksac: May appear as early as late middle stage

Diagnostic characters

See *Isopsetta isolepis* (p. 594); usually pigment on yolksac and in later stages on finfold

LARVAE (see discussion, p. 599)

Preanal length	<50% SL
Length at flexion	~8-10 mm SL
Length at transformation	>22.6 mm SL
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals
Pigment	
	• Postanal body
	—3-4 large spots along ventral midline
	—Three spots along dorsal midline; the first spot more anteriorly placed than the first ventral spot
	• Prominent pigment along edges of dorsal and ventral finfolds
	• Mouth, isthmus
	• Gut pigmented along ventral midline and posterior edge

Diagnostic characters

See *Lepidopsetta bilineata* (p. 596) and *Lepidopsetta 2* (p. 599)

Distinguished from other pleuronectids with pigment bands by





- Distinctive pigment along edges of dorsal and ventral finfolds and dorsal and ventral body midlines

^a 12 branched rays.

^b English 1961

^c Manzer 1947

^d Smith 1936



Figures A–D, NWAFC originals (B. Vinter).

MERISTICS

Vertebrae	Total: 61-63-64 Precaudal: 17-18-19 Caudal: 43-45-46
Branchiostegal rays	7-7-7
Caudal fin	
Pelvic fin	Thoracic R: 5-6-7
Dorsal fin	R: 83-97-105
Pectoral fin	R: 11-14-15
Anal fin	R: 63-72-79
Gill rakers	U: 3-X-6 L: 11-X-16

LIFE HISTORY

Range	South of southern California to Chukchi Sea, north of 66°N
Ecology	Epi-, meso-, and bathybenthal, 14-2000 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Fall ^a Area: Continental slope (>100 m) ^a Mode: Migration:
Fecundity	Range/function: 15,000-215,000 (Atlantic)/ $F=0.000063 \times L^{4.66}$, $L=FL \text{ cm}^a$
Age at first maturity	13 (females, Okhotsk Sea) ^a
Longevity	>23 yr ^a

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	4.0-4.5 mm
No. of oil globules	None
Oil globule diameter	
Yolk	Homogeneous
Envelope	Smooth, transparent
Hatch size	10-16 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters**LARVAE**

Preanal length	<50% SL
Length at flexion	25-27 mm SL
Length at transformation	45-65 mm SL
Sequence of fin development	Caudal, dorsal and anal, pelvics, pectorals
Pigment	


- Ventrally and posterolaterally on gut
- Light ventrolateral pigment along body from above gut, some dorsal, along peduncle; also in finfold above and below tail (~posterior 1/3)
- Increase in pigmentation in postflexion specimens

Diagnostic characters

- Newly hatched larvae large and unpigmented (except for eye)
- Distinguished from other pleuronectids with >60 myomeres by
- No band pattern and overall lightly pigmented
 - Glyptocephalus zachirus* (p. 586): Four postanal bands with more lateral intensity
 - Embassichthys bathybius* (p. 580): Three postanal bands
- Distinguished from *Hippoglossus stenolepis* (p. 592) (both have large unpigmented yolk sac larvae) by
- Depth of collection (>200 m)
 - High myomere count (61-64)
 - Hatch size (10-16 mm SL)

^aDunn and Sample 1976

Ref: Ahlstrom et al. 1984a, Jensen 1935.



Figures A–F, Jensen 1935 (A–D, reversed; specimens collected from West Greenland waters).

MERISTICS

Vertebrae	Total: 50-51-52 Precaudal: 9-9-9 Caudal: 41-X-43
Branchiostegal rays	6-6-6
Caudal fin	Total rays = 12 ^a
Pelvic fin	Thoracic; 4 rays on eyed side, absent on blind side
Dorsal fin	R: 95-X-106
Pectoral fin	Present during larval period but lacking in juvenile and adult stages
Anal fin	R: 77-X-90
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to Oregon, 42-46°N
Ecology	Nearshore shelf demersal, 1-201 m
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: June-Sept ^b Area: Mode: Migration:
Fecundity	Range/function:
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	0.71-0.78 mm
No. of oil globules	Multiple, 10-23
Oil globule diameter	
Yolk	Homogeneous
Envelope	Smooth, colored
Hatch size	1.9 mm SL
Incubation time/temp.	
Pigment	

Diagnostic characters

- Multiple oil globules

LARVAE

Preanal length	50% NL decreasing with development to 28% SL
Length at flexion	9.4-10.8 mm SL
Length at transformation	19.0-24.2 mm SL
Sequence of fin development	Anterior dorsal; caudal, dorsal, and anal; pelvics ^c (pectorals disappear at metamorphosis)

Pigment

- Small melanophores along dorsal midline, larger along ventral midline
- Single band posteriad on tail
- Large blotches at finfold margin, with development restricted to along the distal edges
- Head, gut, and swimbladder

Diagnostic characters

- Morphology: Large head and tapering body
- Gut mass trails posteriad
- Five exerted dorsal fin rays develop by 6.0 mm SL but they are no longer than other fin rays (in other species of *Symphurus*, elongate anterior dorsal fin rays persist)
- Pelvic fin on blind side begins disappearing at 18.0-22.0 mm SL^c

^aAll rays on hypurals, all unbranched.

^bFitch and Lavenberg 1975

^cE.H. Ahlstrom notes





Figure A, Matarese and Sandknop 1984; B–D, Ahlstrom et al. 1984a.



Tetraodontiformes

Although most fishes of the Tetraodontiformes are tropical and associated with the bottom, some are found in temperate zones and remain pelagic throughout their lifespans. The general body shape is rounded or boxlike and the body may be encased in a bony carapace or covered with sharp spines. Many species are able to inflate themselves with water or air. Eight families, 92 genera, and 329 species make up the order (J. Nelson 1984). Only one family, Molidae, is found in the Northeast Pacific. The most unusual of the tetraodontiforms, molids have no caudal fin and propel themselves with large dorsal and anal fins. Eggs are pelagic, 1.42-1.80 mm in diameter, and have multiple oil globules (Leis 1984). Larvae hatch with a functioning jaw, pigmented eyes, and a dermal sac enclosing the head and trunk. With development, body spines form and the tail atrophies. Molids may have a long prejuvenile stage marked by retention of spines and a shape unlike that of adults (Leis 1984).

Family in study area: **Molidae**

MERISTICS

Vertebrae	Total: 17-X-18 Precaudal: 8-X-9 Caudal: 8-9-9
Branchiostegal rays	6-6-6
Caudal fin	Absent
Pelvic fin	Absent
Dorsal fin	R: 15-X-18
Pectoral fin	R: 11-X-13
Anal fin	R: 14-X-18
Gill rakers	U: X-X-X L: X-X-X

LIFE HISTORY

Range	South of southern California to SE Alaska, 55-59°N
Ecology	Epipelagic
ELH pattern	Oviparous, pelagic eggs, pelagic larvae
Spawning	Season: Spring-summer ^a Area: Mode: Migration:
Fecundity	Range/function: 300 million ^b
Age at first maturity	
Longevity	

EARLY LIFE HISTORY DESCRIPTION**EGGS**

Diameter	
No. of oil globules	
Oil globule diameter	
Yolk	
Envelope	
Hatch size	<1.84 mm TL
Incubation time/temp.	Other members of family hatch in 7-8 days
Pigment	

Diagnostic characters - Family

- Pelagic, large (1.4-1.8 mm), and have multiple oil globules





Larvae

Preanal length	<50% SL increasing with development to >50% SL
Length at flexion	Does not occur
Length at transformation	Long ontogenetic stage between larvae and juveniles
Sequence of fin development	Pectorals and caudal, dorsal and anal
Pigment - Family	
	• Usually heavily pigmented over gut and dorsal surfaces



Diagnostic characters

- Morphology: Wide, deep body
- Body spines with a ventral keel (form soon after hatching)
- Tail in young is normal but soon atrophies and a true caudal never forms; notochord flexion does not take place, so the clavus (gradual thickening near the distal end) is not homologous with the caudal fin^c

^aMartin and Drewry 1978^bHart 1973^cLeis 1984



Body spines reach maximum size at the same time the clavus forms



Figures A–F (A, dorsal view; E, anterior view), Martin and Drewry 1978 (North Atlantic specimens).

Citations

- Abe, T., and S. Kaji**
1972 A record of *Oreosoma atlanticum* (Oreosomatidae, Zeiformes, Teleostei) from Tasman Sea. UO (Sakana) No. 8:5-7, Ichthyol. Soc. Jpn., Tokyo.
- Able, K.W., D.F. Markle, and M.P. Fahay**
1984 Cyclopteridae: Development. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 428-437. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Aboussouan, A., and R. Rasonarivo**
1986 Capture d'une larve de *Spectrunculus grandis* (Günther, 1877) dans l'ouest de l'océan Indien, île de la Réunion (Pisces, Ophidiiformes, Ophidiidae). Cybium 10:206-207 [in French].
- Ahlstrom, E.H.**
1965 Kinds and abundance of fishes in the California Current region based on egg and larval surveys. Calif. Coop. Oceanic Fish. Invest. Rep. 10:31-52.
1969 Remarkable movement of oil globules in eggs of bathylagid smelts during embryonic development. J. Mar. Biol. Assoc. India 11: 206-217.
1972 Distributional atlas of fish larvae in the California Current region: Six common mesopelagic fishes, *Vinciguerria lucetia*, *Triphoturus mexicanus*, *Stenobranchius leucopsarus*, *Leuroglossus stilbius*, *Bathylagus wesethi*, and *Bathylagus ochotensis*. Calif. Coop. Oceanic Fish. Invest. Atlas 17, 306 p.
1974 The diverse patterns of metamorphosis in gonostomatid fishes—An aid to classification. In Blaxter, J.H.S. (ed.), The early life history of fish, p. 659-674. Springer-Verlag, NY.
- Ahlstrom, E.H., and O.P. Ball**
1954 Description of eggs and larvae of jack mackerel (*Trachurus symmetricus*) and distribution and abundance of larvae in 1950 and 1951. U.S. Fish Wildl. Serv., Fish. Bull. 56:209-245.
- Ahlstrom, E.H., and R.C. Counts**
1955 Eggs and larvae of the Pacific hake, *Merluccius productus*. U.S. Fish Wildl. Serv., Fish. Bull. 56:295-329.
- Ahlstrom, E.H., and H.G. Moser**
1975 Distributional atlas of fish larvae in the California Current region: Flatfishes, 1955 through 1960. Calif. Coop. Oceanic Fish. Invest. Atlas 23, 207 p.
1976 Eggs and larvae of fishes and their role in systematic investigations and in fisheries. Rev. Trav. Inst. Pêches Marit. 40:379-398.
1980 Characters useful in identification of pelagic marine fish eggs. Calif. Coop. Oceanic Fish. Invest. Rep. 21:121-131.
- Ahlstrom, E.H., and E. Stevens**
1976 Report of neuston (surface) collections made on an extended CalCOFI cruise during May 1972. Calif. Coop. Oceanic Fish. Invest. Rep. 18:167-180.
- Ahlstrom, E.H., J.L. Butler, and B.Y. Sumida**
1976 Pelagic stromateoid fishes (Pisces, Perciformes) of the eastern Pacific: Kinds, distributions, and early life histories and observations on five of these from the Northwest Atlantic. Bull. Mar. Sci. 26:285-402.
- Ahlstrom, E.H., K. Amaoka, D.A. Hensley, H.G. Moser, and B.Y. Sumida**
1984a Pleuronectiformes: Development. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 640-669. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
1984b Argentinoidae: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 155-168. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Ahlstrom, E.H., W.J. Richards, and S.H. Weitzman**
1984c Families Gonostomatidae, Sternoptychidae, and associated stomiiform groups: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 184-198. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Alaska Department of Fish and Game**
1985 Alaska habitat management guide. South central region, Vol. 1: Life histories and habitat requirements of fish and wildlife. Alaska Dep. Fish Game, Juneau, AK 99811, 429 p.
- Alderdice, D.F., and C.R. Forrester**
1971 Effects of salinity and temperature on embryonic development of the petrale sole (*Eopsetta jordani*). J. Fish. Res. Board Can. 28:727-744.
- Allen, L.G.**
1979 Larval development of *Gobiesox rhesodon* (Gobiesocidae) with notes on the larva of *Rimicola muscarum*. Fish. Bull., U.S. 77: 300-304.
1984 Gobiesociformes: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 629-636. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Allen, L.G., and J.M. Ilg**
1983 Larval development of the northern clingfish, *Gobiesox maeandricus*. Copeia 1983:551-554.
- Allen, M.J., and G.B. Smith**
1988 Atlas and zoogeography of common fishes in the Bering Sea and northeastern Pacific. NOAA Tech. Rep. NMFS 66. Natl. Oceanic Atmos. Adm., Natl. Mar. Fish. Serv., Seattle, WA 98115-0070, 151 p.
- Alton, M.S., and R.A. Webber**
1976 Sablefish (family Anoplopomatidae). In Pereya, W.T., et al. (eds.), Demersal fish and shellfish resources in the eastern Bering Sea in the baseline year 1975, p. 425-438. Proc. Rep., Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070.
- Ambrose, D.A., J.L. Butler, H.G. Moser, B.Y. Sumida, E.M. Sandknop, and E.G. Stevens**
1983 Description of the larvae of the cusk eels *Ophidion scrippsae* and *Chilara taylori* (family Ophidiidae). Calif. Coop. Oceanic Fish. Invest. Rep. 24:226-234.
- Anderson, M.E.**
1984a On the anatomy and phylogeny of the Zoarcidae (Teleostei: Perciformes). Ph.D. diss., Coll. William and Mary, Williamsburg, VA 23185, 254 p.
1984b Zoarcidae: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 578-581. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Anderson, M.E., and G.M. Cailliet**
1974 Crab and snailfish commensalism in Monterey Bay. Underwater Nat. 8(3):29-31.
- Andriashev, A.P.**
1954 Fishes of the northern seas of the U.S.S.R. Tr. Zool. Inst. Akad. Nauk SSSR 53. [In Russ., transl. by Isr. Prog. Sci. Transl., Jerusalem, 1964; avail. U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA 22161, as OTS63-11160.]
- Arita, G.S.**
1969 Sexual dimorphism in the cyclopterid fish *Eumicrotremus orbis*. J. Fish. Res. Board Can. 26:3262-3265.
- Arora, H.L.**
1951 An investigation of the California sanddab, *Citharichthys sordidus* (Girard). Calif. Fish Game 37:3-42.
- Atkinson, C.E.**
1939 Notes on the life-history of the tidepool johnny (*Oligocottus maculosus*). Copeia 1939:23-30.
- Auer, N.A. (editor)**
1982 Identification of larval fishes of the Great Lakes Basin with emphasis on the Lake Michigan drainage. Spec. Publ. 82-3, Great Lakes Fish Comm., Ann Arbor, MI 48105, 744 p.

- Badcock, J., and R.C. Baird**
1980 Remarks on systematics, development, and distribution of the hatchetfish genus *Sternopyx* (Pisces, Stomiatoidei). *Fish. Bull.*, U.S. 77:803-820.
- Badcock, J., and N.R. Merrett**
1976 Midwater fishes in the eastern North Atlantic. I. Vertical distribution and associated biology in 30 degrees N, 23 degrees W, with developmental notes on certain myctophids. *Prog. Oceanogr.* 7:3-58.
- Bailey, K.M., R.C. Francis, and P.R. Stevens**
1982 The life history and fishery of the Pacific whiting, *Merluccius productus*. *Calif. Coop. Fish. Invest. Rep.* 23:81-98.
- Bain, H., and A.D. Sekerak**
1978 Aspects of the biology of Arctic cod, *Boreogadus saida*, in the central Canadian Arctic. Report prepared for Polar Gas Project by LGL Limited, Environ. Res. Assoc., Toronto, Ontario, Canada, 104 p.
- Barnhart, P.S.**
1932 Notes on the habits, eggs, and young of some fishes of southern California. *Bull. Scripps Inst. Oceanogr. Univ. Calif.* 3(4):87-99.
- Barsukov, V.V.**
1959 The wolffish (Anarhichadidae). *Fauna SSSR Moscow, N.S.* 73. [Transl. by Smithson. Inst., Wash., D.C., 1972; avail. U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA 22161, as TT67-59074.]
- Bauchot, M.**
1959 Etude des larves leptocephalus du groupe *Leptocephalus lanceolatus* Stromman et identification a la famille des Serrivomeridae. *Dana-Rep. Carlsberg Found.* 48, 148 p. [in French].
- Baxter, J.L.**
1967 Summary of biological information on the northern anchovy *Engraulis mordax* Girard. *Calif. Coop. Oceanic Fish. Invest. Rep.* 11:110-116.
- Baxter, R.**
1975 Inshore marine resources of Bristol Bay, Alaska. *Alaska Dep. Fish Game, Bethel, AK* 99559, 97 p.
- Beamish, R.J.**
1979 Differences in the age of Pacific hake (*Merluccius productus*) using whole otoliths and sections of otoliths. *J. Fish. Res. Board Can.* 36:141-151.
- Beebe, W.**
1933 Deep-sea fishes of the Bermuda Oceanographic Expeditions. No. 2, Family Alepocephalidae. *Zoologica (NY)* 16(2):15-93.
1934 Deep-sea fishes of the Bermuda Oceanographic Expeditions. Family Iriacanthidae. *Zoologica (NY)* 16(4):97-147.
- Beebe, W., and J. Crane**
1939 Deep-sea fishes of the Bermuda Oceanographic Expeditions. Family Melanostomiidae. *Zoologica (NY)* 24(2):65-238.
- Belyanina, T.N.**
1983 Developmental sequences of *Sternopyx* species (Sternopychidae). *J. Ichthyol.* 23(4):73-86.
1984 Larvae of hatchetfishes of the genus *Argyropelecus* (Sternopychidae). *J. Ichthyol.* 24(2):7-20.
- Berrien, P.L.**
1978 Eggs and larvae of *Scomber scombrus* and *Scomber japonicus* in continental shelf waters between Massachusetts and Florida. *Fish. Bull.*, U.S. 76:95-115.
- Bertelsen, E.**
1951 The ceratioid fishes. Ontogeny, taxonomy, distribution, and biology. *Dana-Rep. Carlsberg Found.* 39, 276 p.
1984 Ceratioidei: Development and relationships. *In* Moser, H.G., et al. (eds.), *Ontogeny and systematics of fishes*, p. 325-333. *Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol.* Allen Press, Lawrence, KS.
- Bertelsen, E., G. Krefft, and N.B. Marshall**
1976 The fishes of the family Notosudidae. *Dana-Rep. Carlsberg Found.* 86, 114 p.
- Best, E.A.**
1963 Contribution to the biology of the Pacific hake, *Merluccius productus* (Ayres). *Calif. Coop. Oceanic Fish. Invest. Rep.* 9:51-56.
- Bigelow, H.B., and W.C. Schroeder**
1953 Fishes of the Gulf of Maine. *U.S. Fish Wildl. Serv., Fish. Bull.* 74(vol. 53), 577 p.
- Blackburn, J.E.**
1973 A survey of the abundance, distribution, and factors affecting distribution of ichthyoplankton in Skagit Bay. *M.S. thesis, Univ. Wash., Seattle, WA* 98195, 136 p.
- Boehlert, G.W., and M.M. Yoklavich**
1984 Reproduction, embryonic energetics, and the maternal-fetal relationship in the viviparous genus *Sebastes* (Pisces: Scorpaenidae). *Biol. Bull. (Woods Hole)* 167:354-370.
- Bolin, R.L.**
1936 Embryonic and early larval stages of the California anchovy. *Calif. Fish Game* 22:314-321.
- Bond, C.E., and D.L. Stein**
1984 *Opaephacus acrogeneius*, a new genus and species of Zoarcidae (Pisces: Osteichthyes) from the Bering Sea. *Proc. Biol. Soc. Wash.* 97(3):522-525.
- Breder, C.M., and D.E. Rosen**
1966 Modes of reproduction in fishes. *T.F.H. Publ.*, Jersey City, NJ, 941 p.
- Brothers, E.B.**
1975 The comparative ecology and behavior of three sympatric California gobies. *Ph.D. diss., Univ. Calif., San Diego, CA* 92037, 370 p.
- Budd, P.L.**
1940 Development of the eggs and early larvae of six California fishes. *Calif. Dep. Fish Game, Fish Bull.* 56, 50 p.
- Burge, R.T., and S.A. Schultz**
1973 The marine environment in the vicinity of Diablo Cove with special reference to abalones and bony fishes. *Calif. Dep. Fish Game, Mar. Res. Tech. Rep.* 19, 239 p.
- Castle, P.H.**
1965 Leptocephali of the Nemichthyidae, Serrivomeridae, Synphobranchidae, and Nettastomatidae in Australian waters. *Trans. R. Soc. N.Z. Zool.* 5:131-146.
1984 Notacanthiformes and Anguilliformes: Development. *In* Moser, H.G., et al. (eds.), *Ontogeny and systematics of fishes*, p. 62-93. *Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol.* Allen Press, Lawrence, KS.
- Castle, P.H., and N.S. Raju**
1975 Some rare leptocephali from the Atlantic and Indo-Pacific Oceans. *Dana-Rep. Carlsberg Found.* 85, 25 p.
- Chapman, W.M.**
1939 Eleven new species and three new genera of oceanic fishes collected by the International Fisheries Commission from the northeastern Pacific. *Proc. U.S. Natl. Mus.* 86(3062):501-542.
- Chapman, W.M., and L.D. Townsend**
1938 The osteology of *Zaprora silenus* Jordan, with notes on its distribution and early life history. *Ann. Mag. Nat. Hist.* 11:89-117.
- Chen, L.**
1986 Meristic variation in *Sebastes* (Scorpaenidae) with an analysis of character association and bilateral pattern and their significance in species separation. *NOAA Tech. Rep. NMFS* 45, *Natl. Oceanic Atmos. Adm., Natl. Mar. Fish. Serv., Seattle, WA* 98115-0070, 17 p.
- Chilton, D.E., and R.J. Beamish**
1982 Age determination methods for fishes studied by the groundfish program at the Pacific Biological Station. *Can. Spec. Publ. Fish. Aquat. Sci.* 60, 102 p.
- Chung, M.**
1977 The fishes of Korea. *Il Ji Sa Publ. Co., Seoul, Korea.*
- Clark, F.N., and J.B. Phillips**
1952 The northern anchovy (*Engraulis mordax*) in the California fishery. *Calif. Fish Game* 38:189-207.
- Clemens, W.A., and G.V. Wilby**
1961 Fishes of the Pacific coast of Canada. *Fish. Res. Board Can. Bull.* 68, 2d ed., 443 p.
- Cohen, D.M.**
1958 A revision of the fishes of the subfamily Argentininae. *Bull. Fla. State Mus. Biol. Ser.* 3:93-172.

- 1960 New records of the opisthoproctid genus *Bathylychnops*, with a notice of neoteny in the related genus *Dolichopteryx*. *Copeia* 1960:147-149.
- Cohen, D.M., and J.G. Nielsen**
1978 Guide to the identification of genera of the fish order Ophidiiformes, with a tentative classification of the order. NOAA Tech. Rep. NMFS Circ. 417, Natl. Oceanic Atmos. Adm., Natl. Mar. Fish. Serv., Seattle, WA 98115-0070, 72 p.
- Collette, B.B.**
1984 Atherinomorpha: Introduction. In Moser, H.G., et al. (eds.), *Ontogeny and systematics of fishes*, p. 334. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Collette, B.B., and C.E. Nauen**
1983 Scombrids of the world. FAO Fish. Synop. 125, Vol. 2, 137 p.
- Collette, B.B., G.E. McGowen, N.V. Parin, and S. Mito**
1984a Beloniformes: Development and relationships. In Moser, H.G., et al. (eds.), *Ontogeny and systematics of fishes*, p. 335-354. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Collette, B.B., T. Pothoff, W.J. Richards, S. Ueyanagi, J.L. Russo, and Y. Nishikawa**
1984b Scombroidei: Development and relationships. In Moser, H.G., et al. (eds.), *Ontogeny and systematics of fishes*, p. 591-619. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Craig, P.C., W.B. Griffiths, L. Haldorson, and H. McElderry**
1982 Ecological studies of Arctic cod (*Boreogadus saida*) in Beaufort Sea coastal waters, Alaska. *Can. J. Fish. Aquat. Sci.* 39:395-406.
- DeLacy, A.C., C.R. Hitz, and R.L. Dryfoos**
1964 Maturation, gestation, and birth of rockfish (*Sebastes*) from Washington and adjacent waters. *Wash. Dep. Fish., Fish. Res. Pap.* 2(3):51-67.
- DeMartini, E.E.**
1976 The adaptive significance of territoriality and egg cannibalism in the painted greenling, *Oxylebius pictus* Gill, a northeastern Pacific marine fish. Ph.D. diss., Univ. Wash., Seattle, WA 98195, 286 p.
1978 Spatial aspects of reproduction in buffalo sculpin, *Enophrys bison*. *Environ. Biol. Fishes* 3:331-336.
- Dunn, J.R.**
1983 Development and distribution of young of northern smooth-tongue, *Leuroglossus schmidti* (Bathylagidae) in the Northeast Pacific, with comments on the systematics of the genus *Leuroglossus* Gilbert. *Fish. Bull., U.S.* 81:23-40.
1986 A catalog of Northwest and Alaska Fisheries Center ichthyoplankton cruises 1965-1985. Proc. Rep. 86-08, Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070, 164 p.
- Dunn, J.R., and A.C. Matarese**
1984 Gadidae: Development and relationships. In Moser, H.G., et al. (eds.), *Ontogeny and systematics of fishes*, p. 283-289. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
1987 A review of the early life history of Northeast Pacific gadoid fishes. *Fish. Res. (Amst.)* 5:163-184.
- Dunn, J.R., and T.M. Sample**
1976 Greenland halibut (family Pleuronectidae). In Pereya, W.T., et al. (eds.), *Demersal fish and shellfish resources in the eastern Bering Sea in the baseline year 1975*, p. 475-487. Proc. Rep., Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070.
- Dunn, J.R., and B.M. Vinter**
1984 Development of larvae of saffron cod, *Eleginus gracilis*, with criteria for identification of gadid larvae in Pacific and Arctic waters contiguous to Canada and Alaska. *Can. J. Fish. Aquat. Sci.* 41: 304-318.
- D'Vincent, S., H.G. Moser, and E.H. Ahlstrom**
1980 Description of the larvae and early juveniles of Pacific butterflyfish, *Pepritis similimus* (family Stromateidae). *Calif. Coop. Oceanic Fish. Invest. Rep.* 21:172-179.
- Ebeling, A.W.**
1962 Melamphaidae. I. Systematics and zoogeography of the species in the bathypelagic fish genus *Melamphaes* Günther. *Dana-Rep. Carlsberg Found.* 58, 164 p.
- Ebert, E.E., and C.H. Turner**
1962 The nesting behavior, eggs, and larvae of the bluespot goby. *Calif. Fish Game* 48:249-252.
- Efremenko, V.N., and L.A. Lisovenko**
1970 Morphological features of intraovarian and pelagic larvae of some *Sebastes* species inhabiting the Gulf of Alaska. In Moiseev, P.A. (ed.), *Soviet fisheries investigations in the northeastern Pacific*, Pt. V, p. 267-286. *Izv. Tikhookean. Nauchno-Issled. Inst. Rybn. Khoz. Okeanogr. (TINRO)* 72; Tr. Vses. Nauchno-Issled. Inst. Morsk. Rybn. Khoz. Okeanogr. (VNIRO) 70. [Transl. from Russ.; avail. U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA 22161, as TT71-50127.]
- English, T.S.**
1961 An inquiry into distributions of planktonic fish eggs in a restricted area of Puget Sound. Ph.D. diss., Univ. Wash., Seattle, WA 98195, 227 p.
- Eschmeyer, W.N., E.S. Herald, and H. Hammann**
1983 A field guide to Pacific Coast fishes of North America from the Gulf of Alaska to Baja California. Houghton Mifflin Co., Boston, 336 p.
- Evseenko, S.A.**
1982 Ichthyoplankton of slope and Gulf Stream waters off Nova Scotia in late autumn 1974. *J. Northwest Atl. Fish. Sci.* 3:127-139.
- Faber, D.J.**
1976 Identification of four northern blennioid fish larvae in the Canadian Atlantic Ocean (Stichaeidae, Lumpeninae). *J. Fish. Res. Board Can.* 33:1798-1802.
- Fadeev, N.S.**
1965 Comparative outline of the biology of flatfishes in the southeastern part of the Bering Sea and condition of their resources. In Moiseev, P.A. (ed.), *Soviet fisheries investigations in the Northeast Pacific*, Pt. IV, p.112-119. *Izv. Tikhookean. Nauchno-Issled. Inst. Rybn. Khoz. Okeanogr. (TINRO)* 53; Tr. Vses. Nauchno-Issled. Inst. Morsk. Rybn. Khoz. Okeanogr. (VNIRO) 58. [Transl. from Russ. by Isr. Prog. Sci. Transl., Jerusalem, 1968; avail. U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA 22161, as TT67-51206.]
- Fahay, M.P.**
1983 Guide to the early stages of marine fishes occurring in the western North Atlantic Ocean, Cape Hatteras to the southern Scotian Shelf. *J. Northwest Atl. Fish. Sci.* 4, 423 p.
- Fahay, M.P., and D.F. Markle**
1984 Gadiformes: Development and relationships. In Moser, H.G., et al. (eds.), *Ontogeny and systematics of fishes*, p. 265-282. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Fast, T.N.**
1960 Some aspects of the natural history of *Stenobranchius leucopsarus* Eigenmann and Eigenmann. Ph.D. diss., Stanford Univ., Stanford, CA 94305, 107 p.
- Feder, H.M., C.H. Turner, and C. Limbaugh**
1974 Observations on fishes associated with kelp beds in southern California. *Calif. Dep. Fish Game, Fish Bull.* 160, 144 p.
- Fedorov, V.V.**
1973 Ikhtiofauna materikovogo sklona beringova morya i nekotorye aspekty ee proiskhozhdeniya i formirovaniya. *Izv. Tikhookean. Nauchno-Issled. Inst. Rybn. Khoz. Okeanogr. (TINRO)* 87: 3-41. [Transl. by Transl. Bur. (DCM), Multilingual Serv. Div., Dep. of Secretary of State of Canada, Transl. Ser. 3345, 70 p.; avail. Can. Dep. Fish. Oceans, Fish. Res. Br., Pac. Biol. Stn., Nanaimo, B.C., Canada V9R 5K6.]

- Fink, W.L.**
1984 Stomiiforms: Relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 181-184. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Fink, W.L., and S.H. Weitzman**
1982 Relationships of the stomiiform fishes (Teleostei), with a description of *Diplophos*. Bull. Mus. Comp. Zool. Harv. Univ. 150(2): 31-93.
- Fisher, J.P., and W.G. Pearcy**
1983 Reproduction, growth and feeding of the mesopelagic fish *Tacostoma macropus* (Melanostomiidae). Mar. Biol. 74:257-267.
- Fitch, J.E., and R.J. Lavenberg**
1968 Deep-water fishes of California. Univ. Calif. Press, Berkeley, 155 p.
1971 Marine food and game fishes of California. Univ. Calif. Press, Berkeley, 179 p.
1975 Tidepool and nearshore fishes of California. Univ. Calif. Press, Berkeley, 156 p.
- Forrester, C.R.**
1964 Rate of development of eggs of rock sole (*Lepidopsetta bilineata* Ayres). J. Fish. Res. Board Can. 21:1533-1534.
1969 Life history information on some groundfish species. Fish. Res. Board Can. Tech. Rep. 105, 17 p.
- Francis, R.C., and K.M. Bailey**
1983 Factors affecting recruitment of selected gadoids in the Northeast Pacific and eastern Bering Sea. In Wooster, W.S. (ed.), From year to year: Interannual variability of the environment and fisheries of the Gulf of Alaska and the eastern Bering Sea. Rep WSG-WO 83-3, Wash. Sea Grant Prog., Univ. Wash., Seattle, WA 98195.
- Frey, H.W. (editor)**
1971 California's living marine resources and their utilization. Calif. Dep. Fish Game, Sacramento, CA 95814, 148 p.
- Fritzsche, R.A.**
1978 Development of fishes of the Mid-Atlantic Bight: An atlas of egg, larval and juvenile stages. Vol. V, Chaetodontidae through Ophidiidae. U.S. Fish. Wildl. Serv. Biol. Rep. FWS/OBS-78/12.
1984 Gasterosteiformes: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 398-404. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Garrison, K.J., and B.S. Miller**
1982 Review of the early life history of Puget Sound fishes. Unpubl. rep. FRI-UW-8216, Fish. Res. Inst., Univ. Wash., Seattle, WA 98195, 729 p.
- Giorgi, A.E.**
1981 The environmental biology of the embryos, egg masses, and nesting sites of the lingcod, *Ophiodon elongatus*. Proc. Rep. 81-06, Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070, 107 p.
- Goldberg, S.R.**
1980a Seasonal spawning cycle of the longspine combfish, *Zaniolepis latipinnis*, with notes on spawning of the shortspine combfish, *Zaniolepis frenata* (Zaniolepidae). Copeia 1980:882-884.
1980b Seasonal spawning cycles of two marine cottid fishes, *Chitonotus pugetensis* and *Icelinus quadricornis*, from southern California. Bull. Mar. Sci. 30:131-135.
- Gorbunova, N.N.**
1954 Reproduction and development of the walleye pollock, *Theragra chalcogramma* (Pallas). Tr. Inst. Okeanol. Akad. Nauk SSSR 11: 132-195. [In Russ., transl. by S. Pearson, 1972, Natl. Mar. Mammal Lab., NMFS, 7600 Sand Point Way N.E., Seattle, WA 98115-0070.]
1962 Spawning and development of greenlings (family Hexagrammidae). In Rass, T.S. (ed.), Greenlings: Taxonomy, biology, inter-oceanic transplantation. Tr. Inst. Okeanol. Akad. Nauk SSSR 59: 121-185. [Engl. transl. avail. U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA 22161, as TT69-55097.]
- 1964** Breeding and development of hemilepidotine sculpins (Cottidae, Pisces). In Rass, T.S. (ed.), Fishes of the Pacific and Indian Oceans. Biology and distribution, p. 249-266. Tr. Inst. Okeanol. Akad. Nauk SSSR 73. [Isr. Prog. Sci. Transl., Jerusalem, Transl. 1411.]
- 1982a** Larvae of Pacific species of *Cyathothone* genus. Tr. Inst. Okeanol. Akad. Nauk SSSR 118:120-132 [in Russ.]
1982b Larvae of trichiurid fishes from the collection of the International Mexican Biological Center (Gempylidae, Trichiuridae, Pisces). Proc. P.P. Shirshov. Inst. Oceanol. 118:85-104 [in Russ.].
- Gordon, D.J., D.F. Markle, and J.E. Olney**
1984 Ophidiiformes: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 308-319. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Grant, W.S.**
1986 Biochemical genetic divergence between Atlantic, *Clupea harengus*, and Pacific, *C. pallasii*, herring. Copeia 1986:714-719.
- Grey, M.**
1964 Family Gonostomatidae. In Bigelow, H.B. (ed.), Fishes of the western North Atlantic, p. 78-240. Mem. 1 (pt. 4), Sears Found. Mar. Res., Yale Univ., New Haven, CT.
- Grossman, G.D., and V. DeVlaming**
1984 Reproductive ecology of female *Oligocottus snyderi*, a North American intertidal sculpin. J. Fish. Biol. 25:231-240.
- Gunderson, D.R., P. Callahan, and B. Goiney**
1980 Maturation and fecundity of four species of *Sebastes*. Mar. Fish. Rev. 42(3-4):74-79.
- Hagerman, F.B.**
1952 The biology of the dover sole, *Microstomus pacificus* (Lockington). Calif. Dep. Fish Game, Fish Bull. 85, 48 p.
- Hamada, T.**
1966 Studies on fluctuations in the abundance of larval sand-lance in the Harima-nada and Osaka Bay. I. Relation between the progeny abundance and the age composition of parent fish. Bull. Jpn. Soc. Sci. Fish. 32(5):393-398 [in Jpn., Engl. summ.].
- Hardy, G.S.**
1983 A revision of the fishes of the family Pentacerotidae (Perciformes). N.Z. J. Zool. 10:177-220.
- Harling, W.R., M.S. Smith, and N.A. Webb**
1971 Preliminary report on maturity, spawning season, and larval identification of rockfishes (Scorpaenidae) collected during 1970. Fish. Res. Board Can., Manuscr. Rep. 1137, 26 p.
- Harry, G.Y.**
1959 Time of spawning, length of maturity, and fecundity of the English, petrale, and Dover soles (*Parophrys vetulus*, *Eopsetta jordani*, and *Microstomus pacificus*, respectively). Fish. Comm. Oreg. Res. Briefs 7(1):5-13 (Portland, OR 97201).
- Hart, J.L.**
1967 Fecundity and length-weight relationship in lingcod. J. Fish. Res. Board Can. 24:2485-2489.
1973 Pacific fishes of Canada. Bull. Fish. Res. Board Can. 180, 740 p.
- Hart, J.L., and J.L. McHugh**
1944 The smelts (Osmeridae) of British Columbia. Bull. Fish. Res. Board Can. 64, 27 p.
- Haryu, T., and T. Nishiyama**
1981 Larval form of zaprorid fish *Zaprora silenus* from the Bering Sea and the northern North Pacific. Jpn. J. Ichthyol. 28:313-319.
- Hattori, S.**
1964 Studies on fish larvae in the Kuroshio and adjacent waters. Bull. Tokai Reg. Fish. Res. Lab. 40, 158 p. [in Jpn., Engl. synop.].
- Hearne, M.E.**
1983 Identification of larval and juvenile smelts (Osmeridae) from California and Oregon using selected morphometric characters. M.S. thesis, San Francisco State Univ., San Francisco, CA 94132, 142 p.
1984 Osmeridae: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 153-154. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.

- Hedgpeth, J.W.**
1957 Classification of marine environments. Geol. Soc. Am. Mem. 67(1):17-28.
- Hickman, C.P.**
1959 The larval development of the sand sole (*Psetichthys melanostictus*). Wash. Dep. Fish., Fish. Res. Pap. 2(2):38-47.
- Hinckley, S.**
1986 Spawning dynamics and fecundity of walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea. M.S. thesis, Univ. Wash., Seattle, WA 98195, 103 p.
- Hirschberger, W.A., and G.B. Smith**
1983 Spawning of twelve groundfish species in the Gulf of Alaska and Pacific Coast regions, 1975-81. Tech. Memo. NMFS F/NWC-44, Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070, 50 p.
- Hitz, C.R.**
1962 Seasons of birth of rockfish (*Sebastes* spp.) in Oregon coastal waters. Trans. Am. Fish. Soc. 91:231-233.
- Hollister, G.**
1934 Clearing and dyeing fish for bone study. Zoologica 12(10): 89-101.
- Horn, M.H.**
1984 Stromateoidei: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 620-628. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Hosie, M.J., and H.F. Horton**
1977 Biology of the rex sole, *Glyptocephalus zachirus*, in waters off Oregon. Fish. Bull., U.S. 75:51-60.
- Hubbs, C.L., and T. Iwamoto**
1977 A new genus (*Mesobius*), and three new bathypelagic species of Macrouridae (Pisces, Gadiformes) from the Pacific Ocean. Proc. Calif. Acad. Sci. 41:233-251.
- Hubbs, C.L., W.I. Follett, and L.J. Dempster**
1979 List of the fishes of California. Occas. Pap. Calif. Acad. Sci. 133:1-51.
- Humphreys, R.L., Jr., G.A. Winans, and D.T. Tagami**
1989 Synonymy and proposed life history of North Pacific pelagic armorhead, *Pseudopentaceros wheeleri* Hardy (Pisces: Pentacero-tidae). Copeia 1989:142-153.
- Ida, H.**
1976 Removal of the family Hypoptychidae from the suborder Am-modytoidei, order Perciformes, to the suborder Gasterosteioidei, order Syngnathiformes. Jpn. J. Ichthyol. 23:33-42.
- Idyll, C.P.**
1964 Abyss: The deep sea and the creatures that live in it. Thomas Y. Cromwell Co., NY, 395 p.
- Inoue, A., S. Takamori, K. Kuniyaki, S. Kobayashi, and S. Nishina**
1967 Studies on fishery biology of the sand-lance, *Ammodytes per-sonatus* (Girard). Bull. Naikai Reg. Fish. Res. Lab. 25(121):1-335 [in Jpn., Engl. summ.].
- Iwamoto, T.**
1975 The abyssal fish *Antimora rostrata* (Günther). Comp. Biochem. Physiol. 52b:7-11.
- Jensen, A.**
1935 The greenland halibut (*Reinhardtius hippoglossoides*), its development and migration. K. Dan. Vidensk. Selsk. Skr. 9:1-32.
- Jespersen, P., and A.V. Tåning**
1926 Mediterranean Sternoptychidae. Rep. Dan. Oceanogr. Exped. Mediterr. 2(A.12), 59 p.
- Jewell, E.D.**
1968 Scuba diving operations on lingcod spawning at a Seattle break-water. Wash. Dep. Fish., Fish. Res. Pap. 3(1):27-36, Olympia, WA 98504.
- Johnson, C.L.**
1970 Notes on the intertidal life history of the northern clingfish, *Gobiesox maeandricus* (Girard). Am. Midl. Nat. 83(2):625-627.
- Johnson, G.D.**
1984 Percoidae: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 464-498. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Johnson, R.K.**
1974a A revision of the alepisaurid family Scopelarchidae (Pisces: Myctophiformes). Fieldiana Zool. 66:1-249.
1974b A *Macristium* larva from the Gulf of Mexico with additional evidence for synonymy of *Macristium* with *Bathysaurus* (Mycto-phiformes: Bathysauridae). Copeia 1974:973-977.
1984 Scopelarchidae: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 245-249. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Jones, A.C.**
1962 The biology of euryhaline fish *Leptocottus armatus armatus* (Girard). Univ. Calif. Publ. Zool. 67(4):321-368.
- Jow, T.**
1969 Results of English sole tagging off California. Pac. Mar. Fish. Comm. Bull. 7:15-33.
- Kabata, Z., and C.R. Forrester**
1974 *Atheresthes stomias* (Pisces: Pleuronectiformes) and its eye parasite *Phrizocephalus cincinnatus* (Copepods: Lernaeoceridae) in Canadian Pacific waters. J. Fish. Res. Board Can. 31:1589-1595.
- Karp, W.A.**
1982 Biology and management of Pacific cod (*Gadus macrocephalus*) in Port Townsend, Washington. Ph.D. diss., Univ. Wash., Seattle, WA 98195, 119 p.
- Katz, M.**
1942 The herring races of Washington, with a note on the fecundity of the Seal Rock population. M.S. thesis, Univ. Wash., Seattle, WA 98195, 65 p.
- Kawaguchi, K., and R. Marumo**
1967 Biology of *Gonostoma gracile* (Gonostomatidae). 1. Morphology, life history, and sex reversal. In Information bulletin on plank-tology in Japan, Commemoration number of Dr. Y. Matsui's sixtieth birthday, p. 253-269. Plankton Soc. Jpn., Tokyo.
- Kawaguchi, K., and H.G. Moser**
1984 Stomiatoidea: Development. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 169-180. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Keene, M.J., and K.A. Tighe**
1984 Beryciformes: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 383-392. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Kendall, A.W., Jr., and R.J. Behnke**
1984 Salmonidae: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 142-149. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Kendall, A.W., Jr., and J. Clark**
1982 Ichthyoplankton off Washington, Oregon, and northern California April-May 1980. Proc. Rep. 82-11, Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070, 44 p.
- Kendall, A.W., Jr., and A.C. Matarese**
1987 Biology of eggs, larvae, and epipelagic juveniles of sablefish, *Anoplopoma fimbria*, in relation to their potential use in management. Mar. Fish. Rev. 49(1):1-13.
- Kendall, A.W., Jr., and B. Vinter**
1984 Development of hexagrammids (Pisces, Scorpaeniformes) in the northeastern Pacific Ocean. NOAA Tech. Rep. NMFS 2, Natl. Oceanic Atmos. Adm., Natl. Mar. Fish. Serv., Seattle, WA 98115-0070, 44 p.
- Kendall, A.W., Jr., C.D. Jennings, T.M. Beasley, R. Carpenter, and B.L. Somayajulu**
1983 Discovery of a cluster of unhatched fish eggs of a zoarcid buried 10 to 12 cm deep in continental slope sediments off Washington state, USA. Mar. Biol. 75:193-199.

- Kendall, A.W., Jr., E.H. Ahlstrom, and H.G. Moser**
1984 Early life history stages of fishes and their characters. In Moser, H.G., et al. (eds.), *Ontogeny and systematics of fishes*, p. 11-22. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Ketchen, K.S.**
1947 Studies on lemon sole development and egg production. Fish. Res. Board Can., Prog. Rep. Pac. Coast Stn. 73:68-70.
1956 Factors influencing the survival of lemon sole (*Parophrys vetulus*) in Hecate Strait, British Columbia. J. Fish. Res. Board Can. 13:647-694.
1961 Observations on the ecology of the Pacific cod (*Gadus macrocephalus*) in Canadian waters. J. Fish. Res. Board Can. 18:513-558.
- Khan, N.Y.**
1972 Comparative morphology and ecology of the pelagic larvae of nine Cottidae (Pisces) of the Northwest Atlantic and St. Lawrence Drainage. Ph.D. diss., Univ. Ottawa, Ottawa, Ontario, Canada, 234 p.
- Kido, K.**
1983 New and rare liparidid species from the Okhotsk and Bering Seas and their adjacent waters. Jpn. J. Ichthyol. 29:374-384.
1984 Occurrence of the liparidid fish, *Paraliparis pectoralis*, in the Bering Sea. Jpn. J. Ichthyol. 31:203-204.
1985 New and rare species of the genus *Careproctus* (Liparididae) from the Bering Sea. Jpn. J. Ichthyol. 32:6-7.
- Kido, K., and D. Kitagawa**
1986 Development of larvae and juveniles of *Rhinoliparis barbifer* (Liparididae). In Uyeno, T., et al. (eds.), *Indo-Pacific fish biology: Proceedings of the second international conference on Indo-Pacific fishes*, p. 697-702. Ichthyol. Soc. Jpn., Tokyo.
- Kimura, D.K., and A.R. Millikan**
1977 Assessment of the population of Pacific hake (*Merluccius productus*) in Puget Sound, Washington. Wash. Dep. Fish. Tech. Rep. 35, 46 p.
- Kobayashi, K.**
1961a Young of the wolf-fish, *Anarhichas orientalis* Pallas. Bull. Fac. Fish. Hokkaido Univ. 12:1-4 [in Jpn., Engl. abstr.].
1961b Larvae and young of the quill-fish, *Ptilichthys goodei*. Bull. Fac. Fish. Hokkaido Univ. 12:5-8 [in Jpn., Engl. abstr.].
1961c Larvae and young of the sand-lance, *Ammodytes hexapterus*. Bull. Fac. Fish. Hokkaido Univ. 12:111-120 [in Jpn., Engl. abstr.].
1962 Larvae of the smooth lumpsucker, *Aptocyclus ventricosus* (Pallas), with discussion on revision of the taxonomy of the species. Bull. Fac. Fish. Hokkaido Univ. 13:153-164.
- Kobayashi, K., M. Mikawa, and J. Ito**
1968 Descriptions of the young and one immature adult specimen of coster dory, *Alloctytus verrucosus* (Gilchrist) from the northern part of the Pacific. Bull. Fac. Fish. Hokkaido Univ. 19:1-6.
- Kramer, D.**
1960 Development of eggs and larvae of Pacific mackerel and distribution and abundance of larvae 1952-56. U.S. Fish Wildl. Serv., Fish. Bull. 60:393-438.
1970 Distributional atlas of fish eggs and larvae in the California Current region: Pacific sardine, *Sardinops caerulea* (Girard), 1951 through 1966. Calif. Coop. Oceanic Fish. Invest. Atlas 12, 277 p.
- Kramer, D., and E.H. Ahlstrom**
1968 Distributional atlas of fish larvae in the California Current region: Northern anchovy, *Engraulis mordax* Girard, 1951 through 1965. Calif. Coop. Oceanic Fish. Invest. Atlas 9, 282 p.
- Kramer, D.E., M.J. Kalin, E.G. Stevens, J.R. Thrailkill, and J.R. Zweifel**
1972 Collecting and processing data on fish eggs and larvae in the California Current region. NOAA Tech. Rep. NMFS Circ. 370, Natl. Oceanic Atmos. Adm., Natl. Mar. Fish. Serv., Seattle, WA 98115-0070, 38 p.
- Kyushin, K.**
1968 The embryonic and larval stages of *Hemitripteris villosus* (Pallas). Bull. Fac. Fish. Hokkaido Univ. 18:277-289.
- 1970** Embryonic development and larvae of *Gymnocanthus hertzensteini* Jordan and Starks. Jpn. J. Ichthyol. 17:74-79.
- LaRiviere, M.G., D.D. Jessup, and S.B. Mathews**
1981 Lingcod, *Ophiodon elongatus*, spawning and nesting in San Juan Channel, Washington. Calif. Fish Game 67:231-239.
- Laroche, W.A.**
In prep. Guide to larval and juvenile rockfishes (*Sebastes*) of North America, 311 p. Box 216, Enosburg Falls, VT 05450.
- Laroche, W.A., and S.L. Richardson**
1980 Development and occurrence of larvae and juveniles of the rockfishes *Sebastes flavidus* and *Sebastes melanops* (Scorpaenidae) off Oregon. Fish. Bull., U.S. 77:901-924.
1981 Development of larvae and juveniles of the rockfishes *Sebastes entomelas* and *S. zacentrus* (family Scorpaenidae) and occurrence off Oregon, with notes on head spines of *S. mystinus*, *S. flavidus*, and *S. melanops*. Fish. Bull., U.S. 79:231-258.
- Laroche, W.A., W.F. Smith-Vaniz, and S.L. Richardson**
1984 Carangidae: Development. In Moser, H.G., et al. (eds.), *Ontogeny and systematics of fishes*, p. 510-521. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Lasker, R., and P.E. Smith**
1977 Estimation of the effects of environmental variations on the eggs and larvae of northern anchovy. Calif. Coop. Oceanic Fish. Invest. Rep. 19:128-137.
- Lavenberg, R.J., G.E. McGowen, and R.E. Woodsum**
1984 Preservation and curation. In Moser, H.G., et al. (eds.), *Ontogeny and systematics of fishes*, p. 57-59. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Lea, R.N. and L.J. Dempster**
1982 Status and nomenclatural history of *Agonus vulsus* Jordan and Gilbert, 1880 (Pisces-family Agonidae). Calif. Fish Game 68: 249-252.
- Lea, R.N., and L.F. Quirolo**
1986 First record of *Hemitripteris bolini*, the bigmouth sculpin, from California waters. Calif. Fish Game 72:117-126.
- Lea, R.N., and R.H. Rosenblatt**
1987 Occurrence of the family Notacanthidae (Pisces) from the marine waters of California. Calif. Fish Game 73:51-53.
- Lee, J.U.**
1985 Studies on the fishery biology of the Atka mackerel *Pleurogrammus monopterygius* (Pallas) in the North Pacific Ocean. Bull. 34, Fish. Res. Dev. Agency, Pusan, Korea [in Korean].
- Leis, J.M.**
1984 Tetraodontoidei: Development. In Moser, H.G., et al. (eds.), *Ontogeny and systematics of fishes*, p. 447-449. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Leis, J.M., and D.S. Rennis**
1983 The larvae of Indo-Pacific coral reef fishes. New South Wales Univ. Press, Sydney, 269 p.
- Leis, J.M., and W.J. Richards**
1984 Acanthuroidei: Development and relationships. In Moser, H.G., et al. (eds.), *Ontogeny and systematics of fishes*, p. 547-551. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Limbaugh, C.**
1962 Life history and ecological notes on the tubenose, *Aulorhynchus flavidus*, a hemibranch fish of western North America. Copeia 1962:549-555.
- Loeb, V.J.**
1979 Larval fishes in the zooplankton community of the North Pacific central gyre. Mar. Biol. 53:73-191.
- MacCall, A.D., and G.D. Stauffer**
1983 Biology and fishery potential of jack mackerel (*Trachurus symmetricus*). Calif. Coop. Oceanic Fish. Invest. Rep. 24:46-56.
- MacGregor, J.S.**
1966 Fecundity of the Pacific hake, *Merluccius productus* (Ayres). Calif. Fish Game 52:111-116.

- 1986d** Early life history of the family Myctophidae in the ocean off southern Japan. In Ozawa, T. (ed.), Studies on the oceanic ichthyoplankton in the western North Pacific, p. 114-187. Kyushu Univ. Press, Fukuoka, Japan.
- 1986e** The larvae of the family Trichiuridae in the ocean off southern Japan. In Ozawa, T. (ed.), Studies on the oceanic ichthyoplankton in the western North Pacific, p. 290-300. Kyushu Univ. Press, Fukuoka, Japan.
- Ozawa, T., and S. Aono**
1986 Early ontogeny of melanostomiid fishes in the western North Pacific. In Ozawa, T. (ed.), Studies on the oceanic ichthyoplankton in the western North Pacific, p. 30-50. Kyushu Univ. Press, Fukuoka, Japan.
- Ozawa, T., and K. Oda**
1986 The larvae of the gonostomatid genus *Cyclothone* in the western North Pacific. In Ozawa, T. (ed.), Studies on the oceanic ichthyoplankton in the western North Pacific, p. 52-67. Kyushu Univ. Press, Fukuoka, Japan.
- Pacific Fisheries Management Council**
1978 Northern anchovy fishery: Final environmental impact statement and fishery management plan. Pac. Fish. Manage. Council., 526 S. Mill St., Portland, OR 97201.
- Patten, B.G.**
1973 Biological information on copper rockfish in Puget Sound, Washington. Trans. Am. Fish. Soc. 102:412-416.
1980 Short term thermal resistance of hexagrammid eggs and planktonic larvae from Puget Sound. Trans. Am. Fish. Soc. 109: 427-432.
- Paxton, J.R.**
1972 Osteology and relationships of the lanternfishes (Family, Myctophidae). Bull. Los Ang. Cty. Mus. Nat. Hist. Sci. 13, 81 p.
- Pearcy, W.G.**
1962 Egg masses and early developmental stages of the scorpaenid fish, *Sebastolobus*. J. Fish. Res. Board Can. 19:1169-1173.
- Peden, A.E.**
1978 A systematic revision of the hemilepidotinae fishes (Cottidae). Sysis 11:11-49.
- Peden, A.E., and G.W. Hughes**
1986 First records, confirmatory records, and range extensions of marine fishes of Canada's west coast. Can. Field-Nat. 100:1-9.
- Peden, A.E., W. Ostermann, and L.J. Pozar**
1985 Fishes observed at Canadian weather ship ocean station Papa [50 N, 145 W] with notes on the transpacific cruise of the CSS *Endeavor*. Brit. Col. Prov. Mus. Herit. Rec. 18, Min. Prov. Sec. and Gov. Serv., Victoria, B.C., Canada.
- Peppar, J.L.**
1965 Some features of the life history of the cockscomb prickleback, *Anoplarchus purpureus* Gill. M.S. thesis, Univ. Brit. Columbia, Vancouver, B.C., Canada, 159 p.
- Pertseva-Ostroumova, T.A.**
1961 The reproduction and development of far-eastern flounders. Tr. Inst. Okeanol. Akad. Nauk SSSR, 484 p. [Transl. avail. Fish. Res. Board Can., Pac. Biol. Stn., Nanaimo, B.C., Canada V9R 5K6, Transl. Ser. 856, 1967.]
1964 Some morphological characteristics of myctophid larvae (Myctophidae, Pisces). In Rass, T.S. (ed.), Fishes of the Pacific and Indian Oceans, Biology and distribution, p. 79-97. Tr. Inst. Okeanol. Akad. Nauk SSSR 73. [Transl. avail. U.S. Dep. Commer., Natl. Tech. Inf. Serv., Springfield, VA 22161, as TT65-5120.]
- Phillips, J.B.**
1959 A review of the lingcod, *Ophiodon elongatus*. Calif. Fish Game 45:19-27.
1964 Life history studies on ten species of rockfish (genus *Sebastes*). Calif. Dep. Fish Game, Fish Bull. 126, 70 p.
- Phillips, J.B., and S. Imamura**
1954 The sablefish fishery of California. Pac. Mar. Fish. Comm. Bull. 3:6-37.
- Pietsch, T.W.**
1976 Dimorphism, parasitism and sex: Reproductive strategies among deepsea ceratioid anglerfishes. Copeia 1976:781-793.
1978 Evolutionary relationships of the sea moths (Teleostei: Pegasidae) with a classification of gasterosteiform families. Copeia 1978:517-529.
1984 Lophiiformes: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 320-324. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
1986 Systematics and distribution of bathypelagic anglerfishes of the family Ceratiidae (order: Lophiiformes). Copeia 1986:479-493.
- Pietsch, T.W., and D. Grobecker**
1987 Frogfishes of the world. Systematics, zoogeography, and behavioral ecology. Stanford Univ. Press, Stanford, CA 94305, 420 p.
- Pillsbury, R.W.**
1957 Avoidance of poisonous eggs of the marine fish *Scorpaenichthys marmoratus* by predators. Copeia 1957:251-252.
- Powles, H., and D.F. Markle**
1984 Identification of larvae. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 31-33. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Prasad, R.R.**
1958 Reproduction in *Clevelandia ios* with an account of the embryonic and larval development. Proc. Natl. Inst. Sci. India 25B:12-30.
- Rass, T.S.**
1949 Composition of the ichthyoplankton of the Barents Sea. In Rass T.S., et al. (eds.), Material on the reproduction and development of fishes of the northern seas, p. 1-68. Tr. Vses. Nauchno-Issled. Inst. Morsk. Rybn. Khoz. Okeanogr. (VNIRO) 17 [in Russ.].
- Regan, C.T.**
1916 Larval and post-larval fishes. Nat. Hist. Rep. Br. Antarct. Terra Nova Exped. Zool. 1:125-156.
- Richardson, S.L.**
1977 Larval fishes in ocean waters off Yaquina Bay, Oregon: Abundance, distribution, and seasonality, January 1971 to August 1972. Publ. ORES-T-77-003, Oregon State Univ. Sea Grant Coll. Prog., Corvallis, OR 97331, 73 p.
1981a Current knowledge of larvae of sculpins (Pisces: Cottidae and allies) in Northeast Pacific genera with notes on intergeneric relationships. Fish. Bull., U.S. 79:103-121.
1981b Pelagic eggs and larvae of the deep sea sole, *Embassichthys bathybius* (Pisces: Pleuronectidae), with comments on generic affinities. Fish. Bull., U.S. 79:163-170.
- Richardson, S.L., and C. Bond**
1978 Two unusual cottoid fishes from the Northeast Pacific. Paper presented at Annu. Meet., Am. Soc. Ichthyol. Herpetol., Tempe, Arizona, 33 p. [Avail. A.C. Matarese, Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, 7600 Sand Point Way N.E., Seattle, WA 98115-0070.]
- Richardson, S.L., and D.A. Dehart**
1975 Records of larval, transforming, and adult specimens of quillfish, *Ptilichthys goodei*, from waters off Oregon. Fish. Bull., U.S. 73: 681-684.
- Richardson, S.L., and W.A. Laroche**
1979 Development and occurrence of larvae and juveniles of the rockfishes *Sebastes crameri*, *Sebastes pinniger*, and *Sebastes helvomaculatus* (family Scorpaenidae) off Oregon. Fish. Bull., U.S. 77:1-46.
- Richardson, S.L., and W.G. Pearcy**
1977 Coastal and oceanic fish larvae in an area of upwelling off Yaquina Bay, Oregon. Fish. Bull., U.S. 75:125-145.
- Richardson, S.L., and B.B. Washington**
1980 Guide to the identification of some sculpin larvae from marine and brackish waters off Oregon and adjacent areas of the Northeast Pacific. NOAA Tech. Rep. NMFS Circ. 430, Natl. Oceanic Atmos. Adm., Natl. Mar. Fish. Serv., Seattle, WA 98115-0070, 56 p.
- Richardson, S.L., J.R. Dunn, and N.A. Naplin**
1980 Eggs and larvae of butter sole, *Isopsetta isolepis* (Pleuronectidae), off Oregon and Washington. Fish. Bull., U.S. 78:401-417.

- Robertson, D.A.**
1977 Planktonic eggs of the lanternfish *Lampanyctodes hectoris* (family Myctophidae). Deep-Sea Res. 24:1-4.
- Robins, C.R., R.M. Bailey, C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea, and W.B. Scott**
1980 A list of common and scientific names of fishes from the United States and Canada. Spec. Publ. 12, Am. Fish. Soc., Bethesda, MD 20014, 174 p.
- Rofen, R.R.**
1966a Family Paralepididae. In Mead, G.W. (ed.), Fishes of the western North Atlantic, p. 205-461. Mem. 1 (pt. 5), Sears Found. Mar. Res., Yale Univ., New Haven, CT.
1966b Family Anotopteridae. In Mead, G.W. (ed.), Fishes of the western North Atlantic, p. 498-510. Mem. 1 (pt. 5), Sears Found. Mar. Res., Yale Univ., New Haven, CT.
1966c Family Omosudidae. In Mead, G.W. (ed.), Fishes of the western North Atlantic, p. 462-481. Mem. 1 (pt. 5), Sears Found. Mar. Res., Yale Univ., New Haven, CT.
- Rosenblatt, R.H., and R.R. Wilson**
1987 Cutlassfishes of the genus *Lepidopus* (Trichiuridae) with two new eastern Pacific species. Jpn. J. Ichthyol. 33:342-351.
- Roule, L., and F. Angel**
1930 Larves et alevins de poissons provenant des croisières du Prince Albert I de Monaco. Result. Camp. Sci. Prince Albert I 79:1-148 [in French].
- Rudomilov, O.I.**
1972 Fecundity of herring of the eastern part of the Bering Sea. Izv. Tikhookean. Nauchno-Issled. Inst. Rybn. Khoz. Okeanogr. (TINRO) 82:321-332 [in Russ., Engl. abstr.].
- Rugen, W.C., and A. C. Matarese**
1988 Spatial and temporal distribution and relative abundance of Pacific cod (*Gadus macrocephalus*) larvae in the western Gulf of Alaska. Proc. Rep. 88-18, Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070, 53 p.
- Ruple, D.**
1984 Gobioidae: Development. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 582-587. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Ryder, J.A.**
1987 On the development of osseous fishes, including marine and freshwater forms. Rep. U.S. Comm. Fish. 13(1885):489-603.
- Sakamoto, K.**
1984 Interrelationships of the family Pleuronectidae (Pisces: Pleuronectiformes). Mem. Fac. Fish., Hokkaido Univ. 31(1,2):95-215.
- Salveson, S.J.**
1976 Flathead sole (family Pleuronectidae). In Pereya, W.T., et al. (eds.), Demersal fish and shellfish resources in the eastern Bering Sea in the baseline year 1975, p. 497-510. Proc. Rep., Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070.
- Salveson, S.J., and M.S. Alton**
1976a Pollock (family Gadidae). In Pereya, W.T., et al. (eds.), Demersal fish and shellfish resources in the eastern Bering Sea in the baseline year 1975, p. 369-392. Proc. Rep., Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070.
1976b Yellowfin sole (family Pleuronectidae). In Pereya, W.T., et al. (eds.), Demersal fish and shellfish resources of the eastern Bering Sea in the baseline year 1975, p. 439-459. Proc. Rep., Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070.
- Sanzo, L.**
1931a Sottordinatae: Stomiatoidei. In Uova, larvae e stadi giovanili di Teleostei, p. 42-49. Fauna e Flora del Golfo di Napoli. Monogr. 38, 1064 p. + plates [in Ital.].
1931b Uova stadi embrionali e post-embionali di *Naucrates ductor* L. Mem. Res. Com. Talass. Ital. 185 [in Ital.].
- 1935** Uova, sviluppo, embrionale, stadi larvali, postlarvali e giovanili di Sternoptichidae e Stomiatiidae. Monogr. Res. Com. Talass. Ital. 2:123-180 [in Ital.].
- 1939** Nuovo contributo alla conoscenza dello sviluppo di *Myctophum rissoi* (Cocco). Atti. Accad. Gioenia Sci. Nat. Catania (6)3(25):1-8 [in Ital.].
- Saruwatari, T., K. Betsui, and M. Okiyama**
1987 Occurrence of the grunt sculpin (*Rhamphocottus richardsoni*) larvae from northern central Japan. Jpn. J. Ichthyol. 34:387-392.
- Sazonov, Yu. I.**
1981 *Talismania brachycephala* sp. N. (Alepocephalidae, Salmoniformes)—A new species from the northwestern Pacific Ocean. J. Ichthyol. 21(6):151-153.
- Scattergood, L.W., C.J. Sindermann, and B.E. Skud**
1959 Spawning of North American herring. Trans. Am. Fish. Soc. 88:164-168.
- Schaefer, M.B.**
1937 Notes on the spawning of the Pacific herring, *Clupea pallasii*. Copeia 1937:57.
- Schaefer, K.M.**
1980 Synopsis of biological data on the chub mackerel, *Scomber japonicus* Houttuyn, 1782, in the Pacific Ocean. In Bayliff, W.H. (ed.), Synopses of biological data on eight species of scombrids, p. 394-530. Spec. Rep. 2, Inter-Am. Trop. Tuna Comm., Scripps Inst. Oceanogr., La Jolla, CA 92093.
- Schmidt, J., and A. Strubberg**
1918 Mediterranean Bramidae and Trichuridae. Rep. Dan. Oceanogr. Exped. Mediterr. 2(A.6), 15 p.
- Schmitt, C.C., and B.E. Skud**
1978 Relation of fecundity to longterm changes in growth, abundance, and recruitment. Int. Pac. Halibut Comm. Sci. Rep. 66, 31 p.
- Schultz, L.P.**
1961 Revision of the marine silver hatchetfish (Family Sternoptichidae). Proc. U.S. Natl. Mus. 112(3449):587-649.
- Schultz, L.P., and A.C. DeLacy**
1932 The eggs and nesting habits of the crested blenny, *Anoplarchus*. Copeia 1932:143-147.
- Serobaba, I.I.**
1968 Spawning of the Alaskan pollock, *Theragra chalcogramma* (Pallas) in the northeastern Bering Sea. Probl. Ichthyol. 8(6):789-798.
- Shiino, S.M.**
1976 List of common names of fishes of the world, those prevailing among English-speaking nations. Sci. Rep. Shima Marineland 4, 206 p.
- Shiogaki, M.**
1982 Life history of the stichaeid fish *Opisthocentrus ocellatus*. Jpn. J. Ichthyol. 29:77-85.
- Shmidt, P.Y.**
1950 Fishes of the Sea of Okhotsk. Tr. Zool. Inst. Akad. Nauk SSSR 6:1-392. [Transl. by Isr. Prog. Sci. Transl., Jerusalem, Transl. 1263.]
- Shvetsov, F.C.**
1979 Reproduction of the flounder, *Lepidopsetta bilineata bilineata*, off the Okhotsk Sea coast near Paramoshiro and Shumushu Islands. J. Ichthyol. 19(5):61-62.
- Simenstad, C.A.**
1971 The feeding ecology of the rock greenling, *Hexagrammos lagocephalus* in the inshore waters of Amchitka Island, Alaska. M.S. thesis, Univ. Wash., Seattle, WA 98195, 131 p.
- Smith, D. G.**
1979 Guide to the leptocephali (Elopiformes, Anguilliformes, and Notacanthiformes). NOAA Tech. Rep. NMFS Circ. 424, Natl. Oceanic Atmos. Adm., Natl. Mar. Fish. Serv., Seattle, WA 98115-0070, 39 p.
1984 Elopiformes, Notacanthiformes and Anguilliformes: Relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 94-101. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.

- Smith, J.L.B.**
1964 Fishes of the family Pentacerotidae. Ichthyol. Bull. Rhodes Univ. 29:567-579.
- Smith, R.T.**
1936 Report on the Puget Sound otter trawl investigations. Wash. Dep. Fish. Biol. Rep. 36B, 61 p.
- Smoker, W., and W.G. Pearcy**
1970 Growth and reproduction of the lanternfish *Stenobrachius leucop-sarus*. J. Fish. Res. Board Can. 27:1265-1275.
- Soldatov, V.K., and G.J. Lindberg**
1930 A review of the fishes of the seas of the far east. Izv. Tikhoo-kean. Nauchno-Issled. Inst. Rybn. Khoz. Okeanogr. (TINRO) 5, 576 p. [in Russ.].
- St. Pierre, G.**
1984 Spawning locations and season for Pacific halibut. Int. Pac. Halibut Comm. Sci. Rep. 70, 46 p.
- Stahl-Johnson, K.L.**
1985 Descriptive characteristics of reared *Sebastes caurinus* and *S. auriculatus* larvae. In Kendall, A.W., Jr., and J.B. Marliave (eds.), Description of early life history stages of selected fishes: From the 3rd international symposium on the early life history of fishes and 8th annual larval fish conference, p. 65-76. Can. Tech. Rep. Fish. Aquat. Sci. 1359.
- Stein, D.L.**
1980a Description and occurrence of macrourid larvae and juveniles in the Northeast Pacific Ocean off Oregon, U.S.A. Deep-Sea Res. 27a:889-900.
1980b Aspects of reproduction of liparid fishes from the continental slope and abyssal plain off Oregon, with notes on growth. Copeia 1980:687-699.
- Stein, D.L., and C.E. Bond**
1985 Observations on the morphology, ecology, and behavior of *Bathylchnops exilis* Cohen. J. Fish Biol. 27:215-228.
- Stein, R.**
1972 Identification of some larval Pacific cottids. M.S. thesis, Humboldt State Univ., Arcata, CA 95521, 41 p.
1973 Description of laboratory-reared larvae of *Oligocottus maculosus* Girard (Pisces: Cottidae). Copeia 1973:373-377.
- Stepien, C.A.**
1986 Life history and larval development of the giant kelpfish, *Hetero-stichus rostratus* Girard, 1854. Fish. Bull., U.S. 84:809-826.
- Stevens, E.G., A.C. Matarese, and W.W. Watson**
1984 Ammodytoidei: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 574-575. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Steyskal, G.C.**
1980 The grammar of family-group names as exemplified by those of fishes. Proc. Biol. Soc. Wash. 93(1):168-177.
- Sulak, K.J., C.A. Wenner, G.R. Sedberry, and L.V. Guelpen**
1985 The life history and systematics of deep-sea lizard fishes, genus *Bathysaurus* (Synodontidae). Can. J. Zool. 63:623-642.
- Sumida, B.Y., E.H. Ahlstrom, and H.G. Moser**
1979 Early development of seven flatfish of the eastern North Pacific with heavily pigmented larvae (Pisces, Pleuronectiformes). Fish. Bull., U.S. 77:105-145.
- Takahura, T.**
1954 The behavior of the spawning pollock schools recorded by fish detector. Bull. Jpn. Soc. Sci. Fish. 20(1):10-12 [in Jpn., Engl. summ.].
- Tåning, A.V.**
1918 Mediterranean Scopelidae (*Saurus aulopus*, *Chlorophthalmus*, and *Myctophum*). Rep. Dan. Oceanogr. Exped. Mediterr. 2(A.7), 153 p.
- Templeman, W.**
1948 The life history of the caplin (*Mallotus villosus* O. F. Müller) in Newfoundland waters. Res. Bull. Div. Fish. Res. Newfoundland 17, 151 p.
- Thompson, J.A.**
1962 On the fecundity of Pacific cod (*Gadus macrocephalus*) from Hecate Strait, British Columbia. J. Fish. Res. Board Can. 19: 497-500.
- Thompson, W.F., and R. Van Cleave**
1936 Life history of the Pacific halibut. 2. Distribution and early life history. Rep. Int. Fish. Comm. 9, 184 p.
- Tighe, K.A., and M.J. Keene**
1984 Zeiformes: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 393-397. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Toole, C.L.**
1982 Widow rockfish. Marine Advisory Programs Newsletter, Calif. Sea Grant, Univ. Calif., Davis, CA 95616, p. 1-2.
- Trumble, R.J.**
1973 Distribution, relative abundance, and general biology of selected underutilized fishery resources of the eastern North Pacific Ocean. M.S. thesis, Univ. Wash., Seattle, WA 98195, 178 p.
- Tyler, J.C.**
1980 Osteology, phylogeny, and higher classification of the fishes of the order Plectognathi (Tetraodontiformes). NOAA Tech. Rep. NMFS Circ. 434. Natl. Oceanic Atmos. Adm., Natl. Mar. Fish. Serv., Seattle, WA 98115-0070, 422 p.
- Uchida, K., S. Imai, S. Mito, S. Fujita, M. Ueno, Y. Shojima, T. Senta, M. Tahuka, and Y. Dotsu**
1958 Studies on the eggs, larvae, and juveniles of Japanese fishes. Series 1. Second Laboratory of Fisheries Biology, Fish. Dep., Fac. Agric., Kyushu Univ., Fukuoka, Japan [in Jpn.].
- Waldron, K.D.**
1968 Early larvae of the canary rockfish, *Sebastes pinniger*. J. Fish. Res. Board Can. 25:801-803.
- Wales, L.H.**
1952 Life history of the blue rockfish, *Sebastes mystinus*. Calif. Fish Game 38:485-498.
- Walters, G.E.**
1984 Ecological aspects of larval and juvenile Pacific cod (*Gadus macrocephalus*), walleye pollock (*Theragra chalcogramma*), and Pacific tomcod (*Microgadus proximus*) in Port Townsend, Wash-ington. M.S. thesis, Univ. Wash., Seattle, WA 98195, 129 p.
- Wang, J.C.**
1981 Taxonomy of the early life stages of fishes: Fishes of the Sacra-mento-San Joaquin estuary and Moss Landing harbor-Elkhorn slough, California. Ecological Analysts, Inc., Concord, CA 94520, 168 p.
1986 Fishes of the Sacramento-San Joaquin estuary and adjacent waters, California: A guide to the early life histories. Interagency Ecological Study Program for the Sacramento-San Joaquin Estuary. Tech. Rep. 9, 602 p. [Avail. Ecological Analysts, Inc., 2150 John Glenn Drive, Concord, CA 94520.]
- Washington, B.B.**
1981 Identification and systematics of larvae of *Artemis*, *Clinocottus*, and *Oligocottus* (Scorpaeniformes: Cottidae). M.S. thesis, Oregon State Univ., Corvallis, OR 97331, 205 p.
1986 Systematic relationships and ontogeny of the sculpins *Artemis*, *Clinocottus*, and *Oligocottus* (Cottidae: Scorpaeniformes). Proc. Calif. Acad. Sci. 44(9):157-224.
- Washington, B.B., W.N. Eschmeyer, and K.M. Howe**
1984a Scorpaeniformes: Relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 438-447. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Washington, B.B., H.G. Moser, W.A. Laroche, and W.J. Richards**
1984b Scorpaeniformes: Development. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 405-427. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Washington, P.M., R. Gowan, and D.H. Ito**
1978 A biological report on eight species of rockfish (*Sebastes* spp.) from Puget Sound, Washington. Proc. Rep., Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070, 50 p.

- Watson, W.W.**
 1982 Development of eggs and larvae of the white croaker, *Genyonemus lineatus* Ayres (Pisces: Sciaenidae) off the southern California coast. Fish. Bull., U.S. 80:403-417.
- Webber, R.A., and M.S. Alton**
 1976 Pacific halibut (family Pleuronectidae). In Pereya, W.T., et al. (eds.), Demersal fish and shellfish resources of the eastern Bering Sea in the baseline year 1975, p. 511-522. Proc. Rep., Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070.
- Weitzman, S.H.**
 1974 Osteology and evolutionary relationships of the Sternoptychidae with a new classification of stomiatoid families. Bull. Am. Mus. Nat. Hist. 153:327-478.
- Wenner, C.A.**
 1978 Making a living on the continental slope and in the deep sea: Life history of some dominant fishes of the Norfolk Canyon area. Ph.D. diss., Coll. William and Mary, Williamsburg, VA 23186, 294 leaves, 105 leaves of plates.
- Westrheim, S.J.**
 1975 Reproduction, maturation, and identification of larvae of some *Sebastes* (Scorpaenidae) species in the Northeast Pacific Ocean. J. Fish. Res. Board Can. 32:2399-2411.
- Westrheim, S.J., and A.R. Morgan**
 1963 Results from tagging a spawning stock of Dover sole, *Microstomus pacificus*. Bull. Pac. Mar. Fish. Comm. 6:13-21.
- Westrheim, S.J., W.R. Harling, and D. Davenport**
 1968a Preliminary report on maturity, spawning season, and larval identification of rockfishes (*Sebastes*) collected off British Columbia in 1967. Fish. Res. Board Can., Manuscr. Rep. Ser. 951, 23 p.
- Westrheim, S.J., W.R. Harling, D. Davenport, and M.S. Smith**
 1968b Preliminary report on maturity, spawning season, and larval identification of rockfishes (*Sebastes*) collected off British Columbia in 1968. Fish. Res. Board Can., Manuscr. Rep. Ser. 1005, 28 p.
- White, B.N., R.J. Lavenberg, and G.E. McGowen**
 1984 Atheriniformes: Development and relationships. In Moser, H.G., et al. (eds.), Ontogeny and systematics of fishes, p. 355-361. Spec. Publ. 1, Am. Soc. Ichthyol. Herpetol. Allen Press, Lawrence, KS.
- Wiley, J.W.**
 1973 Life history of the western North American goby *Coryphopterus nicholsi* (Bean). Trans. San Diego Soc. Nat. Hist. 17(14):187-208.
- Wilimovsky, N.J., A. Peden, and J. Peppar**
 1967 Systematics of six demersal fishes of the North Pacific Ocean. Fish. Res. Board Can. Tech. Rep. 34, 95 p.
- Wingert, R.C.**
 1974 Comparative reproductive cycles and growth in two species of *Xiphister* (Pisces, Stichaeidae), from San Simeon, California. M.A. thesis, Calif. State Univ., Fullerton, CA 92631, 91 p.
- Wisner, R.L.**
 1974 The taxonomy and distribution of lanternfishes (family Myctophidae) of the eastern Pacific Ocean. NORDA Rep.-3, Navy Ocean Research and Development Activity, Bay St. Louis, MS, 229 p.
- Wolotira, R.J.**
 1985 Saffron cod (*Eleginus gracilis*) in western Alaska: The resource and its potential. Tech. Memo. NMFS F/NWC-79, Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070, 119 p.
- Wolotira, R.J., T.M. Sample, and M. Morin, Jr.**
 1977 Demersal fish and shellfish resources of Norton Sound, the southeastern Chukchi Sea, and adjacent waters in the baseline year 1976. Proc. Rep., Northwest Alaska Fish. Cent., Natl. Mar. Fish. Serv., NOAA, Seattle, WA 98115-0070, 292 p.
- Wooster, W.S.**
 1983 From year to year: Interannual variability of the environment and fisheries of the Gulf of Alaska and the eastern Bering Sea. Rep. WSG-WO 83-3, Wash. Sea Grant Prog., Univ. Wash., Seattle, WA 98195, 208 p.
- Wyllie Echeverria, T.**
 1987 Thirty-four species of California rockfishes: Maturity and seasonality of reproduction. Fish. Bull., U.S. 85:229-250.
- Yatsu, A.**
 1985 Phylogeny of the family Pholididae (Blennioidei) with a redescription of *Pholis scopoli*. Jpn. J. Ichthyol. 32:273-282.
 1986 Phylogeny and zoogeography of the subfamilies Xiphisterinae and Cebidichthyinae (Blennioidei, Stichaeidae). In Uyeno, T., et al. (eds.), Indo-Pacific fish biology: Proceedings of the second international conference on Indo-Pacific fishes, p. 663-678. Ichthyol. Soc. Jpn., Tokyo.
- Yusa, T.**
 1957 Eggs and larvae of flatfishes in the coastal waters of Hokkaido. I. Embryonic development of the starry flounder *Platichthys stellatus* (Pallas). Bull. Hokkaido Reg. Fish. Res. Lab. 15:1-14.
- Zama, A., M. Asai, and F. Yasuda**
 1977a Records of the pelagic armorhead, *Pentaceros richardsoni*, from Hachijo Island and the Ogasawara Islands. Jpn. J. Ichthyol. 24: 57-60.
 1977b Changes with growth in bony cranial projections and colour patterns in the Japanese boarfish, *Pentaceros japonicus*. Jpn. J. Ichthyol. 24:26-34.

Index to scientific names

- Acantholiparis* 474
Acantholiparis caecus 474
Acantholiparis opercularis 474
Acantholumpenus mackayi 507
Acanthopsetta 573-575
Acanthopsetta nadeshnyi 567, 574, 575
Acanthuroidei 18, 556-557
Agonidae 19, 267, 456, 457, 458-463
Agonidae A 4, 456, 457
Agonomalus mozinoi 4, 456, 458, 459
Agonopsis vulsa 456
Agonus acipenserinus 456
Agonus decagonus 456
Albatrossia pectoralis 200-201
Alectridium aurantiacum 507
Alectrini 504, 505, 507
Alepisauridae 19, 125, 142-143
Alepisaurus ferox 142, 143
Alepocephalidae 19, 51, 76
Alepocephalus tenebrosus 76
Alloctytus sp. 257-258
Allolumpenus hypochromus 507
Allosmerus elongatus 79
Ammodytes hexapterus 4, 27, 540, 541
Ammodytidae 19, 26, 27, 483, 540-541
Ammodytoidei 18, 540-541
Anarhichantidae 19, 483, 524-527
Anarhichas orientalis 4, 524, 525
Anarrhichthys ocellatus 524, 526, 527
Anguilliformes 18, 31-41
Anisarchus medius 507
Anoplagonus inermis 456
Anoplarchus insignis 507
Anoplarchus purpureus 4, 27, 505, 507, 512, 513, 514
Anoplogaster cornuta 250, 251
Anoplogastridae 19, 249-251
Anoplopoma fimbria 4, 20, 24, 338, 339
Anoplopomatidae 19, 267, 338-339
Anoplopomatoidei 18, 267, 338-339
Anotopteridae 19, 125, 140-141
Anotopterus pharao 140, 141
Antimora 182
Antimora microlepis 182
Aphanopus carbo 551
Aphyonidae 19, 211, 217
Apodichthys flavidus 4, 522, 523
Aptocyclus ventricosus 465-466, 467
Archaulus 361
Arctoscopus japonicus 532
Argentina sialis 52, 53
Argentinidae 19, 51-55
Argentinoidei 18, 51-77
Argyropelecus 84, 85, 92-101
Argyropelecus affinis 93-94, 95, 100
Argyropelecus hemigymnus 92-93, 96, 97, 100
Argyropelecus lychnus 93, 98, 99
Argyropelecus sladeni 93, 98, 100, 101
Aristostomias 106, 118-119
Aristostomias scintillans 106, 118, 119
Arteidiellichthys 361
Arteidielliscus 361
Arteidiellus 361
Arteidius 361-362, 384-385, 414-421
Arteidius corallinus 420
Arteidius fenestralis 414, 415, 416, 418, 420
Arteidius harringtoni 363, 416, 417
Arteidius lateralis 414, 416, 418, 419, 420
Arteidius meanyi 361-362, 384, 385
Arteidius notospilotus 420
Arteidius spp. 414, 416, 418, 420
Arteidius 3 414, 416, 418, 420, 421
Ascelichthys 361-363, 386-387
Ascelichthys rhodorus 363, 386, 387
Asemichthys 361
Aspidophoroides bartoni 456
Aspidophoroides olriki 456
Atheresthes 573, 576-579
Atheresthes evermanni 567, 576, 577
Atheresthes stomias 5, 567, 576, 578, 579
Atherinidae 19, 235, 238-241
Atheriniformes 18, 235, 238-241
Atherinomorpha 235-241
Atherinops affinis 238, 239, 240
Atherinopsis californiensis 238, 240, 241
Aulopiformes 18, 125-143
Aulorhynchidae 19, 261-263
Aulorhynchus flavidus 262, 263
Avocettina infans 33, 38, 39, 40
Barathronus pacificus 217
Barbourisia rufa 256
Barbourisiidae 19, 249, 256
Bassozetus sp. 212
Bathophilus 106, 110-111
Bathophilus flemingi 106, 110, 111
Bathyagonus alascanus 456
Bathyagonus infraspinatus 456
Bathyagonus nigripinnis 456
Bathyagonus pentacanthus 456
Bathylaco nigricans 76
Bathylagidae 19, 51, 56-69
Bathylagus bericoides 56, 57, 60
Bathylagus milleri 3, 58, 59, 66
Bathylagus ochotensis 3, 56, 60, 61
Bathylagus pacificus 3, 56, 60, 62, 63
Bathylagus wesethi 58, 64, 65
Bathylachnops exilis 70, 71, 72, 74
Bathymaster A 497
Bathymaster caeruleofasciatus 496

Bathymaster leurolepis 496
Bathymaster signatus 496
 Bathymasteridae 19, 26, 27, 483, 496-497
 Bathysauridae 19, 125, 130-131
Bathysaurus mollis 130, 131
 Beloniformes 18, 235-237
Benthalbella dentata 126, 127
Benthalbella linguidens 126, 127
Benthodesmus elongatus 551-552, 553
Benthodesmus spp. 552, 553
Benthodesmus tenuis 551-552
Bertella idiomorpha 223
 Beryciformes 18, 249-256
 Blennioidei 18, 535-537
Blepsias 20, 361, 364, 448-451
Blepsias bilobus 4, 448, 449, 450
Blepsias cirrhosus 4, 22, 448, 450, 451
Boreogadus saida 3, 189-190, 191
Bothragonus swani 456, 460, 461, 462
Bothrocara brunneum 498
Bothrocara hollandi 498, 499
Bothrocara molle 498
Bothrocara pusillum 498
Bothrocara remigerum 498
Brama japonicus 488, 489
 Bramidae 19, 483, 488-489
Brosomphycis marginata 3, 217-218, 219
Bryozoichthys lysimus 507
Bryozoichthys marjorius 507
 Bythitidae 19, 211, 217-219
 Bythitoidei 211, 217-219
 Carangidae 19, 483-487
Careproctus 470, 472, 474
Careproctus abbreviatus 474
Careproctus attenuatus 474
Careproctus bowersianus 474
Careproctus cameliae 474
Careproctus canus 474
Careproctus colletti 474
Careproctus cypselurus 474
Careproctus ectenes 474
Careproctus filamentosus 474
Careproctus furcellus 474
Careproctus gilberti 474
Careproctus longifilis 474
Careproctus melanurus 474
Careproctus microstomus 474
Careproctus mollis 474
Careproctus opisthotremus 474
Careproctus oregonensis 474
Careproctus ostentum 474
Careproctus ovigerum 474
Careproctus pellucidus 474
Careproctus phasma 474
Careproctus pycnosoma 474
Careproctus rastrinus 474
Careproctus scottae 474
Careproctus simus 474
Careproctus spectrum 474
Careproctus zachirus 474
 Caristiidae 19, 490, 491
Caristius macropus 490
Cataetyx rubrirostris 217
Cebidichthys violaceus 507
 Centrolophidae 19, 483, 558-559
Ceratias holboelli 222-223
Ceratias sp. 222-223
 Ceratiidae 221-223
 Ceratioidei 221-227
Ceratiscopelus 146-147, 160-161
Ceratiscopelus townsendi 146, 160, 161
 Cetomimidae 249, 256
 Cetomimoidei 249, 256
Cetomimus 256
Cetostoma regani 256
Chaenophryne longiceps 223-224, 225
Chaenophryne melanorhabus 223
 Chauliodontidae 19, 83, 106-109
Chauliodus 106-109
Chauliodus macouni 106, 108, 109
Chilara taylori 212, 214, 215
 Chirolophini 500, 501, 507
Chirolophis decoratus 507
Chirolophis nugator 507
Chirolophis snyderi 507
Chirolophis tarsodes 507
Chitonotus 361-362, 388-389
Chitonotus pugetensis 4, 388, 389
Citharichthys sordidus 5, 567-568, 569
Citharichthys stigmaeus 567, 570, 571
Clevelandia ios 4, 543-544, 545, 548
Clidoderma asperrimum 567
 Clinidae 19, 483, 535-537
Clinocottus 361-362, 422-428
Clinocottus acuticeps 360, 422, 423
Clinocottus embryum 424, 425
Clinocottus globiceps 426, 427
Clinocottus recalvus 426, 428, 429
Clupea pallasi 3, 18, 27, 44, 45, 46, 48
 Clupeidae 19, 26, 27, 43-47
 Clupeiformes 18, 43-49
Cololabis saira 3, 20, 25, 236, 237
 Congridae 19, 31-33, 36, 37
Coryphaenoides acrolepis 200-202, 203
Coryphaenoides armatus 200-201
Coryphaenoides cinereus 200-201
Coryphaenoides filifer 200-201, 204, 205
Coryphaenoides leptolepis 200-201, 206, 207
Coryphaenoides liocephalus 200
Coryphaenoides longifilis 200
Coryphaenoides yaquinae 200
Coryphopterus nicholsi 4, 543-544, 546, 547, 548
 Cottidae 18-19, 267, 360-362, 363, 364, 365, 366-455
 Cottoid A 436, 438, 439

Cottoidei 18, 267, 360-455
Cottus 361, 364, 444-445
Cottus aleuticus 444
Cottus asper 444, 445
 Cryptacanthodidae 19, 483, 520, 521
Crystallichthys cyclospilus 474
Crystallichthys mirabilis 474
 Cyclopteridae 4, 18-19, 267, 465-481
 Cyclopterinae 465-469
Cyclopteropsis phrynoides 465
Cyclothone 84, 86-88, 89, 90
Cyclothone acclinidens 87-88, 89
Cyclothone atraria 87-88, 89
Cyclothone pallida 87-88, 89
Cyclothone pseudopallida 87-88, 89
Cyclothone signata 87-88, 89
Cyema atrum 33-34, 35
 Cyematidae 19, 31-34, 35
 Cynoglossidae 19, 565, 567, 624-625
Danaphos 84, 86, 92, 104-105
Danaphos oculatus 3, 93, 104, 105
Dasycottus 361, 364, 440-441
Dasycottus setiger 440, 441
Delolepis gigantea 4, 20, 25, 520, 521
Derepodichthys alepidotus 498
 Diaphini 146, 176-177
Diaphus 146-147, 176-177
Diaphus theta 3, 146, 172, 176, 177
Dicrolene filamentosa 212
Dolichopteryx sp. 72, 73, 74
Dorsadena 146-147
Dorsadena yaquinae 146
Elassodiscus caudatus 474
Elassodiscus tremebundus 474
Electrona 146-149
Electrona rissoi 146, 148, 149
 Electronini 146-153
Eleginus gracilis 189, 192, 193
 Elopomorpha 31-41
Embassichthys 573, 580-581
Embassichthys bathybius 212, 567, 580, 581, 622
 Engraulididae 19, 26, 27, 43, 48-49
Engraulis mordax 27, 44, 46, 48, 49
Enophrys 361-362, 390-391
Enophrys bison 390, 391
Enophrys diceraus 390
Enophrys lucasi 390
Eopsetta 567, 573, 582-583
Eopsetta jordani 567, 582, 583
Ericara salmoneum 76
Erilepis zonifer 338
Eumesogrammus praecisus 507
Eumicrotremus andriashevi 465
Eumicrotremus barbatus 465
Eumicrotremus birulai 465
Eumicrotremus gyrynops 465
Eumicrotremus orbis 465, 468, 469
Eumicrotremus soldatovi 465
Eumicrotremus taranetzi 465
Eurymen 361
 Eurypharyngidae 19, 32-33, 36, 37
Eurypharynx pelecanooides 33, 37
Eustomias 106, 112, 113
Eustomias sp. 106, 112, 113
 Gadidae 18-19, 181, 189-199
 Gadiformes 18, 181-209
Gadus macrocephalus 3, 189, 194, 195, 196, 198
 Gasterosteidae 261
 Gasterosteiformes 18, 261-265
Genyonemus lineatus 492, 493
Gibbonsia metzi 535
Gibbonsia montereyensis 535
Gilbertidia 361, 364, 435-436
Gilbertidia sigalutes 434, 435, 436
Glyptocephalus 567, 573, 584-587
Glyptocephalus stelleri 567, 580, 584, 585
Glyptocephalus zachirus 5, 567, 580, 584, 586, 587, 622
 Gobiesocidae 19, 229-233
 Gobiesociformes 18, 229-233
Gobiesox maeandricus 3, 230, 231, 232
 Gobiidae 19, 483, 543-549
 Gobioidei 18, 543-549
 Gonichthyini 146, 156-159
Gonostoma 84, 86, 88, 90
Gonostoma atlanticum 87-88, 90, 91
Gonostoma gracile 84, 87-88, 90, 91
 Gonostomatidae 19, 83, 87-91
 Gonostomatoidei 86-105
Gymnelis hemifasciatus 498
Gymnelis popovi 498
Gymnelis viridis 498, 499
Gymnocanthus 361-362, 392, 393, 394-395
Gymnocanthus A 392, 393
Gymnocanthus detruscus 392
Gymnocanthus galeatus 392
Gymnocanthus pistilliger 392
Gymnocanthus tricuspis 392, 394, 395
Gymnoclinus cristulatus 507
 Gymnoscopelini 146, 178-179
Gyrinichthys minytrems 474
Gyrinomimus sp. 256
Hemilepidotus 361-362, 368-381
Hemilepidotus gilberti 368, 370, 372, 373
Hemilepidotus hemilepidotus 368, 369, 370, 371, 374, 375, 376, 380
Hemilepidotus jordani 4, 368, 369, 370, 371, 376, 377, 380
Hemilepidotus papilio 368, 370
Hemilepidotus spinosus 21, 363, 368, 370, 371, 378, 379
Hemilepidotus spp. 20, 369, 370, 382
Hemilepidotus zapus 368, 369, 370, 371, 380, 381
Hemitripterus 361, 364, 452-453
Hemitripterus bolini 452
Hemitripterus villosus 20, 23, 365, 452, 453

Heterostichus rostratus 535-536, 537
 Hexagrammidae 19, 267, 341-359
 Hexagrammoidei 18, 267, 341-359
Hexagrammos 20, 341, 350, 351, 352-359
Hexagrammos decagrammus 341, 350, 351, 352, 353
Hexagrammos lagocephalus 341, 350, 351, 354, 355
Hexagrammos octogrammus 24, 341, 350, 351, 356, 357
Hexagrammos stelleri 341, 350, 351, 358, 359
Hippoglossoides 573, 588-591
Hippoglossoides elassodon 15, 18, 567, 588, 589, 590
Hippoglossoides robustus 567, 590, 591
Hippoglossus 573, 592-593
Hippoglossus stenolepis 5, 567, 592, 593, 622
Histiobranchus bathybius 33-34
Holcomycteronus profundissimus 212
Holtbyrnia innesi 77
Holtbyrnia latifrons 77
Hypomesus pretiosus 79
Hypsagonus quadricornus 456, 457
Icelinus 361-362, 384, 388, 396, 397
Icelinus A 396, 397
Icelinus B 396, 397
Icelinus borealis 396
Icelinus burchami 396
Icelinus filamentosus 396
Icelinus fimbriatus 396
Icelinus tenuis 396
Icelus 361-362, 399
Icelus canaliculatus 399
Icelus euryops 399
Icelus scutiger 399
Icelus spatula 399
Icelus spiniger 399
Icelus uncinialis 399
Icichthys lockingtoni 558, 559, 560
 Icosteidae 19, 483, 538-539
 Icosteoidi 18, 538-539
Icosteus aenigmaticus 538, 539
 Idiacanthidae 19, 83, 106, 120-123
Idiacanthus 106, 120-123
Idiacanthus antrostomus 106, 120, 121, 122
Idiacanthus fasciola 120, 122, 123
Inopsetta ischyra 567
Isopsetta 567, 573, 594-595
Isopsetta isolepis 567, 594, 595
Jordania 361
Krusensterniella pavlovskii 498
Laemonema 182
Laemonema longipes 182
Lampadena 146-147, 162-163
Lampadena urophaos 146, 162, 163
 Lampanyctinae 146-147, 160-179
 Lampanyctini 146, 160-175
Lampanyctus 147, 165-169
Lampanyctus fernae 146, 165
Lampanyctus jordani 146, 165
Lampanyctus regalis 3, 146, 165-166, 167, 168
Lampanyctus ritteri 146, 165, 168, 169
 Lampridae 19, 243-245
 Lampriformes 18, 243-247
Lampris guttatus 244, 245
Lepidogobius lepidus 4, 543-544, 548, 549
Lepidopsetta 567, 573, 596-599, 600-601
Lepidopsetta bilineata 5, 18, 567, 596, 597, 599, 600-601
Lepidopsetta 2 5, 599, 600-601
Lepidopus fitchi 551
Leptochilichthys agassizi 76
Leptocottus 361, 364, 446-447
Leptocottus armatus 365, 446, 447
Lestidiops ringens 134, 135, 136, 138
Lethotremus muticus 465
Leuroglossus schmidti 66, 67, 68
Leuroglossus stilbius 66-68, 69
Limanda 567, 573, 602-603
Limanda aspera 567, 602, 603
Limanda proboscidea 567, 602, 603
Limanda sakhalinensis 602
Liopsetta 567, 573, 604-605
Liopsetta glacialis 567, 604, 605
 Liparidinae 465, 470, 471-473, 474-481
Liparis 470, 471, 474-475
Liparis bristolensis 474
Liparis callyodon 474
Liparis catharus 475
Liparis cyclopus 475
Liparis dennyi 475
Liparis florum 475
Liparis fucensis 475
Liparis gibbus 475
Liparis grebnitzki 475
Liparis mednius 475
Liparis megacephalus 475
Liparis micraspidophorus 475
Liparis mucosus 475
Liparis ochotensis 475
Liparis pulchellus 475
Liparis rutteri 475
Liparis tunicatus 475
Lipariscus nanus 475
 Lophiiformes 18, 221-227
Loweina 147, 156-157
Loweina rara 146, 156, 157, 158
Lumpenella longirostris 503, 507
 Lumpeninae 502-505, 507
 Lumpenini 502, 503, 507
Lumpenus fabricii 507
Lumpenus maculatus 503, 507
Lumpenus medius 507
Lumpenus sagitta 4, 503, 507
 Luvaridae 19, 483, 556-557
Luvarus imperialis 556, 557
Lycenchelys altus 498
Lycenchelys camchaticus 498
Lycenchelys crotalinus 498

Lycenchelys hippopotamus 498
Lycenchelys jordani 498
Lycenchelys longirostris 498
Lycenchelys microporus 498
Lycenchelys pliciferus 498
Lycenchelys rassi 498
Lycenchelys ratmanovi 498
Lycenchelys roseus 498
Lycenchelys volki 498
Lycodapus derjugini 498
Lycodapus dermatinus 498
Lycodapus endemoscotus 498
Lycodapus fierasfer 498
Lycodapus leptus 498
Lycodapus mandibularis 498
Lycodapus pachysoma 498
Lycodapus parviceps 498
Lycodapus poecilus 498
Lycodapus psarosomatus 498
Lycodes brevipes 498
Lycodes concolor 498
Lycodes cortezianus 498
Lycodes diapterus 498
Lycodes mucosus 498
Lycodes pacifica 498
Lycodes palearis 498
Lycodes raridens 498
Lycodes turneri 498
Lyconectes aleutensis 4, 20, 25, 520, 521
Lyconema barbatum 498
Lyopsetta 573, 606-607
Lyopsetta exilis 567, 586, 606, 607, 608
Macropinna microstoma 3, 72, 74, 75
Macrouridae 18-19, 181, 200-209
Malacocottus 361, 364, 442-443
Malacocottus kinkaidi 442
Malacocottus zonurus 365, 442, 443
Malacosteidae 19, 83, 106, 118-119
Mallotus villosus 26, 27, 79-80, 81
Maulisia argipalla 77
Megalocottus 361
Melamphaes 252, 253, 254-255
Melamphaes lugubris 252, 253, 254, 255, 336
Melamphaes parvus 254
Melamphidae 19, 249, 252, 253, 254-255
Melanonidae 19, 181, 184-185
Melanonus zugmayeri 3, 184, 185
Melanostigma pammelas 498
Melanostomiidae 19, 83, 106, 110-117
Merlucciidae 19, 181, 186-187
Merluccius productus 1, 186, 187, 486
Microcottus 361
Microgadus proximus 189, 196, 197
Microstomus 573, 608-609
Microstomus pacificus 5, 567, 586, 608, 609
Mola mola 18, 628, 629
Molidae 19, 627-629
Moridae 19, 181-182, 183
Myctophidae 19, 125, 146-179
Myctophiformes 18, 125, 144-179
Myctophinae 146-159
Myctophini 146, 154-155
Myoxocephalus 361-362, 400, 401, 402-403
Myoxocephalus axillaris 400
Myoxocephalus B 400, 401
Myoxocephalus brandti 400
Myoxocephalus G 20, 21, 363, 400, 401
Myoxocephalus jaok 400
Myoxocephalus niger 400
Myoxocephalus polyacanthocephalus 400, 401
Myoxocephalus quadricornis 400
Myoxocephalus scorpioides 400
Myoxocephalus scorpius 400, 402, 403
Myoxocephalus stelleri 400
Myoxocephalus verrucosus 400, 402
Nalbantichthys elongatus 498
Nansenia candida 3, 52, 54, 55
Narctes stomias 76
Naucrates ductor 484, 485
Nautichthys 361, 364, 454-455
Nautichthys oculo-fasciatus 4, 20, 23, 454, 455
Nautichthys pribilovius 454
Nautichthys robustus 454
Nectoliparis pelagicus 4, 470, 475-476, 477
Nemichthyidae 19, 31-33, 38-41
Nemichthys larseni 33, 40
Nemichthys scolopaceus 33, 40, 41
Neoscopelidae 19, 125, 144, 145
Neoscopelus macrolepidotus 144-145
Nettastomatidae 19, 31-33, 36, 37
Nezumia stelgidolepis 200, 208-209
Notacanthidae 31-34, 35
Notacanthiformes 18, 31-35
Notacanthus chemnitzi 33-34
Notolepis rissoi 134, 136, 137
Notoscopelus 146-147, 178-179
Notoscopelus japonicus 178
Notoscopelus resplendens 146, 178, 179
Notosudidae 19, 125, 128-129
Ocella dodecaedron 456
Ocella impi 456
Ocella verrucosa 456, 457
Odontoliparis ferox 475
Odontopyxis trispinosa 456
Oligocottus 361-362, 430-433
Oligocottus maculosus 430, 431, 432
Oligocottus rimensis 430
Oligocottus snyderi 430, 432, 433
Oneirodes bulbosus 223, 226-227
Oneirodes thompsoni 223
Oneirodidae 19, 221-227
Opaeophacus acrogeneius 498
Ophidiidae 3, 19, 211-212, 213, 214-215
Ophidiiformes 18, 211-219

Ophidioidei 211-215
Ophiodon elongatus 20, 24, 341, 346, 347
 Opisthocentrini 504, 505, 507
Opisthocentrus ocellatus 505, 507-508, 509
 Opisthoproctidae 19, 51, 70-75
Opostomias 106, 114-115
Opostomias mitsuui 106, 114, 115
 Oreosomatidae 19, 257-259
 Osmeridae 19, 26, 27, 51, 79-81
Osmerus mordax 79
Osteodiscus cascadiae 475
Oxylebius pictus 341-342, 343, 344
Pachycara bulbiceps 498
Pallasina barbata 456
 Paralepididae 19, 125, 134-139
Paralepis atlantica 134, 138, 139
 Paralichthyidae 19, 565, 567-571
Paraliparis 470, 472, 475, 478-479
Paraliparis cephalus 475
Paraliparis dactylosus 475
Paraliparis deani 470, 475, 478
Paraliparis holomelas 475
Paraliparis latifrons 475
Paraliparis megalopsis 475
Paraliparis melanobranchus 475
Paraliparis mento 475
Paraliparis paucidens 475
Paraliparis pectoralis 475
Paraliparis rosaceus 475
Paraliparis sp. 4, 470, 478, 479
Paraliparis ulochir 475
Paricelinus 361-362, 404-405
Paricelinus hopliticus 404, 405
Parophrys 567, 573, 610-611
Parophrys vetulus 5, 567, 594, 610, 611
Parvilux 147, 170-171
Parvilux ingens 146, 170, 171
Pelagocyclus vitiazii 465
Pellisulus eubranchnus 77
 Pentacerotidae 19, 483, 494, 495
Peprilus simillimus 562, 563
 Perciformes 18, 483-563
Percis japonicus 456
 Percoidei 18, 484-495
Phallocottus 361
 Pholididae 19, 26, 27, 483, 522, 523
Pholis sp. 4, 27, 563
Pholis clemensi 522
Pholis dolichogaster 522
Pholis fasciata 522
Pholis gilli 522
Pholis laeta 522
Pholis ornata 522
Pholis schultzi 522
Phytichthys chirus 507, 512, 514, 515
Platichthys 573, 612-613
Platichthys stellatus 5, 567, 594, 612, 613
 Platytroctidae 19, 51, 77
Plectobranchnus evides 4, 505, 507, 510, 511
Pleurogrammus monopterygius 20, 341, 348, 349
Pleuronectes 567, 573, 614-615
Pleuronectes quadrituberculatus 5, 567, 614, 615
 Pleuronectidae 18-19, 565, 567, 573-623
 Pleuronectiformes 18, 565-625
Pleuronichthys 18, 20, 573, 616-619
Pleuronichthys coenosus 567, 616, 617, 618
Pleuronichthys decurrens 22, 567, 616, 618, 619
Polyacanthonotus challengerii 33-34
Polypera beringiana 475
Polypera greeni 475
Poroclinus rothrocki 503, 507
Porocottus 361
Poromitra 252, 253
Poromitra crassiceps 253
Protomyctophum 147-148, 150-153
Protomyctophum crockeri 146, 150, 151, 152
Protomyctophum thompsoni 146, 150, 152, 153
Psettichthys 573, 599, 600-601, 620-621
Psettichthys melanostictus 5, 567, 594, 599, 600-601, 620, 621
Pseudopentaceros spp. 494, 495
Psychrolutes 361, 364, 436-437
Psychrolutes paradoxus 4, 365, 434, 436, 437
Psychrolutes phrictus 436, 438
 Ptilichthyidae 19, 483, 528-529
Ptilichthys goodei 528, 529
Puzanovia rubra 498
Radulinus 20, 361-362, 406-409
Radulinus asprellus 23, 406, 407, 408
Radulinus boleoides 408, 409
Rastrinus scutigera 399
Reinhardtius 573, 622-623
Reinhardtius hippoglossoides 567, 580, 622, 623
Rhaphiocottus 361-362, 366-367
Rhaphiocottus richardsoni 20, 21, 363, 366, 367
Rhinoliparis attenuatus 475
Rhinoliparis barbulifer 470, 475, 480, 481
Rimicola muscarum 232, 233
Rondeletia loricata 256
 Rondeletiidae 19, 249, 256
Ronquilus jordani 4, 27, 496, 497
Sagamichthys abei 77
 Salmonidae 51
 Salmoniformes 18, 51-81
 Salmonoidei 18, 79-81
Sardinops sagax 44, 46, 47
Sarritor frenatus 456
Sarritor leptorhynchus 456
Sciaonius pedicellaris 217
 Sciaenidae 19, 483, 492-493
 Scomberesocidae 19, 235-237
Scomber japonicus 486, 492, 554, 555
 Scombridae 19, 483, 554-555
 Scombroidei 18, 551-555

- Scopelarchidae* 19, 125-127
Scopeloberyx 252, 253
Scopeloberyx robustus 253
Scopelogadus 252, 253
Scopelogadus bispinosus 253
Scopelomorpha 125-179
Scopelosarus harryi 128, 129
Scorpaenichthys 362-362, 382-383
Scorpaenichthys marmoratus 20, 22, 382, 383, 406
Scorpaenidae 19, 267-337
Scorpaeniformes 18, 267-481
Scorpaenoidei 18, 267-337
Sebastes 267-272, 273-280, 281, 282-287, 288-335
Sebastes aleutianus 269, 271, 273, 283, 288, 289
Sebastes alutus 269, 271, 273, 283
Sebastes auriculatus 269, 271, 273, 285, 290, 291, 296
Sebastes aurora 269, 271, 274, 287, 292, 293
Sebastes babcocki 269, 271, 274, 287, 294, 295
Sebastes borealis 269, 271
Sebastes brevispinis 3, 269, 271, 274, 285
Sebastes caurinus 3, 269, 271, 275, 284, 290, 296, 297
Sebastes chlorostictus 269, 271, 275, 287
Sebastes ciliatus 269, 271, 275
Sebastes crameri 269, 271, 275, 284, 298, 299
Sebastes diploproa 269, 271, 276, 283, 300, 301
Sebastes elongatus 269, 271, 276, 287, 302, 303
Sebastes emphaeus 269, 271, 286, 304, 305
Sebastes entomelas 269, 271, 276, 282, 306, 307
Sebastes flavidus 269, 271, 276, 282, 308, 309, 314
Sebastes glaucus 269, 271
Sebastes goodei 269, 271, 277, 284
Sebastes helvomaculatus 269, 271, 277, 287, 310, 311
Sebastes jordani 3, 269, 271, 277, 286, 312, 313
Sebastes maliger 269, 271, 277
Sebastes melanops 3, 269, 271, 277, 282, 308, 314, 315
Sebastes melanostictus 269, 271
Sebastes melanostomus 269-271, 277, 284, 316, 317
Sebastes miniatus 269, 271, 278, 282, 318, 319
Sebastes mystinus 269, 271, 278, 282, 320, 321
Sebastes nebulosus 269, 271
Sebastes nigrocinctus 269, 271, 284
Sebastes paucispinis 269, 271, 278, 283, 322, 323
Sebastes pinniger 269, 271, 278, 282, 324, 325
Sebastes polyspinis 3, 269, 271, 278, 286
Sebastes proriger 269, 271, 278, 285, 326, 327
Sebastes rastrelliger 269, 271, 285
Sebastes reedi 269, 271, 279, 284, 328, 329
Sebastes ruberrimus 269, 271, 279, 287
Sebastes rufus 3, 269, 271, 279, 286, 330, 331
Sebastes saxicola 269, 271, 279, 284, 332, 333
Sebastes variegatus 3, 269, 271, 280
Sebastes wilsoni 269, 271, 286
Sebastes zacentrus 3, 269, 271-272, 280, 285, 334, 335
Sebastinae 268-335
Sebastobinae 268, 336-337
Sebastolobus 268, 336-337
Sebastolobus alascanus 336, 337
Sebastolobus altivelis 336, 337
Sebastolobus macrochir 336
Sebastolobus spp. 254, 336, 337
Serrivomer jespersenii 33, 36, 37
Serrivomeridae 19, 31-33, 36, 37
Sigmistes 361
Spectrunculus grandis 211-212, 213
Spectrunculus radcliffei 212
Spirinchus starksi 79
Spirinchus thaleichthys 79
Stelgistrum 361
Stellerina xyosterna 456, 457
Stenobranchius 146-147, 172-173
Stenobranchius leucopsarus 3, 146, 172, 173, 176
Stenobranchius nannochir 146, 172
Sternias 361
Sternoptychidae 19, 83, 92-105
Sternoptyx 84, 86, 92, 102, 103
Sternoptyx diaphana 93, 102, 103
Sternoptyx pseudobscura 93, 102, 103
Sternoptyx spp. 102, 104
Stichaeidae 19, 26, 27, 483, 500, 501, 502, 503, 504, 505, 507-519
Stichaeinae 500-501, 507
Stichaeini 496, 500, 501, 507
Stichaeus punctatus 496, 501, 507
Stilegicottus 361
Stomiiformes 18, 83-84, 85, 86-123
Stomioidei 107-123
Stromateidae 19, 483, 562-563
Stromateoidei 18, 558-563
Symbolophorus 147, 154-155
Symbolophorus californiensis 146, 154, 155
Symphurus atricauda 567, 624, 625
Synaphobranchidae 19, 31-34, 35
Synchirus 361-362, 410-411
Synchirus gilli 4, 410, 411
Syngnathidae 19, 261, 264, 265
Syngnathus leptorhynchus 264
Synodontidae 19, 125, 132-133
Synodus lucioceps 132, 133
Taaningichthys 146-147, 174, 175
Taaningichthys bathyphilus 146, 174-175
Tactostoma 106, 116-117
Tactostoma macropus 3, 106, 116, 117
Talismania bifurcata 76
Taractes asper 488
Taranetzella lycoderma 498
Tarletonbeania 147, 158-159
Tarletonbeania crenularis 146, 156, 158, 159
Tarletonbeania taylori 146, 158
Taurocottus 361
Temnocora candida 475
Tetragonuridae 19, 483, 560-561
Tetragonurus cuvieri 558, 560, 561
Tetraodontiformes 18, 627-629
Thalassenchelys coheni 3, 32-34, 35

Thaleichthys pacificus 79
Thecopterus 361
Theragra chalcogramma 1, 3, 189-190, 194, 196, 198, 199
Thyriscus 361
Trachinoidei 18, 532-533
Trachipteridae 19, 243, 246-247
Trachipterus altivelis 3, 244, 246, 247
Trachurus symmetricus 486, 487
Trichiuridae 19, 483, 551-552, 553
Trichodon trichodon 532, 533
Trichodontidae 19, 483, 532-533
Triglops 361-362, 412, 413
Triglops A 412, 413
Triglops B 412, 413
Triglops forficata 412
Triglops jordani 412
Triglops macellus 412
Triglops metopias 412
Triglops pingeli 412
Triglops scepticus 412, 413
Venefica sp. 33, 36
Xeneretmus latifrons 456, 460, 462, 463
Xeneretmus leiops 456
Xeneretmus triacanthus 456
Xenocoelidae 19, 31-34, 35
Xenomystax atrarius 33, 36
Xeropes fucorum 522
Xiphister atropurpureus 4, 505, 507, 516, 517, 518
Xiphister mucosus 507, 516, 518, 519
Xiphisterinae 504-505, 507
Xiphisterini 504, 505, 507
Zaniolepis frenata 341, 344-345
Zaniolepis latipinnis 341, 344-345
Zaniolepis spp. 341-342, 344, 345
Zaprora silenus 20, 24, 530, 531
Zaproridae 19, 483, 530-531
Zeiformes 18, 257-259
Zesticelus 361
Zoarcidae 19, 483, 498, 499
Zoarcoidei 18, 496-531

Italic page numbers indicate illustrations.

Index to common names

- Anchovies 43, 48-49
Anchovy, northern 48, 49
Argentine
 bluethroat 54, 55
 Pacific 52, 53
Argentines 52-55
Armorheads 494, 495
Arrowtail 184, 185
Arrowtails 184-185
Barracudina
 duckbill 138, 139
 ribbon 136, 137
 slender 134, 135
Barracudinas 134-139
Barreleye 74, 75
Bigscale, highsnout 254, 255
Bigscapes 252, 253, 254-255
Blackdragon, Pacific 120, 121
Blackdragons 120-123
Blacksmelt
 Pacific 62, 63
 popeye 60, 61
 robust 58, 59
 snubnose 64, 65
Blenny, ocellated 508, 509
Bocaccio 322, 323
Bottlelights 104, 105
Bristlemouths 87-88, 89, 90, 91
Brotula, red 218, 219
Brotulas 218-219
Butterfishes 562-563
Cabezon 382, 383
Capelin 80, 81
Clingfish
 kelp 232, 233
 northern 230, 231
Clingfishes 229-233
Cockscomb, high 512, 513
Cod
 Arctic 190, 191
 Pacific 194, 195
 saffron 192, 193
Codfishes 189-199
Codlings 182, 183
Combfishes 344, 345
Croaker, white 492, 493
Croakers 492-493
Cusk-eel, spotted 214, 215
Cusk-eels 212, 213, 214-215
Cutlassfishes 551-552, 553
Daggertooth 140, 141
Daggertoos 140-141
Deepsea smelts 51, 56-69
Dragonfish
 highfin 110, 111
 longfin 116, 117
 pitgum 114, 115
Dragonfishes, scaleless 110-112, 113, 114-117
Dreamer, smoothhead 224, 225
Dreamers 222-227
Eelpouts 498, 499
Eels 31-34, 35, 36, 37, 38-41
Fangtooth 250, 251
Fangtoos 250-251
Flashlightfish
 California 150, 151
 chubby 148, 149
 northern 152, 153
Flounder
 Arctic 604, 605
 arrowtooth 578, 579
 Bering 590, 591
 Kamchatka 576, 577
 long 584, 585
 spiny 574, 575
 starry 612, 613
Flounders
 lefteye 568-571
 righteye 573-623
Glowingfish 144-145
Gobies 543-549
Goby
 arrow 544, 545
 bay 548, 549
 blackeye 546, 547
Greenling
 kelp 352, 353
 masked 356, 357
 painted 342, 343
 rock 354, 355
 whitespotted 358, 359
Greenlings 341-350, 351, 352-359
Grenadier
 ghostly 206, 207
 Pacific 202, 203
 threadfin 204, 205
Grenadiers 200-209
Gunnels 522, 523
Hake, Pacific 186, 187
Hakes 186-187
Halibut
 Greenland 622, 623
 Pacific 592, 593
Hatchetfish
 lowcrest 100, 101
 slender 94, 95
 spurred 96, 97
 tropical 98, 99

Hatchetfishes 92-105
 Headlightfish, California 176, 177
 Herring, Pacific 44, 45
 Herrings 43-47
 Irish lord
 banded 372, 373
 brown 378, 379
 longfin 380, 381
 red 374, 375
 yellow 376, 377
 Irish lords 368, 369, 370, 371, 372-381
 Jacks 484-487
 Jacksmelt 240, 241
 Kelpfish, giant 536, 537
 Kelpfishes 535-537
 King-of-the-salmon 246, 247
 Lampfish
 broadfin 168, 169
 dogtooth 160, 161
 giant 170, 171
 patchwork 178, 179
 pinpoint 166, 167
 sunbeam 162, 163
 Lancetfish, longnose 142, 143
 Lancetfishes 142-143
 Lanternfish
 blue 158, 159
 California 154, 155
 northern 172, 173
 Lanternfishes 146-179
 Lingcod 346, 347
 Lizardfish
 California 132, 133
 highfin 130, 131
 Lizardfishes 132-133
 Lizardfishes, deepsea 130-131
 Loosejaw, shiny 118, 119
 Loosejaws 118-119
 Louvar 556, 557
 Louvars 556-557
 Lumpsucker
 Pacific spiny 468, 469
 smooth 466, 467
 Lumpsuckers 465-469
 Mackerel
 Atka 348, 349
 chub 554, 555
 jack 486, 487
 Mackerels 554-555
 Medusafish 558, 559
 Medusafishes 558-559
 Molas 628-629
 Opah 244, 245
 Opahs 244-245
 Oreos 258, 259
 Paperbone, scaly 128, 129
 Paperbones 128-129
 Pearleyes 126, 127
 Pilotfish 484, 485
 Pipefishes 264, 265
 Plaice, Alaska 614, 615
 Poacher
 blacktip 462, 463
 kelp 458, 459
 Poachers 456, 457, 458-463
 Pollock, walleye 198, 199
 Pomfret, Pacific 488, 489
 Pomfrets 488-489
 Pompano, Pacific 562, 563
 Prickleback
 black 516, 517
 bluebarred 510, 511
 ribbon 514, 515
 rock 518, 519
 Pricklebacks 500, 501, 502, 503, 504, 505, 507-519
 Prowfish 530, 531
 Prowfishes 530-531
 Quillfish 528, 529
 Quillfishes 528-529
 Ragfish 538, 539
 Ragfishes 538-539
 Ribbonfishes 246-247
 Rockfish
 aurora 292, 293
 bank 330, 331
 black 314, 315
 blackgill 316, 317
 blue 320, 321
 brown 290, 291
 canary 324, 325
 copper 296, 297
 darkblotched 298, 299
 greenstriped 302, 303
 Puget Sound 304, 305
 redbanded 294, 295
 redstripe 326, 327
 rosethorn 310, 311
 rougeye 288, 289
 sharpchin 334, 335
 shortbelly 312, 313
 splitnose 300, 301
 stripetail 332, 333
 vermilion 318, 319
 widow 306, 307
 yellowmouth 328, 329
 yellowtail 308, 309
 Rockfishes 268-272, 273-280, 281, 282-287, 288-335
 Rockhead 460, 461
 Ronquils 496, 497
 Sablefish 338, 339
 Sablefishes 338-339
 Sanddab
 Pacific 568, 569
 speckled 570, 571

Sanddabs 568-571
 Sandfish, Pacific 532, 533
 Sandfishes 532-533
 Sandlance, Pacific 540, 541
 Sandlances 540-541
 Sardine, Pacific 46, 47
 Sauries 236-237
 Saury, Pacific 236, 237
 Sculpin
 Arctic staghorn 394, 395
 bald 428, 429
 buffalo 390, 391
 calico 424, 425
 crested 448, 449
 darkfin 442, 443
 darter 408, 409
 fluffy 432, 433
 grunt 366, 367
 manacled 410, 411
 mosshead 426, 427
 Pacific staghorn 446, 447
 padded 414, 415
 prickly 444, 445
 Puget Sound 384, 385
 rosylip 386, 387
 roughback 388, 389
 sailfin 454, 455
 scalyhead 416, 417
 sharpnose 422, 423
 shorthorn 402, 403
 silverspotted 450, 451
 slim 406, 407
 smoothhead 418, 419
 soft 434, 435
 spinyhead 440, 441
 tadpole 436, 437
 thornback 404, 405
 tidepool 430, 431
 Sculpins 360-362, 363, 364, 365, 366-455
 Seadevils 222-223
 Sea raven, shaggy 452, 453
 Silversides 238-241
 Skilfish 338
 Slickheads 76
 Smelts 51, 79-81
 Smoothtongue
 California 68, 69
 northern 66, 67
 Snailfish
 longnose 480, 481
 prickly 478, 479
 tadpole 476, 477
 Snailfishes 465, 460, 471-473, 474-481
 Snipe eel
 blackline 38, 39
 slender 40, 41
 Spiny eels 31-35

Sole
 butter 594, 595
 C-O 616, 617
 curlfin 618, 619
 deepsea 580, 581
 Dover 608, 609
 English 610, 611
 flathead 588, 589
 petrale 582, 583
 rex 586, 587
 rock 596, 597
 sand 620, 621
 slender 606, 607
 yellowfin 602, 603
 Spookfish
 javelin 70, 71
 winged 72-73
 Spookfishes 70-72, 73, 74-75
 Squaretail, smalleye 560, 561
 Squaretails 560-561
 Sunfish, ocean 628, 629
 Thornyheads 336, 337
 Tomcod, Pacific 196, 197
 Tonguefish, California 624, 625
 Tonguefishes 624-625
 Topsmelt 238, 239
 Tubeshoulders 77
 Tube-snout 262, 263
 Tubesnouts 262-263
 Veilfin, bigmouth 490
 Veilfins 490, 491
 Viperfish, Pacific 108, 109
 Viperfishes 108-109
 Whalefishes 256
 Wolf-eel 526, 527
 Wolffish, Bering 524, 525
 Wolffishes 524-527
 Wrymouths 521, 522

Italic page numbers indicate illustrations.