

IMMIGRATION OF FISHES THROUGH THE SUEZ CANAL¹

ADAM BEN-TUVIA²

ABSTRACT

The number of Red Sea fishes found in the eastern Mediterranean amounts to 36 species. Twelve immigrants, namely: *Spratelloides delicatulus*, *Herklotsichthys punctatus*, *Tylosurus chorum*, *Sebastapistes nuchalis*, *Epinephelus tauvina*, *Autisthes puta*, *Pelates quadrilineatus*, *Silago sihama*, *Rhonsicus stridens*, *Crenidens crenidens*, *Rastrelliger kanagurta*, *Scomberomorus commerson*, were found in the last 12 yr. The southward migration, from the Mediterranean to the Red Sea is almost negligible. Only *Liza aurata*, *Dicentrarchus punctatus*, and perhaps *Carcharhinus plumbeus* can be regarded as Mediterranean immigrants.

In studying the immigration of fishes through the Suez Canal, three zoocological areas must be taken into consideration: 1) the northern Red Sea; 2) the eastern Mediterranean; and 3) the Suez Canal itself in which many marine animals from the two neighboring areas have found a permanent habitat (Steinitz 1968).

The prevailing hydrographic conditions differ in these three areas, although the salinities and summer temperatures are to some extent similar (Morcos 1967, 1970; El-Saby 1968; Oren 1970; Oren and Hornung 1972). Temperature and salinity are the main abiotic factors influencing the distribution of organisms over large zoogeographical areas. Often they also have a decisive influence on the ecological distribution of species in various biotopes of an area.

The process of immigration is highly selective. Common species of the home seas are not necessarily successful immigrants in a new region. Similar effects have been shown to occur in many forms of colonization (MacArthur and Wilson 1967). The adaptation of a species to a new area requires adjustment of its reproductive processes, especially with regard to the correct timing of spawning in order to ensure suitable physical and ecological conditions for the development and survival of the young stages.

It is evident that the direction of immigration is mainly from the Red Sea into the Mediterranean (Figure 1). The possible causes of such one way immigration have been discussed elsewhere (Aron

and Smith 1971; Ben-Tuvia 1971a, 1973; Por 1971a, b).

Thirty-six Red Sea or cosmopolitan species can be regarded as Suez Canal immigrants. Twelve of them were found within the last 12 yr. Evidently, immigration is a continuous process, and over time the probability of suitable species of fishes entering the Suez Canal and colonizing the new region increases. Time also plays an essential role in the biological processes of adaptation of the species to the modified conditions of life. More resistant species, endowed with greater plasticity of genetic characters, can form local "races" within a few generations by natural selection in the new environment (Kosswig 1974). But first they need a firm foothold on the other side of the Canal, geographically close to the parental stock and in places where conditions are not drastically different from their normal habitat.

Recently I had an opportunity to collect samples from the Gulf of Suez (Ben-Tuvia and Grofit 1973), Suez Canal (Steinitz and Ben-Tuvia 1972), and Bardawil Lagoon (Ben-Tuvia 1975a) which revealed interesting data on the distribution of immigrants. Many of the species which have successfully colonized the eastern Mediterranean, such as *Saurida undosquamis*, *Leiognathus klunzingeri*, *Upeneus moluccensis*, and *U. asymmetricus*, and which are abundant there, are also dominant species on the trawling grounds of the Gulf of Suez.

High percentage of Red Sea fishes found in the hypersaline Bardawil Lagoon on the northern coast of Sinai indicates that it may serve as a stepping stone in the immigration of Red Sea fishes into the Mediterranean, especially if we regard it as a part of the system of lakes and lagoons of the Isthmus of Suez (Por 1971a). Among 55

¹This paper was read at the 17th International Zoological Congress in Monte Carlo, 25-30 September 1972; some changes were introduced to include more recent information on immigrants.

²Department of Zoology, Hebrew University of Jerusalem, Jerusalem, Israel.




FIGURE 1.—Distribution of Red Sea fishes in the Mediterranean Sea: coast of Israel and Sinai with 34 species; coast of Lebanon with 27 species; Aegean Sea with 9 species. Numbers in circles refer to the following fishes: 1. *Signaus luridus*, 2. *S. rivulatus*, 3. *Stephanolepis diaspros*, 4. *Leiognathus klunzingeri*, 5. *Pranesus pinguis*, and 6. *Parexocoetus mento*.

species collected in Bardawil Lagoon, 14 species (25.5%) are Red Sea immigrants in comparison with about 11% estimated for all the fishes collected in the eastern Mediterranean (Ben-Tuvia 1971b).

The following Red Sea immigrants were collected in Bardawil Lagoon: *Hemiramphus far*, *Aphanius dispar*, *Atule djeddaba*, *Pelates quadrilineatus*, *Leiognathus klunzingeri*, *Upeneus moluccensis*, *Crenidens crenidens*, *Liza carinata*, *Pranesus pinguis*, *Siganus luridus*, *S. rivulatus*, *Sphaeroides spadiceus*, and recently *Herklotsichthys punctatus* and *Autisthes puta*.

In addition to the Red Sea immigrants, four cosmopolitan species were also found in Bardawil Lagoon: namely, *Sardinella aurita*, *Lobotes surinamensis*, *Mugil cephalus*, and *Echeneis naucrates*.

RED SEA FISHES IN THE MEDITERRANEAN SEA

In my previous summary of the immigration of Red Sea fishes into the Mediterranean (Ben-Tuvia 1966), 24 species were listed; 12 additional immigrants were since found, most of them are rare fishes at this time. Their names, distribution, and the maximum size observed in the Mediterranean Sea are given in Table 1. As yet, only one specimen of *Tylosurus choram* (Collette and Parin 1970), *Rastrelliger kanagurta* (Collette 1970), *Sebastes nuchalis* (Froiland 1972), and *Silago sihama* (Mouneimne 1977) have been reported from the eastern Mediterranean.

Froiland (1972), who studied scorpaenids from Cyprus in the collection of the Hebrew University Zoological Museum, reported one specimen of *Sebastes nuchalis* (Günther) 58 mm long. This species is known from the Indo-Pacific, including East Africa, but no records are available from the Red Sea, Suez Canal, and other localities in the eastern Mediterranean besides Cyprus. Froiland assumed that this scorpaenid "migrated through the Suez Canal."

Epinephelus tauvina (Ben-Tuvia and Lourie 1969), *Pelates quadrilineatus* (Lourie and Ben-Tuvia 1971), and *Scomberomorus commerson* (George and Athanassiou 1965) were collected on several occasions and do not seem to be very rare. *Herklotsichthys punctatus* and *Rhonciscus stridens* have been found recently in the eastern Mediterranean at several localities (Ben-Tuvia 1967, 1977; Mouneimne 1977). One specimen of

Spratelloides delicatulus (Bennet) 51 mm long was collected with rotenone on 4 June 1973 in a shallow bay about 3 km south of Atlit (Ben-Tuvia, unpubl. data). The occurrence of *Crenidens crenidens* and *Autisthes puta* is restricted to the hypersaline Bardawil Lagoon.

It is of interest to note that some of the Red Sea immigrants have been found in recent years in new localities west of Levant (Figure 1); *Stephanolepis diaspros* in the Gulf of Taranto, south Italy and Gulf of Gabes, Tunisia (Tortonese 1967); *Siganus rivulatus* off Tobrouk, Libya (Tortonese 1970).

With the exception of perhaps two fishes, *Saurida undosquamis* and *Siganus luridus*, very little information is available on the rate of increase of the immigrant population and the ecological influence of their appearance in the new region. The first Mediterranean specimen of *Saurida undosquamis*, 145 mm standard length, was collected in Haifa Bay, by a trawler, in December 1952. Additional specimens (160-171 mm) were collected in Haifa in February 1953. According to my observations in August 1953, taken on the deck of a commercial trawler, this fish was fairly common in the Gaza-El'Arish area, and 10-20 specimens were usually caught in each haul. There is also some information on the trawling activities in the Gulf of Iskenderun and Mersin on the Anatolian coast of Turkey. In August 1952, I participated in a commercial trawling cruise to the Gulf of Mersin between Karadash-Burnuun and Bagase, during which no *Saurida undosquamis* were collected. But in 1956, this fish was common on the same trawling grounds and fished in commercial quantities. According to catch data, *S. undosquamis* started to appear in commercial quantities in the year 1955, first on the southern fishing grounds (El'Arish to Tel Aviv) and towards the end of the same year and especially during 1956 also on the northern fishing grounds such as Haifa Bay, and even in the Gulf of Iskenderun and Mersin (Ben-Yami 1955; Oren 1957).

It is worthwhile noticing that no specimens of *S. undosquamis* were found before December 1952, although Mediterranean fishes were collected in Israel during earlier years by the staff of the Sea Fisheries Research Station, Haifa, and by scientists of the Hebrew University, Jerusalem.

No less interesting is the sudden appearance of *Siganus luridus* (Ben-Tuvia 1964). Not a single specimen was found before February 1955, in spite of extensive collecting activities in Israel during

TABLE 1.—Data on Red Sea fishes found in the Mediterranean.

| Species | Occurrence in eastern Mediterranean | Ecological distribution | SL ¹ (mm) | Geographical distribution | | | |
|--------------------------------------|-------------------------------------|-------------------------|----------------------|---------------------------|---------|------------|----------------------------------|
| | | | | Indo-Pacific | Red Sea | Suez Canal | Mediterranean ² |
| <i>Carcharhinus brevipinna</i> | Common | Inshore-pelagic | 1,200 | + | + | - | Israel |
| <i>Himantura uarnak</i> | Common | Demersal | ³ 1,200 | + | + | + | Anatolia |
| <i>Dussumieria acuta</i> | Very common | Inshore-pelagic | 155 | + | + | + | Anatolia |
| <i>Herklotsichthys punctatus</i> | Rare | Inshore-pelagic | 76 | + | + | + | Lebanon |
| <i>Etrumeus teres</i> | Single record | Inshore-pelagic | 170 | + | + | - | Israel |
| <i>Spratelloides delicatulus</i> | Single record | Inshore-pelagic | 51 | + | + | + | Israel |
| <i>Saurida undosquamis</i> | Very common | Demersal | 335 | + | + | + | Anatolia |
| <i>Paraexocoetus mento</i> | Common | Pelagic | 111 | + | + | - | Aegean Sea; Gulf of Sidra |
| <i>Hemiramphus far</i> | Common | Inshore-pelagic | 340 | + | + | + | Aegean Sea |
| <i>Tylosurus chorum</i> | Single record | Inshore-pelagic | — | + | + | + | Lebanon |
| <i>Aphanius dispar</i> | Common | Sublittoral | 46 | + ⁴ | + | + | Israel |
| <i>Pranesus pinguis</i> | Very common | Inshore-pelagic | 120 | + | + | + | Anatolia; Libya |
| <i>Holocentrus ruber</i> | Common | Inshore-pelagic | 177 | + | + | - | Aegean Sea |
| <i>Sebastapistes nuchalis</i> | Single record | Sublittoral | 58 | + | - | - | Cyprus |
| <i>Platycephalus indicus</i> | Rare | Demersal | 550 | + | + | + | Lebanon |
| <i>Epinephelus tauvina</i> | Rare | Demersal | 790 | + | + | + | Israel |
| <i>Autistes puta</i> | Rare | Sublittoral | 133 | + | + | - | Bardawil Lagoon |
| <i>Pelates quadrilineatus</i> | Rare | Demersal | 116 | + | + | - | Lebanon |
| <i>Apogonichthyoides nigripinnis</i> | Common | Sublittoral | 77 | + | + | + | Lebanon |
| <i>Silago sihama</i> | Single record | Sublittoral | 115 | + | + | - | Lebanon |
| <i>Atule djeddaba</i> | Very common | Inshore-pelagic | 230 | + | + | + | Lebanon |
| <i>Leiognathus klunzingeri</i> | Very common | Demersal | 77 | + | + | + | Aegean Sea; Lampedusa |
| <i>Rhonciscus stridens</i> | Rare | Demersal | 128 | + | + | + | Lebanon |
| <i>Upeneus asymmetricus</i> | Common | Demersal | 140 | + | + | + | Anatolia |
| <i>Upeneus moluccensis</i> | Very common | Demersal | 170 | + | + | + | Aegean Sea |
| <i>Crenidens crenidens</i> | Rare | Demersal | 150 | + | + | + | Bardawil Lagoon |
| <i>Liza carinata</i> | Rare | Inshore-pelagic | 128 | + | + | + | Egypt; Israel |
| <i>Sphyræna chrysaena</i> | Very common | Inshore-pelagic | 225 | + | + | + | Anatolia |
| <i>Callionymus filamentosus</i> | Common | Demersal | 110 | + | + | + | Lebanon |
| <i>Siganus luridus</i> | Common | Sublittoral | 215 | - | + | - | Aegean Sea; Tunisia |
| <i>Siganus rivulatus</i> | Very common | Sublittoral | 223 | + | + | + | Aegean Sea; Libiya |
| <i>Rastrelliger kanagurta</i> | Single record | Pelagic | 215 | + | + | - | Israel |
| <i>Scomberomorus commerson</i> | Rare | Pelagic | 460 | + | + | - | Lebanon |
| <i>Dollfusichthys sinusarabici</i> | Common | Demersal | 110 | - | + | + | Lebanon |
| <i>Stephanolepis diaspros</i> | Common | Demersal | 124 | + ⁵ | + | + | Aegean Sea; south Italy; Tunisia |
| <i>Sphaeroides spadiceus</i> | Rare | Demersal | 245 | + | + | + | Aegean Sea |

¹Maximum length observed in Mediterranean.²Farthest point of distribution.³Length of disc.⁴Indian Ocean only.⁵Persian Gulf only.

the preceding years. Then the fish suddenly appeared to be fairly common along the whole Mediterranean coast of Israel, although it remained inferior in numbers to a previous Red Sea immigrant, *S. rivulatus*. Recent observations and reports show that *S. luridus* spread rapidly in the Mediterranean towards the west and north. It is common in Lebanon (George and Athanassiou 1967), Cyprus, Rhodes, and has even reached Tunisia (Ktari-Chakroun and Bouhlal 1971), a distance of more than 1,000 n.mi. from the Suez Canal.

It is a common feature of invading organisms that after an initial period of successful adaptation to the new and basically favorable environment, they may suddenly increase in number and spread to adjacent areas (Elton 1958). Various factors such as decrease in salinities of the Bitter Lakes and cessation of the Nile flood after the completion of the Aswan Dam may facilitate the passage or dispersion of Red Sea species. There is speculation that this immigration was also favored by a series of warm years (Oren 1956; Ben-Yami and Glaser 1974).

MEDITERRANEAN FISHES IN THE RED SEA

The occurrence in the Red Sea of the two Atlanto-Mediterranean and one cosmopolitan species (Table 2) is assumed to be the result of Suez Canal migration. The discovery of *Dicentrarchus punctatus* in the lagoon of El Bilaiyim, situated about 180 km south of the entrance to the Suez Canal (Ben-Tuvia 1971a), is one of the few indisputable evidences of the immigration of a Mediterranean fish into the Gulf of Suez. However, we have to bear in mind that the conditions of El Bilaiyim differ considerably from those of the Gulf proper. Salinities are much higher (50-60‰ according to measurements taken in June 1968) and most probably the seasonal and diurnal fluctuations are greater than those of surrounding waters. In this particular biotope, less competition is expected than in the open coastal waters. *Dicentrarchus punctatus* is known to inhabit the Bardawil Lagoon on the northern (Mediterranean) coast of Sinai, where salinities may reach 80‰. It was noted already by Tillier (1902) that this fish was common in the Suez Canal and reached its southern entrance. Evidently, it settled in the Canal soon after its opening.

A taxonomic study of Red Sea mugilids (Ben-Tuvia 1975b) revealed that another Mediterranean immigrant, *Liza aurata* is common in the northern Red Sea. An earlier record of its presence was made by Al-Hussaini (1947) who examined the intestine of this mullet that was captured off Ghardaqa. *Liza aurata* is known to be euryhaline and could cross the Suez Canal or an earlier freshwater connection that was established by the ancient Egyptian pharaohs and Persian kings between the Mediterranean and the Red Sea using an arm of the River Nile. This fish was reported in the Suez Canal by Tillier (1902). I found it to be common in Great Bitter Lake (Ben-Tuvia 1975a).

Recently two specimens of a sandbar shark, *Carcharhinus plumbeus*, were found in the Red Sea. One specimen, a male, 1,600 mm total length

was collected on 6 August 1971 in Dahab, Gulf of Aqaba; the second specimen, a gravid female, 1,764 mm was collected on October 1975 in Ras Muhammad at the entrance to the Gulf of Suez (Baranes and Ben-Tuvia in press). Five additional specimens varying in length between 1,500 and 1,800 mm have been found very recently in the same region (unpubl. data). *Carcharhinus plumbeus* (known also under the name of *C. milberti*) is common on both sides of the Atlantic and is well known in the Mediterranean Sea (Tortonese 1956; Ben-Tuvia 1971b; Compagno 1973). The recent appearance of the sandbar shark in the northern Red Sea could be due to immigration through the Suez Canal although the possibility of penetration from the western Indian Ocean should not be excluded.

Special consideration should be given to *Serranus cabrilla*, which is common in all parts of the Mediterranean, in the Suez Canal, and also in the Red Sea. However, it cannot be regarded as an example of Suez Canal immigration, since a 17-cm long specimen was collected by Hemprich and Ehrenberg in the Red Sea before the completion of the Canal (Klunzinger 1884). According to my observations in September 1970, *S. cabrilla* is common in the northern part of the Gulf of Suez. Individuals were easily observed on sandy patches between coral heads and rocks in the shallow coastal waters off Ras Masalla and Ras Sudar. A total of 10 specimens, 52-100 mm, were collected from the Gulf of Suez plus 1 specimen, 70 mm standard length, from the Gulf of Aqaba. The abundance of *S. cabrilla* in the northern section of the Gulf of Suez indicates the possibility that the present distribution might be related to the proximity of the Suez Canal. However, further taxonomic and behavioral studies will be needed to ascertain the relationships between the Red Sea and the Mediterranean populations.

Serranus cabrilla in the Mediterranean shows great plasticity and adaptability to various ecological conditions. This fish is found in shallow coastal waters on sandy beaches and among rocks. It is

TABLE 2.—Data on Mediterranean fishes found in the Red Sea.

| Scientific name | Occurrence in Red Sea | Ecological distribution | SL ¹ (mm) | Geographical distribution | | | |
|--------------------------------|-----------------------|-------------------------|----------------------|---------------------------|---------------|------------|-------------------------------|
| | | | | West Atlantic | East Atlantic | Suez Canal | Red Sea record |
| <i>Carcharhinus plumbeus</i> | Rare | Pelagic | 1,764 | + | + | - | Gulf of Aqaba Gulf of Suez |
| <i>Dicentrarchus punctatus</i> | Rare | Demersal | 245 | - | + | + | Gulf of Suez |
| <i>Liza aurata</i> | Very common | Inshore pelagic | 320 | - | + | + | Gulf of Aqaba Gulf of Suez |

¹Maximum length observed in Red Sea.

also occasionally found on trawling grounds in various depths, at least up to 100 m. It is common throughout the Mediterranean but rare in the Black Sea. In the eastern Atlantic, it occurs from the English Channel to Angola. Smith (1961) quoted its presence in South Africa (Natal), but it seems not to have been recorded from any other part of the Indian Ocean. Gruvel and Chabanaud (1937) reported that this fish is common throughout the Suez Canal.

CONCLUSIONS

The occurrence of large numbers of circum-tropical-cosmopolitan species in the eastern Mediterranean deserves special attention. They demonstrate the distinct faunistic character of the fish population in this area. Of the 290 species of marine fishes identified from the Mediterranean coast of Israel and its immediate neighborhood, 40 species are circumtropical-cosmopolitan. Many of them are found also in the Indo-Pacific and Red Sea regions. Thus, summing up the Red Sea and cosmopolitan fishes in the eastern Mediterranean, there are 76 species which constitute about one-quarter of all fishes identified from this region. The remaining species belong to the Atlanto-Mediterranean fauna.

The Red Sea element in the eastern Mediterranean is particularly pronounced among demersal fishes. This is evident by the large number of demersal immigrants and by their common occurrence. About 17 species are bottom-living fishes, and at least 6 of them are of commercial value. I estimate that they constitute about 21% (by weight) of the Israeli trawl catches and 8% of the inshore fishery (Ben-Tuvia 1973). George and Athanassiou (1967) in their analysis of beach-seine catches of St. George Bay, Lebanon, found that among 26 commercially important fishes, 5 (19%) were Red Sea immigrants.

For a better understanding of Suez Canal immigration, additional taxonomic and biological investigations are required. Comparison of racial characteristics of immigrant fishes could help to clarify the question of the origin and relationship between the Red Sea and the Mediterranean populations. It is suspected that in some cases, exchange of fauna may have taken place before the opening of the Suez Canal as a result of the elevation of sea level and undulation of the Isthmus during the Pleistocene.

Knowledge of the comparative life histories of the immigrant fishes in the two areas is essential for understanding the selective mechanisms controlling passage through the Suez Canal and evaluating extensive ecological changes that the invading species may produce in the new areas of their distribution.

LITERATURE CITED

- AL-HUSSAINI, A. H.
1947. The feeding habits and the morphology of the alimentary tract of some teleosts living in the neighbourhood of the Marine Biological Station, Ghardaqa, Red Sea. Fouad I Univ., Publ. Mar. Biol. Stn. Ghardaqa (Red Sea) 5:1-61.
- ARON, W. I., AND S. H. SMITH.
1971. Ship canals and aquatic ecosystems. *Science* (Wash., D.C.) 174:13-20.
- BARANES, A., AND A. BEN-TUVIA.
In press. The occurrence of the sandbar shark *Carcharhinus plumbeus* (Nardo 1827) in the northern Red Sea. *Isr. J. Zool.*
- BEN-TUVIA, A.
1964. Two siganid fishes of the Red Sea origin in the eastern Mediterranean. *Bull. Sea Fish. Res. Stn. (Haifa)* 37:1-9.
1966. Red Sea fishes recently found in the Mediterranean. *Copeia* 1966:254-275.
1971a. On the occurrence of a Mediterranean serranid fish *Dicentrarchus punctatus* (Bloch) in the Gulf of Suez. *Copeia* 1971:741-743.
1971b. Revised list of the Mediterranean fishes of Israel. *Isr. J. Zool.* 20:1-39.
1973. Man-made changes in the eastern Mediterranean Sea and their effect on the fishery resources. *Mar. Biol. (Berl.)* 19:197-203.
1975a. Comparison of the fish fauna in the Bardawil Lagoon and the Bitter Lakes. *Rapp. Comm. Int. Mer Méditerr.* 23:125-126.
1975b. Mugilid fishes of the Red Sea with a key to the Mediterranean and Red Sea species. *Bamidgeh* 27:14-20.
1976. Fish collections from the eastern Mediterranean, the Red Sea, and inland waters of Israel. *Zool. Mus., Heb. Univ. Jerus.*, 32 p.
1977. New records of Red Sea immigrants in the eastern Mediterranean. *Cybium*, 3e Sér., 1:95-102.
- BEN-TUVIA, A., AND A. GROFIT.
1973. Exploratory trawling in the Gulf of Suez, November 1972. [In Heb., Engl. abstr.] *Fish. Fish Breed. Isr.* 8:8-16.
- BEN-TUVIA, A., AND A. LOURIE.
1969. A Red Sea grouper *Epinephelus tauvina* caught on the Mediterranean coast of Israel. *Isr. J. Zool.* 18:245-247.
- BEN-YAMI, M.
1955. Overfishing or bad season? [In Heb.] *Fishermens' Bull.* 1(6):10-14.
- BEN-YAMI, M., AND T. GLASER.
1974. The invasion of *Saurida undosquamis* (Richardson) into the Levant Basin—An example of biological effect of interoceanic canals. *Fish. Bull., U.S.* 72:359-373.

- COMPAGNO, L. J. V.
1973. Carcharhinidae. In J. C. Hureau and Th. Monod (editors), Check-list of the fishes of the northeastern Atlantic and of the Mediterranean, Clofnam I, p. 23-31. Unesco, Paris.
- COLLETTE, B. B.
1970. *Rastrelliger kanagaruta*, another Red Sea immigrant into the Mediterranean Sea, with a key to the Mediterranean species of Scombridae. Bull. Sea Fish. Res. Stn. (Haifa) 54:3-6.
- COLLETTE, B. B., AND N. V. PARIN.
1970. Needlefishes (Belonidae) of the eastern Atlantic Ocean. Atl. Rep. 11:7-60.
- EL-SABY, K. K.
1968. Effect of the Aswan High Dam on the distribution of salinity in the Suez Canal. Nature (Lond.) 218:758-760.
- ELTON, C. S.
1958. The ecology of invasions by animals and plants. Methuen, Lond., 181 p.
- FRÓILAND, Ó.
1972. Fishes of the family Scorpaenidae from Cyprus, including three new records. Bull. Sea Fish. Res. Stn. (Haifa) 59:5-16.
- GEORGE, C. J., AND V. ATHANASSIOU.
1965. On the occurrence of *Scomberomorus commersoni* (Lacépède) in St. George Bay, Lebanon. Doriana, Ann. Mus. Civico Storia Nat. Genova 4(157):104.
1967. A two-year study of the fishes appearing in the seine fishery of St. George Bay, Lebanon. Ann. Mus. Civico Storia Nat. Genova 76:237-294.
- GRUVEL, A., AND P. CHABANAUD.
1937. Missions A. Gruvel dans le Canal de Suez. II. Poissons. Mem. Inst. Egypt. 35:1-30.
- KLUNZINGER, C. B.
1884. Die Fische des Rothen Meeres. E. Schweizerbart'sche Verlag, Stuttgart, 133 p.
- KOSSWIG, C.
1974. Modificability, a neglected factor for area expansion in marine fish. Istanbul Univ. Fen Fak. Mecm., Ser. B, 39(1-2):1-7.
- KTARI-CHAKROUN, F., AND M. BOUHLAL.
1971. Capture of *Siganus luridus* (Rüppell) dans le Golfe de Tunis. Bull. Inst. Océanogr. Peche Salammbô 2:49-52.
- LOURIE, A., AND A. BEN-TUVIA.
1971. Two Red Sea fishes, *Pelates quadrilineatus* (Bloch) and *Crenidens crenidens* (Forsskal) in the eastern Mediterranean. Isr. J. Zool. 19:203-207.
- MACARTHUR, R. H., AND E. O. WILSON.
1967. The theory of island biogeography. Princeton Univ. Press, Princeton, N.J., 203 p.
- MORCOS, S. A.
1967. Effect of the Aswan High Dam on the current regime in the Suez Canal. Nature (Lond.) 214:901-902.
1970. Physical and chemical oceanography of the Red Sea. Oceanogr. Mar. Biol. Annu. Rev. 8:73-202.
- MOUNEIMNE, N.
1977. Liste des poissons de la côte du Liban (Méditerranée orientale). Cybium, 3e Sér., 1:37-66.
- OREN, O. H.
1957. Changes in temperature of the Eastern Mediterranean Sea in relation to the catch of the Israel trawl fishery during the years 1954-55 and 1955-56. Bull. Inst. Oceanogr. Monaco 1102:1-2.
1970. The Suez Canal and the Aswan High Dam, their effect on the Mediterranean. Underwater Sci. Technol. J. 2:222-229.
- OREN, O. H., AND H. HORNING.
1972. Temperatures and salinities off the Israel Mediterranean coast. Bull. Sea Fish. Res. Stn. (Haifa) 59:17-31.
- POR, F. D.
1971a. One hundred years of the Suez Canal—a century of Lessepsian migration: Retrospects and viewpoints. Syst. Zool. 20:138-159.
1971b. The zoobenthos of the Sirbonian lagoons. Rapp. Comm. Int. Mer Méditerr. 20:247-249.
- SMITH, J. L. B.
1961. The sea fishes of Southern Africa. 4th ed. Central News Agency Ltd., Capetown, 580 p.
- STEINITZ, H.
1968. Remarks on the Suez Canal as pathway and as habitat. Rapp. Comm. Int. Mer Méditerr. 19:139-141.
- STEINITZ, H., AND A. BEN-TUVIA.
1972. Fishes of the Suez Canal. Isr. J. Zool. 21:385-389.
- TILLIER, J. B.
1902. Le Canal de Suez et sa faune ichthyologique. Mém. Soc. Zool. Fr. 15:279-318.
- TORTONESE, E.
1956. Leptocardia, Ciclostomata, Selachii. Fauna Ital. 2, 334 p.
1967. Un pesce plettognato nuovo per i Mari Italiani: *Stephanolepis diaspros* Fr. Br. Ann. Mus. Civico Storia Nat. Genova 4(181):1-4.
1970. On the occurrence of *Siganus* (Pisces) along the coast of North-Africa. Doriana 4(191):1-2.