

Abstract.—We examined 528 bonefish to estimate length and age at sexual maturity and to describe seasonal patterns in gonadal development. These fish ranged from 21 to 702 mm fork length (FL) and were collected in South Florida waters from 1989 to 1995. Gonads of 437 bonefish were examined histologically, and gonadosomatic indices (GSI) were calculated for 449 bonefish. Male bonefish reached 50% sexual maturity (the predicted size and age at which half the individuals are expected to be sexually mature) at 418 mm FL (95% confidence interval 393–443 mm) and an age of 3.6 years (95% confidence interval 3.3–3.9 years). Females reached 50% sexual maturity at 488 mm FL (95% confidence interval 472–504 mm) and 4.2 years (95% confidence interval 3.9–4.6 years). Lengths and ages at 50% maturity for males and females were significantly different. The smallest sexually mature male was 425 mm FL, and the smallest sexually mature female was 358 mm FL. The youngest sexually mature male was 3 years old, and the youngest sexually mature female was 2 years old. Gonadal activity was seasonal and peaked during November–May. Vitellogenic oocytes were present in ovaries in every month except August and September and were most abundant during November–May. Median GSI's were greatest during November–May and least during July–September for both males and females. No fully hydrated ovaries or postovulatory follicles were found, therefore we could not estimate spawning periodicity or batch fecundities. Total fecundity ranged from 0.4 to 1.7 million oocytes and had a significant positive relation to fish weight. The absence of fully hydrated ovaries and postovulatory follicles in the bonefish we sampled suggests that bonefish spawn outside the traditional shallow-water (<2 m) fishing grounds in the Florida Keys.

Maturation and reproductive seasonality in bonefish, *Albula vulpes*, from the waters of the Florida Keys

Roy E. Crabtree*

Derke Snodgrass

Christopher W. Harnden

Florida Marine Research Institute, Department of Environmental Protection
100 Eighth Avenue SE, St. Petersburg, Florida 33701-5095

*E-mail address: crabtree_r@sellors.dep.state.fl.us

Bonefish, *Albula* spp., frequent coastal and inshore waters of tropical seas worldwide and are the basis of economically important recreational fisheries in many areas of their range. Although 23 nominal *Albula* species have been described (Whitehead, 1986), only two Atlantic species, *A. vulpes* and *A. nemoptera*, are recognized (Rivas and Warlen, 1967). In the western Atlantic, *A. vulpes* is common in the Florida Keys, the Bahama Islands, and throughout the Caribbean Sea (Hildebrand, 1963). *Albula nemoptera* appears to have a more restricted distribution than *A. vulpes* and has been reported from the Guianas, Venezuela, Columbia, Panama, Jamaica, and Hispaniola (Rivas and Warlen, 1967; Uyeno et al., 1983). The single record of *A. nemoptera* from Florida waters is considered questionable (Robins and Ray, 1986), and the species has not been reported from the Bahama Islands (Böhlke and Chaplin, 1993).

Bonefish are esteemed for their wariness and fighting abilities, and fishing for them provides an important source of income to Florida Keys and Bahamian fishing guides. Commercial sale of bonefish in Florida is prohibited, and regulations on the recreational fishery restrict catches to one fish per angler per day and the length of captured fish to a minimum total length of

457 mm (390 mm fork length). Most bonefish caught in Florida waters are released.

The life history of bonefish has not been adequately described. Crabtree et al. (1996) described the age and growth of bonefish from South Florida and found that bonefish can attain ages of 19 years. Female and male growth models were significantly different; females were slightly longer than males of the same age. Although the age and growth of Florida Keys bonefish have been studied, important questions remain regarding bonefish reproduction. Bruger (1974) reported sexually mature females as small as 210 mm standard length (221 mm fork length) and as young as 1 year from waters off the Florida Keys, but his sample size was inadequate to determine the age or length at 50% maturity (the predicted size and age at which half the individuals are expected to be sexually mature). Bruger found ripe female bonefish throughout the year in waters off the Florida Keys and concluded that reproduction was not seasonal, but his sample size was small ($n=148$) and his conclusions equivocal. In other areas, bonefish reproduction appears to be seasonal according to patterns of larval and juvenile abundance (Alexander, 1961; Pfeiler, 1984; Pfeiler et al., 1988; Mojica et al., 1995). There

is also no published information on bonefish fecundity. In this article, we estimate the age and length at which sexual maturity is attained and describe the seasonal cycle of gonadal development in bonefish from waters off the Florida Keys. We also estimate the total fecundity of 33 bonefish collected from these waters.

Methods

Sampling

We examined 528 bonefish collected from South Florida waters from February 1989 to April 1995. Most of these bonefish were caught with hook-and-line gear either by biologists or by a single professional bonefish guide and his anglers from waters off the Florida Keys and in Florida and Biscayne Bays. Five bonefish caught with hook-and-line gear were obtained from taxidermists in Fort Lauderdale and five others from tournaments in the Keys. Supplemental collections of small bonefish (<425 mm) were made with various-size seines and gill nets in waters off the Keys. Ages, based on validated sectioned otoliths and growth rates of these bonefish, were described by Crabtree et al. (1996).

Fork length (FL) was measured to the nearest millimeter (mm), and fish were weighed to the nearest gram. Sex, gonad condition, and gonad weight (g) were recorded. Gonad samples for histology and for estimation of fecundity were removed from the fish and preserved in 10% buffered formalin; they were later soaked in water for one hour and then stored in 70% ethanol. Histological sections of gonads from 437 bonefish ranging from 228 to 702 mm were prepared and assessed for reproductive state. Gonad samples were processed histologically with a modification of the periodic acid Schiff's (PAS) stain for glycol-methacrylate sections, with Weigert's iron-hematoxylin as a nuclear stain and metanil yellow as a counterstain (Quintero-Hunter et al., 1991).

Oocyte staging

Oocytes were staged and counted from histological preparations at 100 \times with a compound microscope attached to a digital image-processing system. Three oocyte stages were recognized in bonefish ovaries: primary growth, cortical alveolar, and vitellogenic (Wallace and Selman, 1981; Fig. 1A). In addition, we counted PAS-positive melanomacrophage centers (Ravaglia and Maggese, 1995), which were present in many ovaries (Fig. 1B). When stained with the




Figure 1

(A) A histological section from an ovary of a 677-mm-FL bonefish, *Albula vulpes*, showing oocyte stages. PG = primary growth oocytes, CA = cortical alveolar oocytes, and V = vitellogenic oocytes. Scale bar = 400 microns. (B) A histological section showing PAS-positive melanomacrophage centers (PAS), cortical alveolar oocytes, and primary growth stage oocytes in a regressed ovary from a 692-mm-FL bonefish. When stained with periodic acid Schiff's stain, melanomacrophage centers are brilliant purple. Scale bar = 100 microns. (C) A histological section showing PAS-positive melanomacrophage centers in a regressed testis from a 586-mm-FL bonefish. Scale bar = 50 microns.

PAS stain, these structures are brilliant purple. Melanomacrophage centers are thought to be active in degrading atretic oocytes, postovulatory follicles, and residual cells of the spermatogenic cycle (Chan et al., 1967; Ravaglia and Maggese, 1995). At least 300 combined oocytes and melanomacrophage centers per slide were staged and counted in arbitrarily chosen fields, and frequencies were expressed as a percentage of the total count. We counted all structures that had at least 50% of their area visible in a field before moving to the next field. The presence of atretic hydrated oocytes was also noted.

Length and age at sexual maturity

Females were considered sexually mature if vitellogenic oocytes were present or if the histological sections appeared disorganized, highly vascularized, and contained widespread evidence of atresia. Documentation of atresia followed the classification of Hunter and Macewicz (1985). Immature females had small (gonadosomatic index <0.35), well-organized gonads that contained little evidence of atresia. We interpreted the widespread occurrence of PAS-positive melanomacrophage centers in inactive ovaries as evidence of past gonadal development, and we considered bonefish that had regressed (no vitellogenic oocytes present) ovaries containing many of these structures to be sexually mature (Fig. 1B). Males were considered sexually mature if the testes contained evidence of ongoing spermatogenesis, residual sperm, or widespread PAS-positive melanomacrophage centers associated with gonadal recrudescence (Fig. 1C).

Sometimes distinguishing between the gonads of sexually immature bonefish and the regressed gonads of mature fish was difficult. We reduced the probability of misclassifying regressed and immature fish by eliminating all bonefish collected during June–October from our analyses of age and length at sexual maturity. June–October was the season of minimal gonad development in bonefish, and most of the regressed bonefish in our sample were captured during this period. By excluding the postreproductive months from our analysis, we eliminated

84% of the regressed females and 63% of the regressed males in our sample. Hunter et al. (1992) recommended that only fish collected early in the spawning season be used to estimate the length at 50% maturity, but our sample size was not large enough to allow us to restrict our analysis to this extent.

To describe age and length at sexual maturity, we used nonlinear regression procedures to determine the inflection point of a logistic function fitted to the percentage of males and females that were sexually mature and to their respective lengths and ages. Parameter b in Table 1 is the inflection point and is the estimate of length or age at 50% maturity. Likelihood-ratio tests were used to compare the overall regression models and parameter estimates for males and females (Kimura, 1980).

Seasonality of gonad development

Monthly median gonadosomatic indices (GSI) of sexually mature males and females were plotted to show seasonal reproductive patterns. GSI's were calculated for 449 bonefish ranging from 228 to 702 mm as

$$GSI = (GW/(TW - GW)) \times 100,$$

where GW = total gonad weight (g); and
 TW = total fish weight (g).

We also plotted the monthly frequency of occurrence of the various oocytes stages that we counted for fish

Table 1

Percentage mature-age, percentage mature fork length, and weight-fecundity regressions for bonefish, *Albula vulpes*, from the waters of the Florida Keys. Wt = weight (g), FL = fork length (mm), AGE = age in years, FEC = fecundity. Values in parentheses are standard errors.

Y	X	n	a (1 SE)	b (1 SE)	r^2	Range of X for regressions
$Y = \left(1 / (1 + e^{(-a(X-b)})}\right) \times 100$						
% Mature (Females)	FL	150	0.028 (0.0064)	487.6 (8.14)	0.632	228–702
% Mature (Females)	AGE	143	1.122 (0.2345)	4.24 (0.192)	0.445	1–19
% Mature (Males)	FL	116	0.545 (2.8227)	417.5 (12.59)	0.735	322–687
% Mature (Males)	AGE	109	1.618 (0.3674)	3.60 (0.156)	0.464	2–19
$Y = a + bX$						
\log_{10} FEC	\log_{10} Wt	33	1.936 (0.4708)	1.131 (0.1312)	0.706	1,790–5,790

collected in the years during which we had regular monthly collections.

Fecundity

The total fecundity (the standing stock of advanced yolked oocytes) of 33 bonefish was estimated gravimetrically. Ovaries were subsampled from anterior, middle, and posterior portions of each ovary to evaluate spatial variations in oocyte size within the ovary and between ovaries. Subsamples of ovary containing 1,000–1,500 vitellogenic oocytes were weighed to the nearest 0.01 mg, and total fecundity was calculated on the basis of the mean number of oocytes per gram of ovary. Ovaries that contained widespread atresia, which suggested that partial spawning might have occurred, were not used for fecundity estimation.

Results

Two of the bonefish that we examined were statistically significant outliers (Crabtree et al., 1996); both were exceptionally small for their estimated ages and the weights of their otoliths were exceptionally light. Crabtree et al. excluded both fish from growth models, age-frequency distributions, and otolith weight-age regressions, and we also excluded them from our analyses. One was a 351-mm female that was 7 years old and the other was a 458-mm female that was 18 years old. Both fish were caught with hook-and-line gear on the ocean (Florida Straits) side of North Key Largo, and they were the smallest females examined whose ovaries contained vitellogenic oocytes. The 458-mm female had oocytes that were in the nuclear migratory stage, and these were the most advanced oocytes we found in any bonefish.

Length and age at sexual maturity

Male bonefish reached 50% sexual maturity at a length of 418 mm (95% confidence interval 393–443 mm) and an age of 3.6 years (95% confidence interval 3.3–3.9 years); females reached 50% sexual maturity at a length of 488 mm (95% confidence interval 472–504 mm) and an age of 4.2 years (95% confidence interval 3.9–4.6 years; Fig. 2; Table 1). The lengths at 50% maturity ($\chi^2=124.43$, $df=1$, $P<0.001$) and the ages at 50% maturity ($\chi^2=5.59$, $df=1$, $P=0.018$) for males and females were significantly different. In addition, the overall logistic equations for length at 50% maturity ($\chi^2=51.18$, $df=2$, $P<0.001$) and for age at 50% maturity ($\chi^2=11.55$, $df=2$, $P=0.003$) for males and females were significantly different. The smallest sexually mature male was 425 mm long, and the smallest sexually mature female was 358


mm long. The youngest sexually mature male was 3 years old, and the youngest sexually mature female was 2 years old. All males longer than 477 mm and all females longer than 594 mm were sexually mature. All males older than 5 years and all females older than 7 years were sexually mature.

Primary growth stage oocytes were present in all ovaries in which we counted oocytes (Fig. 3A). Cortical alveolar oocytes were present only in ovaries from fish longer than about 400 mm and older than 2 years and were common only in fish longer than about 475 mm and older than 4 years (Fig. 3B). Vitellogenic oocytes were found only in fish longer than 450 mm and were common only in fish longer than 550 mm and older than 5 years (Fig. 3C). PAS-positive melanomacrophage centers were common only in females longer than about 550 mm and older than 5 years (Fig. 3D): the same length and age as those for females that contained vitellogenic oocytes.

Seasonality of gonad development

Bonefish gonadal activity was seasonal. Vitellogenic oocytes were present in greatest numbers during November–May, and their numbers declined during May–June (Fig. 4). There were no vitellogenic oocytes in any ovaries from females collected during August–September of any of the three summers during which we sampled. Cortical alveolar oocytes were present during all months but were least abundant during July–October. Primary growth stage oocytes were present in all females examined and made up at least 20% of the total number of oocytes present. PAS-positive melanomacrophage centers were most abundant in the gonads of spent and regressed males and females and were most abundant in ovaries at the end of the spawning season in June–August (Fig. 5). They were least abundant in ovaries immediately before the initiation of spawning in November, when recrudescence was complete and most ovaries were ripening to spawn during winter–spring. We saw no evidence of recent or imminent spawning, such as postovulatory follicles or fully hydrated females. Only six ovaries contained atretic hydrated oocytes, and no single histological preparation contained more than a few hydrated oocytes.

Seasonal GSI patterns suggest that bonefish spawned during a prolonged period from November to June (Fig. 6). Median GSI's were greatest during November–May and were least during July–September. The decrease in female GSI's during July–September corresponded with the decrease in the number of vitellogenic oocytes present in ovaries and with the increased abundance of spent and regressed fish during late summer.



Fecundity

Bonefish total fecundity estimates ranged from 0.4 to 1.7 million oocytes and had a significant positive relation to fish weight (Table 1; Fig. 7). Relative fecundity (the number of oocytes per gram fish weight) ranged from 159 to 385 oocytes/g (mean=259 oocytes/g, SD=47.1, $n=33$) for fish ranging in length from 485 to 702 mm. There was no significant relation between relative fecundity and fish length ($n=33$, $r^2=0.014$, $P=0.514$) or weight ($n=33$, $r^2=0.043$, $P=0.247$), but there was a significant positive relation between relative fecundity and age ($n=32$, $r^2=0.248$, $P=0.003$).


Oocyte development among areas within the ovary was homogeneous. We used a two-factor analysis of variance to compare oocyte densities with side (right or left) and position (anterior, middle, or posterior section of the ovary) as the effects. The number of late vitellogenic oocytes per gram of wet ovary weight was not significantly different between left and right ovaries (ANOVA, $df=1$, $P=0.648$) or among subsamples from anterior, middle, or posterior sections of the ovary (ANOVA, $df=2$, $P=0.709$). Furthermore,

we found no significant interaction between side and position from which subsamples were taken (ANOVA, $df=2$, $P=0.702$). Weights of left and right ovaries from sexually mature females were not significantly different (paired t -test, $n=112$, $t=0.480$, $P=0.632$), but right testes from sexually mature males were significantly larger than left testes (mean difference=9.1 g, SD=18.62; paired t -test, $n=98$, $t=4.826$, $P<0.001$).

Discussion

Length and age at sexual maturity

The bonefish we examined reached sexual maturity at an older age and larger size than reported by Bruger (1974). He reported sexually mature females that were 1 year old and ranged from 221 to 352 mm FL (reported as 210 to 338 mm standard length). Bruger considered these small bonefish to be sexually mature on the basis of the presence of vitellogenic oocytes, one of the criteria that we used. He did not report how many of these small sexually mature fe-



males were captured or the typical length and age at sexual maturity. Some of the sexually mature females he collected were smaller than the 351-mm sexually mature female that we considered an outlier and excluded from our analysis. Furthermore, the lengths of Bruger's fish were substantially shorter than our estimated length at 50% maturity for females (488 mm). We also did not find any 1-year-old bonefish that were sexually mature; the youngest sexually mature fish we examined was 2 years old. All of the small mature females reported by Bruger were caught in deeper water (9.1–12.2 m) than that surveyed for bonefish in our sample; most of our fish

were caught in water less than 2 m deep. Both Bruger (1974) and Crabtree et al. (1996) considered the possible existence of a cryptic bonefish species in waters off the Florida Keys as a potential explanation for the presence of exceptionally small and sexually mature bonefish, but additional study is needed to resolve this question.

Little is known regarding bonefish maturation in other areas. Pfeiler et al. (1988) reported 12 *Albula* sp., ranging in length from 205 to 264 mm (SL) from the Gulf of California, that had ripe or ripening gonads; this finding suggests a smaller length at sexual maturity there than we found in the Keys. The *Albula*

species in the Gulf of California is distinct from the Caribbean *A. vulpes* (Pfeiler, 1996).

Most of the bonefish in our sample that were caught with hook-and-line gear were 500–700 mm long (80%) and 3–10 years old (86%; Crabtree et al., 1996). If the length and age composition of our hook-

and-line sample reflects that of the fishery, as suggested by Crabtree et al. (1996), then most of the fish caught in the fishery are longer than our estimate of length at 50% maturity and older than our estimate of age at 50% maturity. The current 390-mm-FL (457-mm-TL) legal minimum fish length imposed upon the fishery is less than our estimate of the length at 50% maturity for males (418 mm FL) and females (488 mm FL). Crabtree et al. (1996) found little evidence of fishing mortality in the Florida Keys because most bonefish caught by recreational anglers are released; thus, the population would probably be insensitive to changes in the legal minimum fish length.

The presence of PAS-positive melanomacrophage centers was related to gonadal activity. They appeared to be associated with the resorption of postovulatory follicles and atretic vitellogenic oocytes, as has been suggested by Ravaglia and Maggese (1995). The presence of vitellogenic oocytes is an unambiguous indication of sexual maturity, and the length and age at which vitellogenic oocytes first developed in bonefish corresponded to the length and age at which PAS-positive melanomacrophage centers first appeared (Fig. 3). Melanomacrophage centers were most abundant at the end of the spawning season, when spent fish were most common, and decreased in abundance during

the postreproductive period (July–November), when recrudescence occurred (Fig. 5). Ravaglia and Maggese (1995) found that the abundance of melanomacrophage centers was also seasonal in *Synbranchus marmoratus* and peaked during the postreproductive season.

Seasonality of gonad development

Bonefish gonadal activity in the Florida Keys was seasonal, and development occurred over about eight months, from November to June. Bonefish were reproductively inactive for only a few months during summer. This period of inactivity roughly corresponds with the period of maximum water temperatures in the Keys (Crabtree et al., 1996). Bruger (1974) reported no evidence of a seasonal pattern of gonadal development for bonefish off the Florida Keys, but his small sample size during July–September ($n=7$ females, $n=4$ males) may have obscured seasonal patterns. Alexander (1961)




Figure 4

Monthly mean percent frequency of occurrence and standard errors of oocyte stages in bonefish, *Albula vulpes*, ovaries ($n=250$) for 1992–94. * = primary growth stage oocytes, \diamond = cortical alveolar oocytes, and \square = vitellogenic oocytes.




Figure 5

Monthly mean percent frequency of occurrence and standard errors of PAS-positive melanomacrophage centers in bonefish, *Albula vulpes*, ovaries. The numbers above the error bars are the numbers of ovaries examined.

found that the abundance of pre-metamorphic bonefish larvae in the West Indies was seasonal and that most larvae (96% of a total collection of 417 premetamorphic larvae) were caught during November–April. She concluded, from the capture of 17 premetamorphic larvae during August, that bonefish spawn throughout the year in the West Indies, but she suggested that bonefish spawn at reduced levels during the hot summer months. The 17 larvae that she reported were captured in August and were caught far south of the Florida Keys—at 13°11'N latitude, where spawning may follow a different seasonal pattern than it does in the Keys.

Studies from other areas also suggest seasonal reproduction in bonefish. Recruitment studies off the Bahama Islands suggest that spawning there has a seasonal pattern that is similar to that of spawning in the Florida Keys. Mojica et al. (1995) backcalculated hatching dates from the analysis of otolith microstructure of field-collected larvae and found that bonefish spawned continuously from mid-October through early January. Recruitment patterns suggest that spawning probably continues until spring, because Mojica et al. (1995) observed recruitment pulses of bonefish larvae as late as June—presumably resulting from April and May spawning. Bonefish spawning in the Gulf of California also appears to be seasonal. Pfeiler et al. (1988) examined gonads from 33 bonefish ranging from 202 to 279 mm (SL) and suggested that *Albula* sp. in the Gulf of California spawn during late spring and early summer. Metamorphic leptocephali are abundant in coastal regions and hypersaline lagoons throughout the Gulf of California during winter and spring (Pfeiler, 1984; Pfeiler et al., 1988). Pfeiler et al. (1988) suggested that the premetamorphic larval phase lasts 6 or 7 months.

The location of bonefish spawning grounds remains unknown, but the absence of females with post-ovulatory follicles or many hydrated oocytes in our samples suggests that bonefish do not spawn in the




Figure 6
Monthly median gonadosomatic indices (GSI) and interquartile ranges for sexually mature male and female bonefish, *Albula vulpes*.




Figure 7
The fecundity-weight (g) relation for bonefish, *Albula vulpes*. The fecundity-weight regression equation is presented in Table 1.

shallow nearshore areas where the fishery exists. It is possible that bonefish had spawned in our sampling area but did not feed before or after spawning and so were not available for capture with hook-and-

line gear until after they had completed the resorption of recognizable postovulatory follicles. This possibility seems unlikely because most collections of premetamorphic bonefish larvae are from offshore waters (Alexander, 1961); thus spawning bonefish probably move out of the shallow waters (<2 m) where fishing usually occurs. Alexander (1961) suggested that bonefish either spawn offshore or in areas where currents are likely to carry the eggs offshore.

Fecundity

We did not examine any bonefish ovaries containing oocytes in the final stages of oocyte maturation or showing definitive evidence of recent spawning, such as postovulatory follicles. Consequently, we do not know if bonefish are isochronal or multiple-batch spawners, and we could not estimate batch fecundity. If annual fecundity in bonefish is indeterminate (Hunter et al., 1992), our estimate of total fecundity may not accurately represent total annual egg production. The bonefish spawning season is prolonged, and the potential exists for additional vitellogenic oocytes to mature from the standing stock of unyolked oocytes during the spawning season. Some ovaries contained vitellogenic oocytes, widespread atresia, and were loosely organized and highly vascularized. These females may have spawned earlier in the season and were developing an additional batch of oocytes that would have been spawned later in the season. It is unclear whether these oocytes were recruited from the standing stock of unyolked oocytes after previous spawning or if they were vitellogenic oocytes already present in the ovary that did not ovulate during previous spawning. Another bias of our fecundity estimates is that we could not correct them for atretic losses of vitellogenic oocytes during the spawning season; these losses could have caused us to overestimate egg production.

Acknowledgments

We thank Captain John Kipp, who provided us with most of the bonefish examined in this study and without whose efforts this work would not have been possible. We also thank Captain Mike Collins, the Florida Keys Fishing Guides Association, and the Islamorada bonefish tournaments for their support. We also thank John Swanson, Bill Gibbs, and the staff at the Keys Marine Laboratory for their assistance; Jim Colvocoresses, John Hunt, and others at the South Florida Regional Laboratory for their cooperation; and David Harshany, Victor Neugebauer, and Connie Stevens for their assistance. Jim Colvocoresses,

Harry Grier, Judy Leiby, Rich McBride, Jim Quinn, and Dana Winkelman made helpful comments that improved the manuscript. This work was supported in part under funding from the Department of the Interior, U.S. Fish and Wildlife Service, Federal Aid for Sportfish Restoration Project Number F-59.

Literature cited

- Alexander, E. C.**
1961. A contribution to the life history, biology and geographical distribution of the bonefish, *Albula vulpes* (Linnaeus). Dana-Rep. Carlsberg Found. 53, 51 p.
- Böhlke, J. E., and C. C. G. Chaplin.**
1993. Fishes of the Bahamas and adjacent tropical waters, 2nd ed. Univ. Texas Press, Austin, TX, 771 p.
- Bruger, G. E.**
1974. Age, growth, food habits, and reproduction of bonefish, *Albula vulpes*, in South Florida waters. Fla. Mar. Res. Publ. 3, 20 p.
- Chan, S. T. H., A. Wright, and J. G. Phillips.**
1967. The atretic structures in the gonads of the rice-field eel (*Monopterus albus*) during natural sex-reversal. J. Zool. (Lond.) 153:527-539.
- Crabtree, R. E., C. W. Harnden, D. Snodgrass, and C. Stevens.**
1996. Age, growth, and mortality of bonefish, *Albula vulpes*, from the waters of the Florida Keys. Fish. Bull. 94: 442-451.
- Hildebrand, S. F.**
1963. Family Albulidae. In H. B. Bigelow (ed.), Fishes of the western North Atlantic, p. 132-147. Mem. Sears Found. Mar. Res. Yale Univ. 1(3).
- Hunter, J. R., and B. J. Macewicz.**
1985. Measurement of spawning frequency in multiple spawning fishes. In R. Lasker, (ed.), An egg production method for estimating spawning biomass of pelagic fish: application to the northern anchovy, *Engraulis mordax*, p. 79-94. U.S. Dep. Commer., NOAA Tech. Rep. NMFS 36.
- Hunter, J. R., B. J. Macewicz, N. C. Lo, and C. A. Kimbrell.**
1992. Fecundity, spawning, and maturity of female Dover sole *Microstomus pacificus*, with an evaluation of assumptions and precision. Fish. Bull. 90:101-128.
- Kimura, D. K.**
1980. Likelihood methods for the von Bertalanffy growth curve. Fish. Bull. 77:765-776.
- Mojica, R., J. M. Shenker, C. W. Harnden, and D. E. Wagner.**
1995. Recruitment of bonefish, *Albula vulpes*, around Lee Stocking Island, Bahamas. Fish. Bull. 93:666-674.
- Pfeiler, E.**
1984. Inshore migration, seasonal distribution and sizes of larval bonefish, *Albula*, in the Gulf of California. Environ. Biol. Fishes 10:117-122.
1996. Allozyme differences in Caribbean and Gulf of California populations of bonefishes (*Albula*). Copeia 1996:181-183.
- Pfeiler, E., M. A. Mendoza, and F. A. Manrique.**
1988. Premetamorphic bonefish (*Albula* sp.) leptocephali from the Gulf of California with comments on life history. Environ. Biol. Fishes 21:241-249.
- Quintero-Hunter, I., H. Grier, and M. Muscato.**
1991. Enhancement of histological detail using metanil yellow as counterstain in periodic acid Schiff's hematoxy-

- lin staining of glycol methacrylate tissue sections. *Bio-technol. Histochem.* 66:169–172.
- Ravaglia, M. A., and M. C. Maggese.**
1995. Melano-macrophage centers in the gonads of the swamp eel, *Synbranchus marmoratus* Bloch, (Pisces, Synbranchidae): histological and histochemical characterization. *J. Fish Dis.* 18:117–125.
- Rivas, L. R., and S. M. Warlen.**
1967. Systematics and biology of the bonefish, *Albula nemoptera* (Fowler). *Fish Bull.* 66:251–258.
- Robins, C. R., and G. C. Ray.**
1986. A field guide to Atlantic coast fishes of North America. The Petersen field guide series, number 32. Houghton Mifflin Co., Boston, MA, 354 p.
- Uyeno, T., K. Matsuura, and E. Fujii.**
1983. Fishes trawled off Suriname and French Guiana. Japan Marine Fishery Resource Research Center, 519 p.
- Wallace, R. A., and K. Selman.**
1981. Cellular and dynamic aspects of oocyte growth in teleosts. *Am. Zool.* 21: 325–343.
- Whitehead, P. J. P.**
1986. The synonymy of *Albula vulpes* (Linnaeus, 1758) (Teleostei, Albulidae). *Cybium* 10:211–230.